PAUS
Pet Animal Ultrasound Service
4paus.com 781-635-5677



SonoPath
Clinical Sonography & Telecytology
EDUCATIONAL TELECONSULTATION SERVICES™
1-800-838-4268 info@sonopath.com SonoPath.com

PATIENT
Avery Kallstrom

SPECIES
Feline

BREED
DLH

SEX
Female Spayed

AGE
11 years

WEIGHT
10.5lbs

INTERPRETED BY
Maggie Machen Lamy, DVM DACVIM (Cardiology)

IMAGING PERFORMED BY
Pamela Harrigan, RDCS

HOSPITAL NAME
Halifax Veterinary Hospital

REFERRING VET
Dr. Hopkins

INVOICE
27072

DATE
10/24/22

PRESENTING CLINICAL SIGNS

History: Grade III/VI heart murmur. No clinical signs Echocardiogram prior to anesthesia for dental prophy. Normal thyroid values. BP: 140mmHg.

ECHOCARDIOGRAM FINDINGS

2D, m-mode, color flow and Doppler imaging is available.
Left ventricle: The LV diameter is normal with adequate myocardial function. The LV wall thicknesses are asymmetric with severe septal thickening. There is a diffusely hyperechoic endocardium consistent with fibrosis. The papillary muscles are mildly hypertrophied. False tendon.
Left atrium: The left atrium is normal. No obvious spontaneous contrast or thrombi seen.
Mitral valve: The mitral valve is normal in structure and mobility. No obvious systolic anterior motion is seen. No MR.
Aortic valve/Aorta: The aortic valve is normal in morphology and mobility. Normal aortic outflow velocity; laminar flow. No aortic insufficiency.
Right ventricle: Normal right ventricular diameter and morphology indicating no overt evidence of pulmonary arterial hypertension.
Right atrium: The right atrium is normal in dimension.
Tricuspid valve: The tricuspid valve appears normal with no tricuspid regurgitation.
Pulmonic valve/Pulmonary artery: The pulmonic valve is normal in morphology and mobility. No pulmonic insufficiency. Pulmonic outflow is normal on Spectral doppler; however, a dynamic outflow obstruction is suspected on 2D/color flow imaging.
Pericardium/other: No pericardial or pleural effusion noted. No obvious cardiac masses.
Heart rhythm: ECG reveals a sinus rhythm with an average HR of 200bpm.

2-Dimensional Measurements

Ao diam (cm)	0.9
LA diam (cm)	1.1
LA:Ao (Swe)	1.2
IVS thickness (cm)	0.77
LVID diastole (cm)	1.2
PW thickness (cm)	0.44
LVID systole (cm)	0.4
FS (%)	70

Doppler Measurements

PV Vmax (m/s)	1.5
AoV Vmax (m/s)	1.5
MR Vmax (m/s)	NA
TR Vmax (m/s)	NA
TR PG (mmHg)	NA

INTERPRETATION OF THE FINDINGS

HCM is a rule out diagnosis, once hypertension and hyperthyroid disease are ruled out. Both have been considered in this case, making primary disease likely. Regardless, the degree of disease is mild, with significant yet isolated septal hypertrophy and no LA dilation. The murmur is benign in origin due to a dynamic RVOT obstruction. No additional issues are identified.

Prognosis is guarded, due to the highly variable rates of progression with subclinical feline cardiomyopathy.

RECOMMENDATIONS

- Given these findings, no medications are indicated.
- Monitor BP and T4 every 6 months.



PATIENT
Avery Kallstrom

SPECIES
Feline

BREED
DLH

SEX
Female Spayed

AGE
11 years

WEIGHT
10.5lbs

INTERPRETED BY
Maggie Machen Lamy, DVM DACVIM (Cardiology)

IMAGING PERFORMED BY
Pamela Harrigan, RDCS

HOSPITAL NAME
Halifax Veterinary Hospital

REFERRING VET
Dr. Hopkins

INVOICE
27072

DATE
10/24/22

- Anesthetic risk is considered mild, however judicious IV fluid rates are advised to avoid fluid overload. Additionally, drugs that stimulate heart rate should be avoided unless clinically necessary (glycopyrrolate, atropine). A reasonable protocol includes opioid/benzodiazepine premedication, propofol induction, isoflurane maintenance.
- Risk for complication with steroid use typically follows LA dilation, which in this case is low. That being said, any cat can experience unexpected signs of intolerance and monitoring of RR/RE is advised particularly in the initiation phase.
- Monitor for any clinical evidence of cardiac compromise, including respiratory changes and/or signs of a blood clot event (paralysis, neurologic changes, etc.).

PLAN

- Recommend recheck echocardiogram in 6-12 months to screen for progression, sooner if any clinical signs arise in the interim.

IMAGES



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Maggie Machen Lamy, DVM
Diplomate of the American College of Veterinary Internal Medicine (Cardiology)
info@sonopath.com