


PATIENT

Bentley Johnston

PRESENTING CLINICAL SIGNS

 History: Coughing, shallow rapid breathing. Liver enzymes elevated.
 -Current medications: Furosemide, spironolactone.
 -Abnormal PE/Chem/CBC/UA Results: Elevated creatinine, severely elevated ALT.

SPECIES

Canine

ECHOCARDIOGRAM FINDINGS

2D, m-mode, color flow and doppler imaging is available. Severe left ventricular dilation with diminished systolic function. Severe left atrial enlargement. The mitral valve appears thickened with lack of coaptation in systole. Mild to moderate central mitral regurgitation secondary to annular stretch. Decreased MR velocity. Decreased LV wall thickness. The tricuspid valve appears normal in form and function. Moderate right atrial and ventricular dilation. Mild tricuspid regurgitation. Borderline TR velocity. The aortic valve is normal in morphology and mobility. No subvalvular ridge present; normal LVOT velocity. No aortic insufficiency. Normal pulmonic valve with no pulmonic insufficiency seen. No pericardial or pleural effusion noted. No obvious cardiac tumors. Rapid irregular rhythm throughout.

BREED

Great Dane

SEX

Male Neutered

AGE

4 years

CARDIAC CHART
WEIGHT

159lbs

CANINE CARDIAC PARAMETERS	MR VMAX (m/s)	TR VMAX (m/s)	LA/AO (Boon method)	LA/AO (Heart Base; Swe)	FS (%)	EF (%)	EPSS (cm)
NORMAL PARAMETER	4.5-5.5	<2.7	1.3	<1.6	28-40	40-100	<0.6
PATIENT	4.0	2.7	NM	2.0	5	7	2.5
CANINE CARDIAC PARAMETERS	HR (BPM)	AV VMAX (m/s)	PV MAX (m/s)	BODY WEIGHT (kg)	LA 2D short axis Base view (cm)	LVIDd Avg; 2D and m-mode short axis (cm)	LVIDs Avg; 2D and m-mode short axis (cm)
NORMAL PARAMETER	50-100	0.7-1.7	0.7-1.6	BELOW	BELOW	BELOW	BELOW
PATIENT	220	1.1	0.9	72.1	4.6	7.2	6.8
*Normal chamber parameters expressed as a mean value (SD)				3	1.27 (5.3)	2.46 (2.46)	1.36 (5.5)
BODY WEIGHT DEPENDENT PARAMETERS				5	1.40 (4.5)	2.74 (5.2)	1.60 (4.7)
<i>*Note: All measurements based upon multi-modal images and methods. An average value is reported.</i>				10	1.50 (3.8)	3.27 (3.5)	2.06 (3.1)
				15	1.83 (2.0)	3.71 (2.4)	2.43 (2.1)
				20	2.02 (1.9)	4.14 (2.2)	2.80 (2.0)
				25	2.18 (2.4)	4.48 (2.9)	3.10 (2.5)
				30	2.33 (3.3)	4.83 (3.9)	3.39 (3.4)
				35	2.48 (4.3)	5.17 (5.0)	3.69 (4.5)
				40	2.62 (5.2)	5.48 (6.1)	3.96 (5.4)
				50	2.88 (7.1)	6.07 (8.3)	4.46 (7.4)

INTERPRETED BY

 Maggie Machen Lamy,
 DVM DACVIM
 (Cardiology)

IMAGING PERFORMED BY

Kelly Reschny, RVT

HOSPITAL NAME

 St Catharines Animal
 Hospital

REFERRING VET

Dr. Masoud

INVOICE


22954

DATE

3/7/22

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Unfortunately, this patient has severe biventricular myocardial failure. There is dilation and overload of all 4 chambers resulting in insufficiency of the mitral and tricuspid valves. The degree of dilation and pump failure is resulting in a rapid irregular tachycardia (suspect AF; intermittent VT cannot be ruled out) and congestive heart failure is highly suspected. The arrhythmia puts the patient at risk for right heart dilation and congestion (causing tachycardia-induced cardiomyopathy typically leads to right-sided congestive



PATIENT

Bentley Johnston

signs), while respiratory distress is typically due to left-sided congestion (pulmonary edema).

SPECIES

Canine

The importance of a screening ECG cannot be stressed enough in order to definitively diagnose the arrhythmia. Without treatment we will not be able to get the congestion controlled despite diuretic therapy. If atrial fibrillation (AF) is confirmed, rate control must be instituted in order to get the rate and secondary congestion under control (typically using Diltiazem +/- Digoxin). If ventricular in origin, sotalol or mexiletine may be necessary. If this is not possible, referral to a tertiary facility should be offered for further evaluation ASAP.

BREED

Great Dane

SEX

Male Neutered

Systolic failure can be primary in nature (DCM) or secondary to taurine deficiency, myocarditis, hypothyroidism, tachycardia-induced cardiomyopathy, or infiltrative disease such as lymphoma. While primary disease is certainly suspected in this breed, consider testing for primary causes that may be treatable. A troponin (cTnI) level can be submitted to further investigate infiltrative/inflammatory contribution (myocarditis). Additionally, a taurine level may be helpful (screen for malabsorption issue), and a thorough diet history given the recent correlation with grain free/boutique brand/exotic ingredient diets. Finally, further systemic evaluation for underlying infiltrative contribution such as neoplasia is also reasonable (abdominal ultrasound, etc.).

AGE

4 years

WEIGHT

159lbs

Regardless of cause, prognosis is poor at this stage in the disease process, with an average survival time of <6mo for canine patients with active congestion on medications, however they generally are able to maintain a good quality of life for that period. Even with diet-related dysfunction, improvement will likely be minimal at this end-stage phase of disease. Immediate initiation of full cardiac supportive medications is recommended as below. Emergency hospitalization for supportive care, O2 support and potentially IV rate control is recommended. Cases of systolic failure are at high risk for malignant tachyarrhythmias (such as AF or VT), and activity restriction is advised. Patient will always be at risk for recurrent CHF, development of arrhythmias/LA tear, syncope and/or sudden death in the future.

INTERPRETED BY

Maggie Machen Lamy,
DVM DACVIM
(Cardiology)

IMAGING PERFORMED BY

Kelly Reschny, RVT

Monitor for development of a cough, worsening labored breathing, exercise intolerance or collapse episodes in the future. Monitoring of sleeping breathing rates at home is recommended to assess response to medications and recurrence of CHF in the future.

HOSPITAL NAME

St Catharines Animal
Hospital

PLAN

An ECG should be performed ASAP due to the rapid irregular heart rate noted during the study in order to dictate anti-arrhythmic therapy. Consider hospitalization for IV diuretic/rate control therapy if needed/indicated. Initiate diuretic furosemide 1-2mg/kg PO q8h if unstable; wean to q12h once breathing normally (2-3 days). Initiate aldosterone antagonist Spironolactone 1-2mg/kg PO q12h. Initiate Pimobendan 0.3mg/kg PO q12h.

REFERRING VET

Dr. Masoud

Once HR is controlled, a recheck renal panel, heart rate/BP is recommended in 5-7 days. If doing well and BP >130mmHg at this time, reinstitute ACEI 0.5mg/kg PO q12h.

INVOICE

22954

Recheck echocardiogram and ECG in 4-6 months to reassess cardiac function.

DATE

3/7/22



PATIENT

Bentley Johnston

SPECIES

Canine

BREED

Great Dane

SEX

Male Neutered

AGE

4 years

WEIGHT

159lbs

INTERPRETED BY

Maggie Machen Lamy,
DVM DACVIM
(Cardiology)

**IMAGING
PERFORMED BY**

Kelly Reschny, RVT

HOSPITAL NAME

St Catharines Animal
Hospital

REFERRING VET

Dr. Masoud


INVOICE

22954

DATE

3/7/22

IMAGES



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Maggie Machen Lamy, DVM
Diplomate of the American College of Veterinary Internal Medicine (Cardiology)
info@sonopath.com