



## PATIENT

Eladio Camilo  
Foundation Brownie  
Blondie

## PRESENTING CLINICAL SIGNS

History: Grade 3/6 heart murmur. Diagnosed with heartworm disease; assess prior to treatment.  
-Abnormal PE/Chem/CBC/UA Results: Mild elevation in BUN.  
-Radiographs: WNL.

## SPECIES

Canine

## BREED

Labrador Mix

## SEX

Male Intact

## AGE

Middle-aged

## WEIGHT

36.8lbs

## INTERPRETED BY

Maggie Machen Lamy,  
DVM, DACVIM  
(Cardiology)

## IMAGING PERFORMED BY

G. Ferrer, DVM

## HOSPITAL NAME

Paseos Veterinary  
Center

## REFERRING VET

Dr. Carrasquillo

## INVOICE

23577

## DATE

4/11/22

## ECHOCARDIOGRAM FINDINGS

2D, m-mode, color flow and doppler imaging is available. Mild thickening of mitral valve with no prolapse into the left atrial lumen. Mild to moderate MR. Minimal LA enlargement. Normal MR velocity. No LV dilation with adequate myocardial function. The MPA and branches are minimally dilated. Concern for adult worm(s) in the distal pulmonary artery near the bifurcation (see below); no worms seen in the RA or RV. Minimal right atrial dilation. RV appears normal with no obvious RVH. Mild tricuspid regurgitation. Normal TR velocity, suggesting normal pulmonary pressures. The pulmonic and aortic valves are normal in morphology and mobility. Normal pulmonic and aortic outflow velocities with laminar flow. No pulmonic or aortic insufficiency. No pericardial or pleural effusion. No cardiac tumors seen.

## CARDIAC CHART

CANINE CARDIAC PARAMETERS	MR VMAX (m/s)	TR VMAX (m/s)	LA/AO (Boon method)	LA/AO (Heart Base; Swe)	FS (%)	EF (%)	EPSS (cm)
NORMAL PARAMETER	4.5-5.5	<2.7	1.3	<1.6	28-40	40-100	<0.6
PATIENT	5.0	2.0	NM	1.3	56	87	0.53
CANINE CARDIAC PARAMETERS	HR (BPM)	AV VMAX (m/s)	PV MAX (m/s)	BODY WEIGHT (kg)	LA 2D short axis Base view (cm)	LVIDd Avg; 2D and m-mode short axis (cm)	LVIDs Avg; 2D and m-mode short axis (cm)
NORMAL PARAMETER	50-100	0.7-1.7	0.7-1.6	BELOW	BELOW	BELOW	BELOW
PATIENT	NM	0.7	0.6	16.7	2.9	4.0	1.5
*Normal chamber parameters expressed as a mean value (SD)				3	1.27 (5.3)	2.46 (2.46)	1.36 (5.5)
<b>BODY WEIGHT DEPENDENT PARAMETERS</b>				5	1.40 (4.5)	2.74 (5.2)	1.60 (4.7)
<i>*Note: All measurements based upon multi-modal images and methods. An average value is reported.</i>				10	1.50 (3.8)	3.27 (3.5)	2.06 (3.1)
				15	1.83 (2.0)	3.71 (2.4)	2.43 (2.1)
				20	2.02 (1.9)	4.14 (2.2)	2.80 (2.0)
				25	2.18 (2.4)	4.48 (2.9)	3.10 (2.5)
				30	2.33 (3.3)	4.83 (3.9)	3.39 (3.4)
				35	2.48 (4.3)	5.17 (5.0)	3.69 (4.5)
				40	2.62 (5.2)	5.48 (6.1)	3.96 (5.4)
				50	2.88 (7.1)	6.07 (8.3)	4.46 (7.4)

Adapted from June Boon, Veterinary Echocardiography, 1998  
Rishniw M and Hollis NE, J Vet Intern Med 2000; 14:429-435  
Hansson et al, Vet Rad and Ultrasound 2002  
Bonagura et al. Echocardiography: principles of interpretation, Vet Clin North Am 15:1177, 1995

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The murmur is due to chronic degenerative valve disease causing mild to moderate mitral and mild tricuspid regurgitation. The left atrium is minimally enlarged, indicating the current risk for complication is low. More importantly, there is suspicion for an adult heartworm infestation in the distal MPA. This is certainly not definitive in this peripheral location; however, as ultrasound is largely insensitive (i.e., adult worms may be easily either missed peripheral or elsewhere). Given a lack of right heart enlargement, the infestation is considered relatively mild without evidence significant pulmonary hypertension. No additional issues are identified.



**PATIENT**

Eladio Camilo  
Foundation Brownie  
Blondie

Heartworms can cause significant damage to the lung tissue leading to pulmonary damage, pulmonary hypertension and clinical signs such as coughing, decreased ability to exercise, or difficulty breathing. Disease severity can range from an asymptomatic dog with few worms to dogs with severe respiratory signs. In the most severe cases, caval syndrome may develop due to a very high worm burden sheering blood cells as they pass through the heart. Caval syndrome is a life-threatening emergency that requires immediate surgical removal of the worms.

**SPECIES**

Canine

Given that this patient has no reported clinical signs and no right heart enlargement, we do have some flexibility when approaching therapy. Medical management with drugs like Sildenafil and prednisone is typically utilized if the patient is showing respiratory signs or syncope (none noted). If extraction would be a possibility from a financial standpoint, I would consider referral to a local cardiologist for advanced echocardiography and evaluation. If that is not a possibility, utilizing the standard approach to heartworm treatment as dictated by the American Heartworm Society is recommended, including 30 days of doxycycline and heartguard prior to the split immiticide protocol. Please see website and protocol for specific information. There is high risk for thromboembolism in any patient, however those with adult worms seen in the PA are no question at elevated risk. At this time, exercise restriction is paramount, including cage rest with leash walks only, as a worm embolus can be a life-threatening complication of the disease. This should be continued for an additional 6-8 weeks following therapy.

**BREED**

Labrador Mix

**SEX**

Male Intact

**AGE**

Middle-aged

Modifications to this protocol are sometimes elected depending on individual circumstances which may involve fewer injections or a "slow kill" method. These are not, however, our standard recommendation as alternate treatment may not result in effective treatment of the infestation.

**WEIGHT**

36.8lbs

Following treatment, retest for heartworm disease 6 months after completing the full course of therapy. Anesthesia is NOT advised prior to completing the protocol, as vasodilation can lead to increased risk for an embolus. Prognosis is guarded, as the right heart/MPA changes are often permanent and may cause clinical signs (exertional syncope/dyspnea, right-sided CHF) in the future.

**INTERPRETED BY**

Maggie Machen Lamy,  
DVM, DACVIM  
(Cardiology)

During therapy, there is high risk for a worm embolus and breathing rate and effort should be monitored closely. Anti-inflammatory prednisone can be used if becomes symptomatic. Patient will be at high risk for developing clinical signs due to pulmonary hypertension with age given the inherent secondary inflammation and damage to the pulmonary vasculature and lungs, and periodic rechecks may be helpful. Monitor for exertional dyspnea or fainting episodes going forward.

**IMAGING PERFORMED BY**

G. Ferrer, DVM

**HOSPITAL NAME**

Paseos Veterinary  
Center

Once heartworm negative, a recheck echocardiogram and chest radiographs are recommended in 6 months to reassess right heart and screen for progression in CVD.

**REFERRING VET**

Dr. Carrasquillo

**INVOICE**

23577

**DATE**

4/11/22



**PATIENT**

Eladio Camilo  
Foundation Brownie  
Blondie

**SPECIES**

Canine

**BREED**

Labrador Mix

**SEX**

Male Intact


**AGE**

Middle-aged

**WEIGHT**

36.8lbs

**IMAGES**



**INTERPRETED BY**

Maggie Machen Lamy,  
DVM, DACVIM  
(Cardiology)

**IMAGING PERFORMED BY**

G. Ferrer, DVM

**HOSPITAL NAME**

Paseos Veterinary  
Center

**REFERRING VET**

Dr. Carrasquillo

**INVOICE**

23577

**DATE**

4/11/22

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Maggie Machen Lamy, DVM  
Diplomate of the American College of Veterinary Internal Medicine (Cardiology)  
info@sonopath.com