


PATIENT

Tango Taverner

PRESENTING CLINICAL SIGNS

History: Low grade heart murmur. No cough. 4Dx Test - Positive for heartworm antigen with Microfilaria. BP: 133mmHg.

-Current medications: Gabapentin 300mg: 1 capsule every 12 hours

SPECIES

Canine

ECHOCARDIOGRAM FINDINGS

2D, m-mode, color flow and doppler imaging is available. Mild thickening of mitral valve with no prolapse into the left atrial lumen. No MR. Normal LA dimension. No LV dilation with adequate myocardial function. The MPA is normal in dimension. Possible HW seen in the distal PA near the bifurcation (see below). Artifact cannot be ruled out. The peripheral branches are not visualized. No worms seen in the RA or RV. No right atrial dilation. RV appears normal with no obvious RVH. No tricuspid regurgitation. The pulmonic and aortic valves are normal in morphology and mobility. Normal pulmonic and mildly elevated aortic outflow velocities with laminar flow. No pulmonic or aortic insufficiency. No pericardial or pleural effusion.

BREED

American Bulldog

SEX

Male Neutered

AGE

8 years

CARDIAC CHART
WEIGHT

78.9lbs

INTERPRETED BY

 Maggie Machen Lamy,
 DVM DACVIM
 (Cardiology)

CANINE CARDIAC PARAMETERS	MR VMAX (m/s)	TR VMAX (m/s)	LA/AO (Boon method)	LA/AO (Heart Base; Swe)	FS (%)	EF (%)	EPSS (cm)
NORMAL PARAMETER	4.5-5.5	<2.7	1.3	<1.6	28-40	40-100	<0.6
PATIENT	NA	NA	1.1	1.1	40	71	0.6
CANINE CARDIAC PARAMETERS	HR (BPM)	AV VMAX (m/s)	PV MAX (m/s)	BODY WEIGHT (kg)	LA 2D short axis Base view (cm)	LVIDd Avg; 2D and m-mode short axis (cm)	LVIDs Avg; 2D and m-mode short axis (cm)
NORMAL PARAMETER	50-100	0.7-1.7	0.7-1.6	BELOW	BELOW	BELOW	BELOW
PATIENT	120	1.8	0.94	35.8	2.5	3.8	2.3
*Normal chamber parameters expressed as a mean value (SD)				3	1.27 (5.3)	2.46 (2.46)	1.36 (5.5)
BODY WEIGHT DEPENDENT PARAMETERS				5	1.40 (4.5)	2.74 (5.2)	1.60 (4.7)
<i>*Note: All measurements based upon multi-modal images and methods. An average value is reported.</i>				10	1.50 (3.8)	3.27 (3.5)	2.06 (3.1)
				15	1.83 (2.0)	3.71 (2.4)	2.43 (2.1)
				20	2.02 (1.9)	4.14 (2.2)	2.80 (2.0)
				25	2.18 (2.4)	4.48 (2.9)	3.10 (2.5)
				30	2.33 (3.3)	4.83 (3.9)	3.39 (3.4)
				35	2.48 (4.3)	5.17 (5.0)	3.69 (4.5)
				40	2.62 (5.2)	5.48 (6.1)	3.96 (5.4)
				50	2.88 (7.1)	6.07 (8.3)	4.46 (7.4)

HOSPITAL NAME

 Governors Road
 Animal Hospital

REFERRING VET

Dr. Farooq

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Possible single adult heartworm near the bifurcation of the pulmonary artery. This is certainly not definitive in this peripheral location however, as ultrasound is largely insensitive (i.e., adult worms may be easily either missed peripheral or elsewhere). Given a lack of right heart or MPA enlargement, the infestation is considered relatively mild without evidence of significant pulmonary hypertension. No additional issues are identified. The murmur appears benign in origin with a borderline LVOT velocity.

INVOICE

30709

DATE

5/10/23



PATIENT

Tango Taverner

SPECIES

Canine

BREED

American Bulldog

SEX

Male Neutered

AGE

8 years

WEIGHT

78.9lbs

INTERPRETED BY

Maggie Machen Lamy,
DVM DACVIM
(Cardiology)

IMAGING PERFORMED BY

Kelly Reschny, RVT

HOSPITAL NAME

Governors Road
Animal Hospital

REFERRING VET

Dr. Farooq

INVOICE

30709

DATE

5/10/23

Heartworms can cause significant damage to the lung tissue leading to pulmonary damage, pulmonary hypertension and clinical signs such as coughing, decreased ability to exercise, or difficulty breathing. Disease severity can range from an asymptomatic dog with few worms to dogs with severe respiratory signs. In the most severe cases, caval syndrome may develop due to a very high worm burden sheering blood cells as they pass through the heart. Caval syndrome is a life-threatening emergency that requires immediate surgical removal of the worms.


Given that this patient has no reported clinical signs and no right heart enlargement, we do have some flexibility when approaching therapy. Medical management with drugs like Sildenafil and prednisone is typically utilized if the patient is showing respiratory signs or syncope (none noted). If extraction would be a possibility from a financial standpoint, I would consider referral to a local cardiologist for advanced echocardiography and evaluation. If that is not a possibility, utilizing the standard approach to heartworm treatment as dictated by the American Heartworm Society is recommended, including 30 days of doxycycline and heartguard prior to the split immiticide protocol. Please see website and protocol for specific information. There is high risk for thromboembolism in any patient, however those with adult worms seen in the PA are no question at elevated risk. At this time, exercise restriction is paramount, including cage rest with leash walks only, as a worm embolus can be a life-threatening complication of the disease. This should be continued for an additional 6-8 weeks following therapy.

Modifications to this protocol are sometimes elected depending on individual circumstances which may involve fewer injections or a "slow kill" method. These are not; however, our standard recommendation as alternate treatment may not result in effective treatment of the infestation. Given that the patient has been on the slow kill for multiple years and there is still concern for a possible adult worm, highly recommend immiticide at this time.

Following treatment, retest for heartworm disease 6 months after completing the full course of therapy. Anesthesia is NOT advised prior to completing the protocol, as vasodilation can lead to increased risk for an embolus. Prognosis is guarded, as the right heart/MPA changes are often permanent and may cause clinical signs (exertional syncope/dyspnea, right-sided CHF) in the future.

During therapy, there is high risk for a worm embolus and breathing rate and effort should be monitored closely. Anti-inflammatory prednisone can be used if becomes symptomatic. Patient will be at high risk for developing clinical signs due to pulmonary hypertension with age given the inherent secondary inflammation and damage to the pulmonary vasculature and lungs, and periodic rechecks may be helpful. Monitor for exertional dyspnea or fainting episodes going forward.

Ideally anesthesia would be avoided in this patient until HW negative. If needed, anesthetic risk is considered elevated, particularly should an adult worm be confirmed. The risk is for an embolus to develop, which can occur at any time yet is exacerbated by heavy sedation or gas anesthetics. Should you elect to proceed, premedicate with a vagolytic due to high vagal tone and ensure a normal response is seen. Cardiac protective drug choices (opioid/benzodiazepine premedication, propofol or alfaxalone induction, isoflurane gas) are recommended. Pre-oxygenate for 5-10 minutes prior to induction and



PATIENT

Tango Taverner

recover in O2 if possible. Monitor for arrhythmias, hypotension, and hypoxia both intra and post-operatively and intervene as necessary. Mild IV fluid restriction is recommended to avoid fluid overload.

SPECIES

Canine

Once heartworm negative, a recheck echocardiogram and chest radiographs are recommended in 6 months to reassess right heart changes.

BREED

American Bulldog

SEX

Male Neutered


AGE

8 years

WEIGHT

78.9lbs

IMAGES



INTERPRETED BY

Maggie Machen Lamy,
DVM DACVIM
(Cardiology)

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

IMAGING PERFORMED BY

Kelly Reschny, RVT

Maggie Machen Lamy, DVM
Diplomate of the American College of Veterinary Internal Medicine (Cardiology)
info@sonopath.com

HOSPITAL NAME

Governors Road
Animal Hospital

REFERRING VET

Dr. Farooq

INVOICE

30709

DATE

5/10/23