



**PATIENT**

Feeny Dotts

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Male Neutered

**AGE**

12.6.10

**WEIGHT**

14.31lbs

**INTERPRETED BY**

Maggie Machen Lamy,  
DVM, DACVIM  
(Cardiology)

**HOSPITAL NAME**

North East Animal  
Hospital

**REFERRING VET**

Dr. Hanlin

**INVOICE**

25748

**DATE**

8.12.22

**PRESENTING CLINICAL SIGNS**

History: Murmur on 2/4/21 exam grade 2/5. None detected on 8/4/22 exam, Indoor/outdoor cat.  
-Bloodwork done 8/4/22: CBC WNL. ProBNP High 628.  
-Current medications: Terramycin Ophthalmic.  
-Sedation used: Not required to complete full diagnostic ultrasound.  
-Pertinent previous ultrasound results: No previous.  
-STAT: Not requested.  
-Imaging performed by: Stephanie Warga RDCS, RVT.

**ELECTROCARDIOGRAPHIC FINDINGS**

A six lead ECG is available at both 25 and 50mm/s; 5mm/mV. The average heart rate is 200bpm with an underlying sinus rhythm. P for every QRS complex and vice versa. The P wave morphology is positive with a normal dimension. Normal PR. The QRS morphology is positive with normal dimension. MEA is borderline, shifted left. VPCs are identified throughout; singles only, monomorphic. No supraventricular beats, pauses or other dysrhythmias observed. ECG diagnosis: Normal sinus tachycardia with isolated VPCs.

**ECHOCARDIOGRAM FINDINGS**

2D, m-mode, color flow and doppler imaging is available. The left ventricular wall is normal in dimension with regions of irregularity. There is a diffusely hyperechoic endocardium consistent with fibrosis. The papillary muscles are remodeled. The mitral valve is normal with no MR. The left atrium is mildly dilated and bulbous in appearance. No obvious smoke. The right atrium is normal. Tricuspid valve is normal with no TR. The right ventricle appears normal. Blood flow through both the LVOT and RVOT is normal in velocity. No pericardial effusion seen. No pleural effusion. No obvious cardiac tumors.

**CARDIAC CHART**

FELINE CARDIAC PARAMETERS	BODY WEIGHT (kg)	HR (BPM)	IVSd (cm) <small>(Moise, Pipers)</small>	LVIDd (cm) <small>(Moise, Pipers)</small>	LWVd (cm) <small>(Moise, Pipers)</small>	FS (%)	EF (%)
NORMAL PARAMETER	-----	150-240	3.5-0.55	<2 (mean 1.5)	3.5-0.55	35-67	80-100
PATIENT	6.5	208	0.53	1.54	0.52	46	80
FELINE CARDIAC PARAMETERS	LA/AO (Boon)	LA/AO HEART BASE (Swe) (Abbott)	LA 2D short axis Base view (cm) (Abbott)		LVOT VEL (m/s)	RVOT VEL (m/s)	E max (m/s)
NORMAL	<1.5	<1.3	<1.2		<1.6	<1.3	<0.9
PATIENT	NM	1.5	1.4		1.0	0.9	NM

Adapted from June Boon, Veterinary Echocardiography, 1998  
Abbott J & MacLean H JVIM 2006;20: 111-119, Moise et al. Am J Vet Res 47:1476, 1986. Pipers et al. Am J Vet Res 40:882, 1979.

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The finding of mild left atrial enlargement in the face of normal LV dimensions is most consistent with Unclassified Cardiomyopathy (UCM); however, some prior infectious or inflammatory issues cannot be ruled out. Mild left atrial dilation is present, which may suggest risk for progression going forward. No additional structural issues are identified.

Additionally, there is an arrhythmia present on the ECG with isolated VPCs identified. The abnormal beats are singles only, albeit relatively frequent. What is seen here in an asymptomatic cat (i.e., not syncopal) does not warrant anti-arrhythmic therapy; however, the arrhythmia should be monitored closely in the future. It is important to note that anti-arrhythmics in cats are difficult to use and should only be instituted if sustained arrhythmias or syncope are noted in the future.

While mild structural disease can lead to development of VPCs, full systemic evaluation should be considered to rule out ancillary systemic issues.

Given what is seen here, no indication for medications at this time. Follow up is certainly advised, as any progressive left atrial enlargement will warrant medical management.

The long-term prognosis given the totality of the findings is guarded; however, there is a highly variable rate of progression in cats with subclinical disease. There will always remain risk for progression to CHF and development of blood clots and/or sudden death in the future. Monitoring is certainly advised, particularly should any respiratory signs, collapse or significant lethargy be noted in the future.


Anesthetic risk is considered moderately elevated due to a combination of mild LAE and VPCs. Drugs that stimulate heart rate should be avoided unless clinically necessary (glycopyrrolate, atropine). Avoid ketamine, telazol, alpha 2 agonists. A reasonable protocol includes opioid/benzodiazepine premedication, propofol induction, isoflurane maintenance. Monitor ECG intra and post-operatively, with careful intervention if ventricular arrhythmias are sustained (i.e., sustained VT) and lead to hemodynamic compromise.

## PLAN

Consider systemic evaluation as discussed.

A recheck echocardiogram and ECG is recommended in 6 months to assess progression.

## IMAGES



**The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.**

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

**Maggie Machen Lamy, DVM**  
**Diplomate of the American College of Veterinary Internal Medicine (Cardiology)**  
**info@sonopath.com**