



PATIENT

Jasper Ashley

SPECIES

Canine

BREED

Pomeranian

SEX

Male Neutered

AGE

10 years

WEIGHT

9.56lbs

INTERPRETED BY

Maggie Machen
Lamy, DVM
DACVIM (Cardiology)

**IMAGING
PERFORMED BY**

Pamela Harrigan,
RDCS

HOSPITAL NAME

Mass Veterinary
Specialty Services

REFERRING VET

Dr. Masloski

INVOICE

20708

DATE

8/24/21

PRESENTING CLINICAL SIGNS

History: Recheck echo. History chronic valvular disease - Stage B1. Currently, No coughing or dyspnea but is a nervous panter. Good appetite and energy. Recently was seen for retinal detachment. At that time, blood pressure was significantly elevated - started on Amlodipine. CV/RESP: NSR, grade IV/VI murmur with PMI left apical area radiating to right, PSS, lung fields clear. BP: 100mmHg x 5.

-Current medications:1) Amlodipine 1mg/ml 0.4mls daily 2) Pred acetate 1 drop OS twice a day 3) Flurbiprofen 1 drop OS twice a day 4) Dorzolamide 1 drop OS twice a day 5) Ocunovis 1 drop OS twice a day.

-Pertinent previous echo findings (1/11/21 MML): LA 1.6 cm; LA:Ao 1.1; LV 2.8 cm; normal LA size; moderate MR; trace TR (2.2 m/s). *Sedated with propofol.

ECHOCARDIOGRAM FINDINGS

2D, m-mode, color flow and Doppler imaging is available.

Left ventricle: Mild LV dilation with adequate myocardial function. LV wall thicknesses are normal.

Left atrium: The left atrium is moderately dilated.

Mitral valve: The mitral valve is diffusely thickened with significant prolapse into the left atrial lumen. Moderate eccentric mitral regurgitation with a normal velocity.

Aortic valve/aorta: The aortic valve is normal in morphology and mobility. Normal aortic outflow velocity; laminar flow. No aortic insufficiency.

Right ventricle: Normal right ventricular diameter and morphology indicating no overt evidence of pulmonary arterial hypertension.

Right atrium: Normal RA dimension.

Tricuspid valve: The tricuspid valve appears normal with trace tricuspid regurgitation; mildly elevated velocity consistent with early pulmonary hypertension.

Pulmonic valve/pulmonary artery: The pulmonic valve is normal in morphology and mobility. No pulmonic insufficiency. Normal RVOT velocity; laminar flow.

Pericardium/other: No pericardial or pleural effusion noted. No obvious cardiac masses.

Heart rhythm: ECG reveals a sinus rhythm with an average HR of 130bpm.

2-Dimensional Measurements


Ao diam (cm)	1.3
LA diam (cm)	2.3
LA:Ao (Swe)	1.77
IVS thickness (cm)	0.6
LVID diastole (cm)	2.8
PW thickness (cm)	0.58
LVID systole (cm)	1.5
FS (%)	46

Doppler Measurements

PV Vmax (m/s)	0.57
AoV Vmax (m/s)	0.95
MR Vmax (m/s)	4.5
TR Vmax (m/s)	3.2
TR PG (mmHg)	41

INTERPRETATION OF THE FINDINGS

Chronic degenerative valve disease persists with evidence of progression. Previously normal LA dimension is now moderately enlarged and there is development of early



PATIENT

Jasper Ashley

pulmonary hypertension. This is concerning for progressive issues going forward, and Pimobendan is warranted at this juncture. In an asymptomatic dog no additional medications are clearly indicated at this time; however, close monitoring at home is advised.

SPECIES

Canine

Prognosis is guarded at this stage (B2), with risk for spontaneous CHF, development of arrhythmias, LA tear and/or sudden death going forward.

BREED

Pomeranian

RECOMMENDATIONS

- Institute Pimobendan 0.2-0.3mg/kg PO q12h.
- Omega fatty acid supplementation and mild salt restriction may be of some long-term benefit.
- Once on Pimobendan for 3-5 days, anesthetic risk is considered mildly elevated. Cardiac protective drug choices (opioid/benzodiazepine premedication, propofol or alfaxalone induction, isoflurane gas) are recommended. Pre-oxygenate for 5-10 minutes prior to induction. Monitor for arrhythmias, hypotension, and hypoxia both intra and post-operatively and intervene as necessary. Mild IV fluid restriction is recommended to avoid fluid overload. Avoid heart rate stimulating drugs such as atropine unless clinically indicated.
- Monitor for development of a cough, labored breathing, exercise intolerance or collapse episodes. Monitoring of sleeping breathing rates is recommended to screen for CHF at home.

SEX

Male Neutered

AGE

10 years

WEIGHT

9.56lbs

INTERPRETED BY

Maggie Machen
Lamy, DVM
DACVIM (Cardiology)

PLAN

- Recommend conservative monitoring with a recheck echocardiogram in 6 months, sooner if any development of clinical signs.

IMAGING

PERFORMED BY

Pamela Harrigan,
RDCS


HOSPITAL NAME

Mass Veterinary
Specialty Services

REFERRING VET

Dr. Masloski

IMAGES



INVOICE

20708

DATE

8/24/21



PATIENT

Jasper Ashley

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

SPECIES

Canine

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

BREED

Pomeranian

Maggie Machen Lamy, DVM

Diplomate of the American College of Veterinary Internal Medicine (Cardiology)

info@sonopath.com

SEX

Male Neutered

Echocardiogram performed by: Pamela Harrigan, RDCS

Pet Animal Ultrasound Service (4paus.com)

AGE

10 years

WEIGHT

9.56lbs

INTERPRETED BY

Maggie Machen
Lamy, DVM
DACVIM (Cardiology)

**IMAGING
PERFORMED BY**

Pamela Harrigan,
RDCS

HOSPITAL NAME

Mass Veterinary
Specialty Services

REFERRING VET

Dr. Masloski

INVOICE

20708

DATE

8/24/21