



PATIENT

Berkley Asselin

SPECIES

Canine

BREED

Boxer

SEX

Female Spayed

AGE

~11 years

WEIGHT

60.7lbs

INTERPRETED BY

Maggie Machen Lamy,
DVM, DACVIM
(Cardiology)

IMAGING PERFORMED BY

Potomac Mobile
Veterinary Ultrasound

HOSPITAL NAME

Banfield Leesburg

REFERRING VET

Dr. Jarrett

INVOICE

21084

DATE

9/17/21

PRESENTING CLINICAL SIGNS

History: Elevation in liver enzymes in the past. Previously suspected to have gall bladder disease and was put on Actigall and did well. Currently acting a bit lethargic/seems a bit off. Leaking urine some. -Abnormal PE/Chem/CBC/UA Results: ALT 338, ALP 523, GGT 152, WBC 5.76 Lymph 0.53.

ELECTROCARDIOGRAPHIC FINDINGS *Note: Single lead ECGs are evaluated as a rhythm strip. Morphology/MEA cannot be definitively commented on.

A single lead ECG is available; 25mm/s, 20mm/mV. The average heart rate is 120bpm (range 107-136bpm). The rhythm is sinus in origin, with a p for every QRS complex and vice versa. The P and QRS morphologies are positive. No ectopic beats, pauses or other dysrhythmias observed. ECG diagnosis: Normal sinus rhythm with respiratory variation.

ECHOCARDIOGRAM FINDINGS

2D, m-mode, color flow and doppler imaging is available. Mild mitral valve thickening with no obvious prolapse into the left atrial lumen. Trace/mild mitral regurgitation. Normal MR velocity. Normal left atrial dimension. Normal LV diameter with normal myocardial function. The tricuspid valve appears subjectively normal, and there is no tricuspid regurgitation. The right heart appears normal (subjective). No overt evidence of pulmonary arterial hypertension. The pulmonic and aortic valves are normal in morphology and mobility. Normal pulmonic and aortic outflow velocities. No aortic or pulmonic insufficiency. No pericardial or pleural effusion noted. No cardiac tumors observed. A single VPC is identified (lead II attached).

CARDIAC CHART

CANINE CARDIAC PARAMETERS	MR VMAX (m/s)	TR VMAX (m/s)	LA/AO (Boon method)	LA/AO (Heart Base; Swe)	FS (%)	EF (%)	EPSS (cm)
NORMAL PARAMETER	4.5-5.5	<2.7	1.3	<1.6	28-40	40-100	<0.6
PATIENT	5.5	NA	NM	1.4	33	62	0.2
CANINE CARDIAC PARAMETERS	HR (BPM)	AV VMAX (m/s)	PV MAX (m/s)	BODY WEIGHT (kg)	LA 2D short axis Base view (cm)	LVIDd Avg; 2D and m-mode short axis (cm)	LVIDs Avg; 2D and m-mode short axis (cm)
NORMAL PARAMETER	50-100	0.7-1.7	0.7-1.6	BELOW	BELOW	BELOW	BELOW
PATIENT	120	1.3	0.9	27.5	2.6	4.1	2.7
*Normal chamber parameters expressed as a mean value (SD)				3	1.27 (5.3)	2.46 (2.46)	1.36 (5.5)
BODY WEIGHT DEPENDENT PARAMETERS				5	1.40 (4.5)	2.74 (5.2)	1.60 (4.7)
*Note: All measurements based upon multi-modal images and methods. An average value is reported.				10	1.50 (3.8)	3.27 (3.5)	2.06 (3.1)
				15	1.83 (2.0)	3.71 (2.4)	2.43 (2.1)
				20	2.02 (1.9)	4.14 (2.2)	2.80 (2.0)
				25	2.18 (2.4)	4.48 (2.9)	3.10 (2.5)
				30	2.33 (3.3)	4.83 (3.9)	3.39 (3.4)
				35	2.48 (4.3)	5.17 (5.0)	3.69 (4.5)
				40	2.62 (5.2)	5.48 (6.1)	3.96 (5.4)
				50	2.88 (7.1)	6.07 (8.3)	4.46 (7.4)

Adapted from June Boon, Veterinary Echocardiography, 1998
Rishniw M and Hollis NE, J Vet Intern Med 2000; 14:429-435
Hansson et al, Vet Rad and Ultrasound 2002
Bonagura et al. Echocardiography: principles of interpretation, Vet Clin North Am 15:1177, 1995



PATIENT

Berkley Asselin

SPECIES

Canine

BREED

Boxer

SEX

Female Spayed

AGE

~11 years

WEIGHT

60.7lbs

INTERPRETED BY

Maggie Machen Lamy,
DVM, DACVIM
(Cardiology)

IMAGING PERFORMED BY

Potomac Mobile
Veterinary Ultrasound

HOSPITAL NAME

Banfield Leesburg

REFERRING VET

Dr. Jarrett

INVOICE

21084

DATE

9/17/21

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Overtly normal cardiac structure and function. A small mitral leak is noted which may reflect early valve disease or may be physiologic in origin. Follow up is advised should a murmur be ausculted in the future. No other significant valve issues are noted. No right heart dilation or structural issues are identified. Systolic function is intact without evidence of DCM.

While no ventricular premature contractions were captured on the screening ECG, a single abnormal beat was identified during the study (single lead ECG attached). A couplet is not ruled out; however, cannot be confirmed. VPCs are generated from abnormal conductive or fibrotic tissue in the ventricles of the heart muscle, and even frequent single VPCs will often cause no clinical signs in dogs. When sustained however, ventricular tachycardia can lead to symptoms such as lethargy and collapse.

VPCs are a very non-specific finding. They can be primary in origin (such as ARVC), be secondary to significant cardiac disease (not present in this study) or be extra-cardiac in origin, i.e., due to pain, stress, inflammation, cancer, GI disease, DIC/sepsis, etc. In a senior Boxer, ARVC is possible (although the most common age of onset is 6-8y). ARVC can occur with or without systolic dysfunction or structural issues, however this should be monitored going forward for any progressive changes. Recommend rule out other differentials for ectopy in this senior dog through AUS, tick titers, troponin, etc. Unfortunately, there is always an elevated risk for collapse and sudden death in any arrhythmic patient, and even on medications this risk unfortunately still persists. ARVC carries a HIGHLY variable prognosis, with some dogs able to remain asymptomatic for extended periods of time, and others developing exercise intolerance, syncopal episode, and refractory arrhythmias/sudden death imminently.

Based upon the arrhythmia present on the available ECG in this asymptomatic dog (rare), anti-arrhythmic therapy is not clearly indicated. A holter monitor can be considered as the next step to allow monitoring of the rhythm throughout 24 hours of a normal day and help determine if treatment is indicated. An alternative approach would be to simply monitor for clinical signs and recheck ECG in 6 months. Discussion with the owner is advised.


Fish oil supplementation is recommended for dogs with arrhythmias (1000mg of omega 3 and 6 once to twice daily). Mild activity/stress restriction is advised.

Monitor at home for collapse, exercise intolerance, and/or lethargy. If a holter monitor is elected, this will dictate whether therapy is needed and follow up protocol. I would not recommend anesthesia until the results are available if elected. If declined, an ECG should be monitored during general anesthesia and lidocaine administered in the event of sustained VT or malignant arrhythmias. Avoid stimulants such as atropine or glycopyrrolate unless indicated.

No cardiac medications are indicated at this time. Monitor for any development of cough, labored breathing or exercise intolerance.

PLAN

Consider extended ECG and/or Holter monitor as discussed. Consider systemic evaluation as discussed. If a holter is declined, recommend a recheck ECG is recommended in 6 months (sooner if any collapse episodes occur).



PATIENT


Berkley Asselin

A recheck echocardiogram is recommended every 6-12 months to screen for development of dilation/dysfunction.

IMAGES

SPECIES

Canine



BREED

Boxer

SEX


Female Spayed

AGE

~11 years

WEIGHT

60.7lbs



INTERPRETED BY

Maggie Machen Lamy,
DVM, DACVIM
(Cardiology)

IMAGING PERFORMED BY

Potomac Mobile
Veterinary Ultrasound

HOSPITAL NAME

Banfield Leesburg

REFERRING VET

Dr. Jarrett

INVOICE

21084

DATE

9/17/21

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. This report was generated using transcription software, and minor dictation errors may be present. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Maggie Machen Lamy, DVM
Diplomate of the American College of Veterinary Internal Medicine (Cardiology)
info@sonopath.com