



PATIENT

Andre Gleaves

SPECIES

Canine

BREED

Miniature poodle

SEX

Male Neutered

AGE

10-23-2013

WEIGHT

16 lbs

INTERPRETED BY

Andrea Nicastrò DVM
Diplomate ACVIM
(Sm Animal Internal Med)

**IMAGING
PERFORMED BY**

Andrea Nicastrò DVM
Diplomate ACVIM
(Sm Animal Internal Med)

HOSPITAL NAME

Ashley Pines AH

REFERRING VET

Andrea Winney, DVM

INVOICE

22543

DATE

2-12-26

PRESENTING CLINICAL SIGNS

Clinical Exam Findings: Patient has a history of diabetes, which is relatively well-controlled. Pt has been intermittently vomiting and lethargic since 1/29, PU/PD.

Diagnosed with DM on 11/14/25. Started on Vetsulin 4 units BID
Insulin has since been increased to 7 units BID.

Abnormal lab-work values: Most recent BW on 1/29 - ALT 555, ALP 409 and GGT 15, 1+ ketones and trace protein in urine. USG 1.050. Fructosamine 535 on 12/30, spot BG on 1/29 303. CBC unremarkable.

Current Medications: Gabapentin 100 mg/ ml - 0.35 ml q8-12 h; Vetsulin 7 units BID; pt was started on Carprofen 25 mg 1/2-tab po BID on 1/15- discontinued on 1/29

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder and visible portion of the pelvic urethra are normal for the degree of luminal distension. The urine is anechoic with no evidence of debris. Cystic calculi and discrete masses are not observed. The region of the trigone and visible portion of the proximal urethra are normal.

The prostate is normal in size (0.72 cm in width) and shape. Parenchyma is homogenous. The prostatic urethra appears normal without evidence of dilation or obstruction.

The left kidney is normal in size (4.69 cm in length) with a normal shape, architecture and smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with mild loss of corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydronephrosis. Renal vasculature is normal.

The right kidney is normal in size (4.93 cm in length) with a normal shape, architecture and smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with mild loss of corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydronephrosis. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is mildly enlarged (0.65 cm at cranial pole) (0.63 cm at caudal pole) with a normal shape and homogenous parenchyma. The glandular echogenicity and detail are unremarkable. Capsule, cortex, and medullary definition are normal. The phrenicoabdominal vein and surrounding vasculature are normal.

The right adrenal gland is mildly enlarged (1.30 cm at cranial pole) (0.62 cm at caudal pole) with a normal shape and homogenous parenchyma. The glandular echogenicity and detail are unremarkable. Capsule, cortex, and medullary definition are normal. The phrenicoabdominal vein and surrounding vasculature are normal.

Spleen

The spleen is normal in size (1.52 cm in width at the level of the hilus) with a normal capsular contour. There is appropriate echogenicity and echotexture. No focal lesions are observed. Splenic vasculature is normal.

Liver

The liver is subjectively enlarged with slightly swollen peripheral contours. The parenchyma is isoechoic relative to the spleen and diffusely homogeneous in appearance. No distinct focal lesions are



PATIENT

Andre Gleaves

observed. Vascular and biliary tracts are of normal volume with no evidence of congestion. The portal vein to caudal vena cava ratio is approximately 1: 1.

SPECIES

Canine

The gallbladder lumen is moderately distended. The wall is thin and smooth. A small amount of echogenic debris is observed within the lumen (most of which is gravity-dependent, and some of which is adhered to the mucosal surface). The cystic and common bile ducts are normal/not seen.

BREED

Miniature poodle

Gastrointestinal

The gastric lumen is mildly fluid-distended. The gastric wall is diffusely thickened (up to 0.46 cm) with retention of the normal layering pattern. pyloric outflow tract is patent. The small intestinal lumen is not dilated. The small intestinal wall is normal in thickness with a normal layering pattern and appropriate mural detail. Discreet masses are not identified. The ileocecolic junction and colonic wall are normal. The colonic lumen contains some shadowing fecal material. There is no obvious evidence of an obstructive pattern.

SEX

Male Neutered

AGE

10-23-2013

Pancreas

The right limb of the pancreas is visible with normal curvilinear peripheral contours. The parenchyma is largely isoechoic relative to surrounding omental fat and slightly heterogenous in appearance. The pancreatic duct is visible but not overtly dilated. There is no evidence of peripancreatic inflammation or effusion.

WEIGHT

16 lbs

Lymph Nodes

A 1.1 x 0.75 cm periportal lymph node is visualized.

INTERPRETED BY

Andrea Nicastrò DVM
Diplomate ACVIM
(Sm Animal Internal Med)

Free Abdomen

There is no obvious evidence of free fluid.

Other

A brief echocardiogram reveals no evidence of pericardial effusion or obvious right atrial/auricular mass.

IMAGING PERFORMED BY

Andrea Nicastrò DVM
Diplomate ACVIM
(Sm Animal Internal Med)

ULTRASONOGRAPHIC FINDINGS

Primary Findings

- The hepatic changes are nonspecific and could be secondary to inflammatory disease (i.e., cholangiohepatitis, chronic hepatitis), Leptospirosis, hepatotoxicosis, infiltrative neoplasia (i.e., lymphoma), vacuolar hepatopathy, regenerative nodular hyperplasia, other hepatopathy, or some combination thereof.

HOSPITAL NAME

Ashley Pines AH

Secondary Findings

- Mild bilateral adrenomegaly
- Mild bilateral nonspecific age-related renal changes
- The pancreatic changes are most consistent with age-related parenchymal remodeling, potentially secondary to a prior inflammatory episode, early fibrosis or chronic pancreatitis.
- The gastric wall changes are most consistent with gastritis, with a lower possibility of emerging neoplasia.

REFERRING VET


Andrea Winney, DVM

INVOICE

22543

DATE

2-12-26



PATIENT

Andre Gleaves

SPECIES

Canine

BREED

Miniature poodle

SEX

Male Neutered

AGE

10-23-2013

WEIGHT

16 lbs

INTERPRETED BY

Andrea Nicastrò DVM
 Diplomate ACVIM
 (Sm Animal Internal Med)

IMAGING PERFORMED BY

Andrea Nicastrò DVM
 Diplomate ACVIM
 (Sm Animal Internal Med)

HOSPITAL NAME

Ashley Pines AH

REFERRING VET

Andrea Winney, DVM

INVOICE

22543


DATE

2-12-26

- The prominent periportal lymph node is likely reactive, with a lower possibility of emerging neoplasia.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

- Consider Leptospirosis testing (i.e., blood and urine PCR, serology), particularly if clinical suspicion for disease is high.
- Cytologic evaluation of the liver should be considered in this patient if clotting status is appropriate. A fine needle aspirate using a 25-gauge needle is recommended. If cytologic evaluation is inconclusive, consider a surgical liver biopsy with aerobic and anaerobic bile cultures and acquisition of additional hepatic tissue samples for copper quantitation.
- If a conservative approach is desired, consider empirical treatment for bacterial cholangiohepatitis (amoxicillin-clavulanic acid, +/-metronidazole, Denamarin). If no improvement in the liver values is seen within 7-10 days of initiating therapy, antibiotics should be discontinued, and hepatic tissue sampling reconsidered. If liver values improve, continue therapy for at least 4-6 weeks and 1 week beyond normalization of the liver values.
- Consider testing for hyperadrenocorticism with a low-dose dexamethasone suppression test or ACTH stimulation test if clinical signs (i.e., PU/PD) develop in the future.





PATIENT

Andre Gleaves

SPECIES

Canine

BREED

Miniature poodle

SEX

Male Neutered

AGE

10-23-2013

WEIGHT

16 lbs

INTERPRETED BY

Andrea Nicastro DVM
 Diplomate ACVIM
 (Sm Animal Internal Med)

IMAGING PERFORMED BY

Andrea Nicastro DVM
 Diplomate ACVIM
 (Sm Animal Internal Med)

HOSPITAL NAME

Ashley Pines AH

REFERRING VET


Andrea Winney, DVM

INVOICE

22543

DATE

2-12-26





PATIENT

Andre Gleaves

SPECIES

Canine

BREED

Miniature poodle

SEX

Male Neutered

AGE

10-23-2013

WEIGHT

16 lbs

INTERPRETED BY

Andrea Nicastrò DVM
 Diplomate ACVIM
 (Sm Animal Internal Med)

IMAGING PERFORMED BY

Andrea Nicastrò DVM
 Diplomate ACVIM
 (Sm Animal Internal Med)

HOSPITAL NAME

Ashley Pines AH

REFERRING VET


Andrea Winney, DVM

INVOICE

22543

DATE

2-12-26





PATIENT

Andre Gleaves

SPECIES

Canine

BREED

Miniature poodle

SEX

Male Neutered

AGE

10-23-2013

WEIGHT

16 lbs

INTERPRETED BY

Andrea Nicastro DVM
Diplomate ACVIM
(Sm Animal Internal Med)

IMAGING PERFORMED BY

Andrea Nicastro DVM
Diplomate ACVIM
(Sm Animal Internal Med)

HOSPITAL NAME

Ashley Pines AH

REFERRING VET

Andrea Winney, DVM

INVOICE

22543

DATE

2-12-26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Andrea Nicastro, MPH, DVM, Diplomate DACVIM (Small Animal Internal Medicine)
info@SonoPath.com