



|   |   |
|---|---|
| <b>PATIENT</b>  | <b>PRESENTING CLINICAL SIGNS</b>  |
| PomPom NJ So Hills  | History: Urinating in the house, 1 week after Convenia inj, increased thirst  |
| <b>SPECIES</b>  | <b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>  |
| Canine  | <b>Urinary System</b><br>The <b>urinary bladder</b> , trigone, and pelvic urethra are normal in thickness and the mucosal surface is smooth. The bladder lumen is moderately distended with anechoic urine. No masses, inflammatory changes or calculi are observed. Ureteral papillae and visualized portion of the proximal urethra, visible to a depth of 2 cm, are normal.                                |
| <b>BREED</b>  |   |
| Pomeranian  | The <b>prostate</b> is normal in size (0.90 cm in width) and shape. Parenchyma is homogenous. The prostatic urethra appears normal without evidence of dilation or obstruction.   |
| <b>SEX</b>  |   |
| Neutered Male   | The <b>left kidney</b> is normal in size (4.20 cm in length) with a normal shape, smooth peripheral margins, and normal internal architecture. There is mild to moderate loss of corticomedullary distinction. Several hyperechoic shadowing diverticular foci are observed. There is no evidence of pyelectasia, infarcts or hydronephrosis. Renal vasculature is normal.                                    |
| <b>AGE</b>  |   |
| 10 years  | The <b>right kidney</b> is normal size (3.82 cm in length); normal shape and architecture with smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with mild loss of corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydronephrosis. Renal vasculature is normal.  |
| <b>WEIGHT</b>   |   |
| 15.8 lbs  | <b>Adrenal Glands</b><br>The <b>left adrenal gland</b> is normal size (0.39 cm at cranial pole) (0.57 cm at caudal pole) (1.50 cm in length); normal shape; homogenous parenchyma. The glandular echogenicity and detail are unremarkable. Capsule, cortex, and medullary definition are normal. The phrenicoabdominal vein and surrounding vasculature are normal.   |
| <b>INTERPRETED BY</b>   |   |
| Andrea Nicastro,<br>DVM, Diplomate<br>ACVIM ( <i>Small Animal<br/>Internal Medicine</i> ) | The <b>right adrenal gland</b> is normal size (0.80 cm at cranial pole) (0.52 cm at caudal pole) (0.52 cm in length); normal shape; homogenous parenchyma. The glandular echogenicity and detail are unremarkable. Capsule, cortex, and medullary definition are normal. The phrenicoabdominal vein and surrounding vasculature are normal.   |
| <b>IMAGING<br/>PERFORMED BY</b>   | <b>Spleen</b><br>The <b>spleen</b> is normal in size (1.37 cm in width at the level of the hilus) with a normal capsular contour. There is appropriate echogenicity and echotexture. No focal lesions are observed. Splenic vasculature is normal.  |
| Jenn  |   |
| <b>HOSPITAL NAME</b>  | <b>Liver</b><br>The <b>liver</b> is normal to slightly prominent in size with normal curvilinear peripheral contours. There is appropriate echogenicity and echotexture. No overt structural evidence of inflammatory, infiltrative, or regenerative pathology is evident. Vascular and biliary tracts are of normal volume with no evidence of congestion. No pathological hepatic lymphadenopathy observed. |
| Rockaway AH   |   |
| <b>REFERRING VET</b>  | The <b>gall bladder</b> lumen is moderately distended. The wall is thin and smooth. A small to moderate amount of mostly gravity dependent debris is observed within the lumen. The cystic and common bile ducts are normal/not seen.   |
| Dr. Maniar  |   |
| <b>INVOICE</b>  | <b>Gastrointestinal</b><br>The <b>stomach and intestine</b> are free of stasis and exhibit normal peristaltic activity. The gastric lumen is not distended. The gastric wall and pylorus are normal in thickness with a normal layering pattern. The pyloric  |
| 11168   |   |
| <b>DATE</b>   |   |
| 6.28.22   |   |

outflow tract is patent. The small intestinal lumen is not dilated. The small intestinal wall thickness is normal with a normal layering pattern and appropriate mural detail. Discreet masses are not identified. The colonic wall is normal. There is no evidence of an obstructive pattern.

### **Pancreas**

The region of the **pancreas** is isoechoic relative to surrounding omental fat. No obvious parenchymal abnormalities are observed. There is no evidence of regional inflammation or effusion.

### **Free Abdomen**

The **peritoneal cavity** is normal. There is no evidence of inflammation or effusion. The abdominal **lymph nodes** are normal/not visible.

## **ULTRASONOGRAPHIC FINDINGS**

### **Primary Findings**


- Mild age-related renal changes with left dystrophic mineralization

\*An obvious cause for the patient's clinical signs is not identified in this study. Considerations include a resistant urinary tract infection, underlying metabolic issue, endocrinopathy, other.

## **INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

- Baseline lab work, including a CBC, chemistry panel, urinalysis and T4 is recommended, if not already performed.
- A urine culture and sensitivity is also recommended, preferably one week after the Convenia injection has worn off.
- Depending on the results of the above diagnostics, further work-up may be warranted.





The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

**Andrea Nicastro, DVM, Diplomate DACVIM (Small Animal Internal Medicine)**  
info@SonoPath.com