

**DATE PRESENTING CLINICAL SIGNS**

8.2.2022

Progressive skin lesions on feet/muzzle. Cytology from nodule on paw contains pyogranulomatous inflammation, culture pending. History of chronic ear infections Suspect atopy with interdigital furunculosis vs less likely filamentous bacterial or fungal/pseudofungal infection. Chest rads unremarkable excepting some mild vertebral spondylosis. Lamé on RFL (lesions are worst on this paw), reduced appetite, 10-pound weight loss.

**PATIENT**

Harper Leonard

**SPECIES**

Canine

Current Medications: Apoquel 16mg 1.5 once daily, Simplicef 200mg once daily, Gabapentin 300mg BID, Mometamax and ear cleaner

Lab Results: Globulin 6.3, albumin 2.7, mild monocytosis, BUN low at 6, 4DX negative.

Date of Previous IntraPet Ultrasound: No previous.

Sedation: Not required to complete full diagnostic ultrasound.

**BREED**

Stat Report: Not requested.

Labrador

Imaging Performed By: Andi Parkinson, BS, RDMS.

**SEX**

Spayed Female

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN****Urinary System**

The **urinary bladder** wall is normal in thickness and the mucosal surface is smooth. The bladder lumen is moderately distended with anechoic urine. No masses, inflammatory changes or calculi are observed. Ureteral papillae and visualized portion of the proximal urethra, visible to a depth of 2 cm, are normal.

**AGE**

3/31/2014

**WEIGHT**

42. 8kg

The **left kidney** is normal size (7.95 cm in length); normal shape and architecture with smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with mild loss of corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter.

The **right kidney** is normal size (7.98 cm in length); normal shape and architecture with smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with mild loss of corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter.

**INTERPRETED BY**

Andrea Nicastro, DMV,  
Diplomate DACVIM  
(Small Animal  
Internal Medicine)

**Adrenal Glands**

The **left adrenal gland** is normal size (0.47 cm at cranial pole) (0.55 cm at caudal pole) (2.64 cm in length); normal shape; homogenous parenchyma. The glandular echogenicity and detail are unremarkable. Capsule, cortex, and medullary definition are normal. The phrenicoabdominal vein and surrounding vasculature are normal.

**HOSPITAL NAME**

Nexus Veterinary  
Specialists

The **right adrenal gland** is normal size (0.71 cm at cranial pole) (0.51 cm at caudal pole) (2.31 cm in length); normal shape; homogenous parenchyma. The glandular echogenicity and detail are unremarkable. Capsule, cortex, and medullary definition are normal. The phrenicoabdominal vein and surrounding vasculature are normal.

**REFERRING VET**

Dr. Steele

**Spleen**

The **spleen** is normal in size (2.57 cm in width at the level of the hilus) with a normal capsular contour. There is appropriate echogenicity and echotexture. No focal lesions are observed. Splenic vasculature is normal.

**INVOICE**

11305

**Liver**

The **liver** is subjectively normal in size with normal contours and structure. There is appropriate echogenicity and echotexture. No overt structural evidence of inflammatory, infiltrative, or regenerative

pathology is evident. Vascular and biliary tracts are of normal volume with no evidence of congestion. No pathological hepatic lymphadenopathy observed.

The **gall bladder** lumen is moderately distended. The wall is thin and smooth. Luminal contents are mostly anechoic. The cystic and common bile ducts are normal/not seen.

### ***Gastrointestinal***

The **gastric lumen** is not distended. The gastric wall is normal in thickness with a normal layering pattern. The small intestinal lumen is not dilated. The small intestinal wall thickness is normal with a normal layering pattern and appropriate mural detail. Discreet masses are not identified. The colonic wall is normal. There is no evidence of an obstructive pattern.

### ***Pancreas***

The region of the **pancreas** is isoechoic relative to surrounding omental fat. No obvious parenchymal abnormalities are observed. There is no evidence of regional inflammation or effusion.

### ***Free Abdomen***

The **peritoneal cavity** is normal. There is no evidence of inflammation or effusion. A 1.94 x 0.65 cm sublumbar **lymph node** is visualized. The node is normal in shape and echogenicity.


## **ULTRASONOGRAPHIC FINDINGS**


### **Primary Findings**

- Minor bilateral age-related renal changes
- The prominent sublumbar lymph node is likely reactive, with a lower possibility of emerging neoplasia.
- \*The abdomen is otherwise unremarkable. An obvious cause for the patient's weight loss is not identified in this study. Considerations include occult fungal disease or neoplasia, maldigestion/malabsorption, other metabolic disease, nonmetabolic issues, other.

## **INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Recommendations regarding this exam to be implemented by Dr. Cara Steele.





The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

**Andrea Nicastro, MPH, DVM, Diplomate DACVIM (Small Animal Internal Medicine)**  
[info@SonoPath.com](mailto:info@SonoPath.com)

