

**DATE PRESENTING CLINICAL SIGNS**

9/26/22

Cryptorchid and has neutered scheduled for October.

PATIENT

Frankie Miller

Current Medications: None.

Lab Results: WNL.

Date of Previous IntraPet Ultrasound: No previous.

Sedation: Not required to complete full diagnostic ultrasound.

Stat Report: Not requested.

Imaging Performed By: Andi Parkinson, BS, RDMS.

SPECIES

Canine

BREED

Boxer

SEX

Male, intact

AGE

10/29/2021

WEIGHT

58.9 lbs.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

**This study was limited to the urinary system. There is a potential for pathology in organs that were not visualized.

Urinary System

The urinary bladder wall is normal in thickness and the mucosal surface is smooth. The bladder lumen is mildly distended with anechoic urine. No masses, inflammatory changes or calculi are observed. The region of the trigone is normal.

The left kidney is normal size (6.07 cm in length); normal shape and architecture with smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with normal corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter.

The right kidney is normal size (5.98 cm in length); normal shape and architecture with smooth peripheral margins. There is a normal 1:3 cortex to medulla ratio with normal corticomedullary distinction. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter.

Other

The left testicle is visualized within the scrotum and is subjectively normal in size (3.48 cm length; 1.77 cm width) with a normal shape and homogeneous parenchyma. No focal lesions are observed.

The right testicle is located intra-abdominally and is subjectively slightly small in size (2.47 cm x 1.57 cm) and slightly irregular in shape with homogeneous parenchyma.

INTERPRETED BY

Andrea Nicastro, DVM,
 Diplomate ACVIM
 (Small Animal Internal
 Medicine)

HOSPITAL NAME

Healing Paws
 Veterinary Wellness
 Center

REFERRING VET

Dr. Preston

ULTRASONOGRAPHIC FINDINGS


Right cryptorchidism - the right testicle is located intra-abdominally.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Castration. An intra-abdominal approach will be necessary to remove the right testicle.

INVOICE

14011



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Andrea Nicastro, MPH, DVM, Diplomate DACVIM (Small Animal Internal Medicine)
info@SonoPath.com