



PATIENT

Meow Hofstrand

SPECIES

Feline

BREED

DSH

SEX

FS

AGE

11 years

WEIGHT

11.84 lbs

INTERPRETED BY

Bradley Harris, DVM,
DACVECC, DACVIM
(cardiology)

IMAGING PERFORMED BY

Sara Hansen

HOSPITAL NAME

Banfield – Salem
Lancaster

REFERRING VET

Dr. Marcberg

INVOICE

10825

DATE

11/25/2025

PRESENTING CLINICAL SIGNS

Clinical Exam Findings: BAR, Cooperative for exam TPR: 100.8 F (aural), 200 bpm, purr/meowing/45 brpm BCS: 6/9, 11.84 lbs Eyes: OU mild nuclear sclerosis and small immature cataracts Ears: clean and clear Nose/throat: wnl Oral: 4/4 dental calculus, gingivitis, CRT <2 sec, mm pink and moist Skin: wnl, no ectoparasites observed, no evidence of dehydration Heart/Lungs: 3/6 parasternal heart murmur, lungs clear, normal respiratory effort *difficult to assess due to loud purring Abd: significant distention of the abdomen M/S: mild generalized muscle wasting, ambulatory on all four limbs Peri: wnl LN: wnl rectal: not performed Muscle score: mild generalized muscle loss.

Abnormal PE/Chem/CBC/UA Results: ABNORMAL Labwork Values DX: CBC- WBC 33.48 H (3.66-16.31), NEU 31.86 H (1.84-11.01), HCT 30.2 L (30.3-49.7), RBC 6.03 L (6.71-11.97) IOF- BUN 15 L (16-36), GLU 190 H (71-159) Electrolytes- Na 168 H (150-165) SDMA- 16 H (0-14) Cardiopet proBNP- sent to ref lab, FedEx Tracking 4816 4553 0301 Fecal- no sample to collect UA- postponed Current Medications Clavamox Drops, Gabapentin Liquid Oral.

ULTRASONOGRAPHIC EXAMINATION OF THE HEART

FELINE CARDIAC PARAMETERS	BODY WEIGHT (kg)	HR (BPM)	IVSd (cm)	LVIDd (cm)	LVWd (cm)	FS (%)	EF (%)
NORMAL PARAMETER	-----	150-240	0.3-0.6	1.0-2.1	0.25-0.6	35-67	80-100
PATIENT	5.38 kg	NM	0.57	1.42	0.53	61%	92%
FELINE CARDIAC PARAMETERS	LA/AO (M-mode)	LA/AO HEART BASE (Sisson)	LAD LA MAX 4 Chamber		LVOT VEL. (m/s)	RVOT VEL. (m/s)	IVRT (m/)
NORMAL PARAMETER	<1.5	1.6	0.7-1.7		<1.6	<1.3	40-60
PATIENT	1.08	1.10	1.42		0.9	1.1	NM
Adapted from June Boon, Veterinary Echocardiography, 1998 Sisson D et al. JVIM 1991; 5: 232, Jacobs et al. Am J Vet Res 1985; 46:1705							

Cardiac Presentation

The left atrium is normal in dimension. There are no distinct left atrial thrombi/clots or spontaneous echo contrast appreciated. The left ventricle is normal in dimension as well as wall thickness, and no evidence of restriction. Left ventricular systolic function is normal, with adequate contractility based on fractional shortening and systolic left ventricular dimensions. The right atrium and ventricle are subjectively normal in dimension and systolic function. There is evidence of systolic anterior motion of the mitral valve with no overt mitral regurgitation. The tricuspid valve leaflets presented normal linear structure, extension in systole, and union in diastole without regurgitation. The left ventricular outflow tract demonstrated turbulent flow and subjective structural valvular integrity. The visible aorta is unremarkable. Pulmonary outflow tract assessment revealed normal valve structure, laminar flow, and appropriate diameter and distensibility. There is no evidence of semilunar valve insufficiency or pulmonary hypertension documented. There is possible pleural effusion noted with the curvilinear probe but not appreciated with the cardiac sector probes.



PATIENT

Meow Hofstrand

SPECIES

Feline

BREED

DSH

SEX

FS

AGE

11 years

WEIGHT

11.84 lbs

INTERPRETED BY

Bradley Harris, DVM,
 DACVECC, DACVIM
 (cardiology)

IMAGING PERFORMED BY

Sara Hansen

HOSPITAL NAME

Banfield – Salem
 Lancaster

REFERRING VET

Dr. Marcberg

INVOICE

10825

DATE

11/25/2025

ULTRASONOGRAPHIC FINDINGS

- These findings are consistent with dynamic subaortic stenosis, as there is SAM present, but no convincing hypertrophy is identified. It is unlikely that any of the clinical/radiographic signs are related to underlying heart disease.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Given the absence of any underlying heart disease, no cardiac therapy will be recommended. In addition, there are no cardiac objections to fluid therapy or steroid use. Owing to the presence of an outflow tract obstruction, a follow up echo is recommended in another 6-12 months to make sure no progression has occurred.

Anesthesia considerations:


If anesthesia is necessary, then alpha-2 agonists, ketamine, high dose acepromazine, and Telazol should be avoided. Fluid therapy during anesthesia should be considered at a reduced rate (e.g., 5 ml/kg/hour) if possible (i.e., if not hypotensive). A shorter anesthetic duration will reduce the risk of complications. Pre-oxygenation is advised. Premedication with an opioid (i.e., butorphanol, hydromorphone, oxymorphone) with or without a benzodiazepine is generally the safest protocol. An induction agent such as Propofol, alfaxalone can be used to effect. Maintenance of anesthesia with isoflurane or sevoflurane is reasonable.

Diet:

No special considerations are necessary. Any high-quality food from Hills, Royal Canin, Science Diet, Eukanuba, Iams, or Purina is reasonable.

Activity:

No special considerations are necessary.





PATIENT

Meow Hofstrand

SPECIES

Feline

BREED

DSH

SEX

FS

AGE

11 years

WEIGHT

11.84 lbs

INTERPRETED BY

Bradley Harris, DVM,
 DACVECC, DACVIM
 (cardiology)

IMAGING PERFORMED BY

Sara Hansen

HOSPITAL NAME

Banfield – Salem
 Lancaster

REFERRING VET


Dr. Marcberg

INVOICE

10825

DATE

11/25/2025



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Bradley Harris, DVM, DACVECC, DACVIM (cardiology)

info@SonoPath.com