



**PATIENT**

Piper Matekovic

**SPECIES**

Canine

**BREED**

French Bulldog

**SEX**

Spayed Female

**AGE**

7.5 Years

**WEIGHT**

14.9 kg

**INTERPRETED BY**

Brad Harris, DVM,  
 DACVECC, DACVIM  
 (cardiology)

**IMAGING PERFORMED BY**

Crystal Hill

**HOSPITAL NAME**

Aldershot Animal  
 Hospital

**REFERRING VET**

Dr. Patton

**INVOICE**

74233

**DATE**

4/6/26

**PRESENTING CLINICAL SIGNS**

Presented for evaluation of collapsing episodes and urinary incontinence. Episodes began about 2 weeks ago and occur multiple times per day when she gets excited. O describes it as wobbling, spinning then falling over, owner feels that she is conscious for these episodes. PE - no ataxia normal mentation, grade 4/6 heart murmur, tachycardia noted, rhythm was regular, no meds. BP today 209/112.206/132

Abnormal PE/Chem/CBC/UA Results: Please read attached ECG

**ULTRASONOGRAPHIC EXAMINATION OF THE HEART**

CANINE CARDIAC PARAMETERS	BW	HR BPM	LAD 4 ch Long	RAD 4 ch Long	La/Ao Heart Base	LVIDd	LVIDs
<b>NORMAL PARAMETER</b>		50-100			<1.6		
<b>PATIENT</b>	14.9	150	2.05	3.86	1.2	2.16	1.41
CANINE CARDIAC PARAMETERS	FS	EPSS	PV V MAX (m/s)	AV V Max (m/sec)	MR Vmax	TR Vmax	RPA distensibility (normal >30%)
<b>NORMAL PARAMETER</b>	28-40	<0.6	0.7-1.6	0.7-1.7	4.5-5.5	< 2.7	
<b>PATIENT</b>	35	0.3	0.7	1.1	4.3	4.8	NM

**Cardiac Presentation**

The left atrium is normal in dimension. The left ventricle is normal in dimension with normal systolic function. The right atrium and ventricle are moderately enlarged with marginal systolic function. The anterior and posterior mitral valve leaflets are thickened and redundant consistent with myxomatous changes, and there is no significant prolapse. There is trivial mitral regurgitation identified. The tricuspid valve leaflets are thickened and redundant, with moderate tricuspid regurgitation and evidence of severe pulmonary hypertension. The left ventricular outflow tract demonstrated normal laminar flow and the visible aorta is unremarkable. The right ventricular outflow tract assessment revealed normal laminar flow, with appropriate main pulmonary artery diameter and right pulmonary artery distensibility. There is no pulmonic and no aortic valve insufficiency identified. There is no visible pericardial, pleural, or free peritoneal fluid documented. No evidence of hepatic venous congestion is noted. The cardiac chambers, pericardial, and visible extra-cardiac regions were free of masses, spontaneous echo contrast, or thrombi.

**ECG:**

There is a six-lead ECG available for review. The underlying rhythm is regular at an average rate of 150BPM. The rhythm appears to be sinus in origin with narrow QRS complexes (<70ms). There is no atrial or ventricular ectopy and no conduction delay or block identified. This is most consistent with a normal sinus rhythm.



**PATIENT**

Piper Matekovic

**SPECIES**

Canine

**BREED**

French Bulldog

**SEX**

Spayed Female

**AGE**

7.5 Years

**WEIGHT**

14.9 kg

**INTERPRETED BY**

Brad Harris, DVM,  
 DACVECC, DACVIM  
 (cardiology)

**IMAGING PERFORMED BY**

Crystal Hill

**HOSPITAL NAME**

Aldershot Animal  
 Hospital

**REFERRING VET**

Dr. Patton

**INVOICE**

74233

**DATE**

4/6/26

**ULTRASONOGRAPHIC FINDINGS**

These findings identify significant pulmonary hypertension (PH) in the presence of minimal degenerative mitral valve disease. Therefore, cor pulmonale secondary to primary pulmonary disease/PH is the likely cause for morbidity. Pulmonary hypertension in dogs is most commonly secondary to primary respiratory disease (chronic bronchitis, pulmonary fibrosis, or other forms of pulmonary interstitial disease). Pulmonary hypertension can also develop in dogs with severe heartworm disease or secondary to pulmonary thromboembolism (PTE). Less commonly, pulmonary hypertension is identified in dogs as an idiopathic condition. The degree of PH has resulted in right sided cardiac enlargement (cor pulmonale), and commonly causes syncope. The clinical signs are likely attributable to this condition.

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Given the degree of right-sided cardiac enlargement, cardiac therapy is reasonable at this time. Treatment for the PH/presumed respiratory disease is also warranted, as clinical signs are present. Therapy should include Vetmedin (0.25-0.35 mg/kg BID), sildenafil (2 mg/kg BID), and enalapril (0.5 mg/kg BID assuming normotension and lack of renal insult). Baseline thoracic radiographs, blood pressure and chemistry panel should be performed now, and again in 1-2 weeks. A repeat echocardiogram, thoracic radiographs, blood pressure, and chemistry panel is indicated in another 3-6 months, or sooner if progression is suspected, clinical signs develop/worsen, or additional cardiac therapy is being contemplated.

Anesthesia considerations:

Anesthesia should be avoided if possible. If anesthesia is necessary, then alpha-2 agonists, ketamine, and Telazol should be avoided. If an ACE inhibitor (enalapril, benazepril) or spironolactone is being given, it should not be administered on the morning of general anesthesia. Other cardiac medications should be administered per the normal dosing schedule. Fluid therapy during anesthesia should be considered at a reduced rate (5 ml/kg/hour) if possible (i.e., if not hypotensive). A shorter anesthetic duration will reduce the risk of complications. Pre-oxygenation is mandatory. Premedication with an opioid (e.g., butorphanol, hydromorphone, oxymorphone) with or without a benzodiazepine is generally the safest protocol. An induction agent such as Propofol, alfaxalone, or diazepam/etomidate can be used to effect. Maintenance of anesthesia with isoflurane or sevoflurane is reasonable.

Diet:

A high-quality food from Hills, Royal Canin, Science Diet, Eukanuba, Iams, or Purina that is highly palatable with adequate protein and calories for maintaining optimal body condition with mild dietary sodium restriction (< 100 mg/100 kcal) is recommended. Consider omega-3 fatty acid supplementation. Ensure the patient is not currently receiving a boutique, exotic, or grain-free diet.

Activity:

Moderate physical activity (meandering walks, exploring the backyard, playing with toys inside, getting excited when family gets home, etc.) is encouraged, but periods of strenuous aerobic activity (jogging, strenuous outdoor ball play, prolonged play at the dog park, etc.) should be avoided, especially during periods of high heat (> 80 F) and humidity. Dogs with heart disease tend to tolerate cool and cold temperatures much better than high temperatures. Avoid sudden increases in activity (e.g. 2 block walks during the week but 2 mile walks followed by 30 minutes at the dog park on the weekends) as this may be difficult for the cardiovascular system to deal with.



**PATIENT**

Piper Matekovic

**SPECIES**

Canine

**BREED**

French Bulldog

**SEX**

Spayed Female

**AGE**

7.5 Years

**WEIGHT**

14.9 kg

**INTERPRETED BY**

Brad Harris, DVM,  
DACVECC, DACVIM  
(cardiology)

**IMAGING PERFORMED BY**

Crystal Hill

**HOSPITAL NAME**

Aldershot Animal  
Hospital

**REFERRING VET**


Dr. Patton

**INVOICE**

74233

**DATE**

4/6/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Brad Harris, DVM, DACVECC, DACVIM (cardiology)**

[info@SonoPath.com](mailto:info@SonoPath.com)