

PATIENT PRESENTING CLINICAL SIGNS

PATIENT Riley Jensen
SPECIES BAR, nervous, euhydrated. No LN enlargement.
 Canine
BREED Vizsla
SEX Spayed female
 History: Riley came in on 5/5/2026. Riley has been vomiting a white foam about 2-3 times per day. It is difficult to tell if it is sputum v.s vomiting.
 No coughing, sneezing or diarrhea. Eating and drinking is normal. No issues with urination or defecation. No murmur, or arrhythmia auscultated in the cardiac field. No crackles or wheezes in the lung field. Abdominal auscultation yielded no significant findings. Bloodwork and radiographs were not taken; O wants to do U/S first.
 Patient was prescribed 2 tablets of cerenia SID x 2d, patient was responsive to treatment. Vomiting returned after medication was stopped.
 Today, abdominal breathing.
 Patient was prescribed famotidine BID x 7d. Patient has been unresponsive to this as a monopharmacy. Current Medications Famotidine

AGE ULTRASONOGRAPHIC EXAMINATION OF THE HEART

13 years
WEIGHT 27.2 kg
INTERPRETED BY Bradley Harris, DVM, DACVECC, DACVIM (cardiology)
 The left atrium is normal to small. The left ventricle is small with normal systolic function. The right atrium and ventricle are subjectively normal in dimension and display evidence of diastolic collapse/compression. The anterior and posterior mitral and tricuspid valve leaflets presented normal linear structure, extension in systole, and union in diastole without prolapse or myxomatous changes noted. There is no significant mitral or tricuspid valve regurgitation noted. The left ventricular outflow tract demonstrated normal laminar flow and the visible aorta is unremarkable. The right ventricular outflow tract assessment revealed normal laminar flow, and appropriate diameter and distensibility. There is no evidence of semilunar valve insufficiency. There is mild pericardial, but no pleura or free peritoneal fluid noted. There is a mixed echogenic mass at the wall of the right atrioventricular junction. The remaining cardiac chambers and visible extra-cardiac regions were free of masses, spontaneous echo contrast, or thrombi.

IMAGING PERFORMED BY

Kelly Reschny

HOSPITAL NAME

Joshua Creek AH

REFERRING VET

Al-Zubaidy


INVOICE

75403

DATE

5/13/26

CANINE CARDIAC PARAMETERS	Body Weight kg	HR BPM	LAD 4 ch Long	RAD 4 ch Long	La/Ao Heart Base	LVIDd	LVIDs
NORMAL PARAMETER		50-100			<1.6		
PATIENT	27.2 kg	160	3.9	2.74	0.89	3.12	2.23
CANINE CARDIAC PARAMETERS	FS	EPSS	PV V MAX (m/s)	AV V Max (m/sec)	MR Vmax	TR Vmax	RPA distensibility (normal >30%)
NORMAL PARAMETER	28-40	<0.6	0.7-1.6	0.7-1.7	4.5-5.5	< 2.7	
PATIENT	29	0.1	1.2	1.0	NM	NM	NM



PATIENT

Riley Jensen

SPECIES

Canine

BREED

Vizsla

SEX

Spayed female

AGE

13 years

WEIGHT

27.2 kg

INTERPRETED BY

Bradley Harris, DVM,
 DACVECC, DACVIM
 (cardiology)

IMAGING PERFORMED BY

Kelly Reschny

HOSPITAL NAME

Joshua Creek AH

REFERRING VET

Al-Zubaidy

INVOICE

75403

DATE

5/13/26

ULTRASONOGRAPHIC FINDINGS

These findings identify pericardial effusion in the setting of a mass lesion in the wall of the right atrium. The location/appearance of the mass is consistent with hemangiosarcoma. The pericardial effusion is most likely neoplastic in origin.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is a therapeutic benefit to tapping the pericardium (if there is enough fluid). Evaluation of the fluid will occasionally be of diagnostic value, but in most cases the results are simply compatible with hemorrhage. If it is hemangiosarcoma, the prognosis is very poor, as recurrent effusions are likely, as is the presence of neoplasia elsewhere in the body. Ideally, an abdominal ultrasound should be considered to identify evidence of neoplasia elsewhere in the body, which (or may not) affect prognosis. If no additional masses can be identified, there may be merit to considering surgical intervention, either by taking the entire pericardium (subtotal pericardiectomy), or creation of a smaller hole via approaches less invasive than a median sternotomy. The value of surgery is simply to avoid the effects of recurrent effusion and need for multiple taps; unfortunately surgery does not alter the natural course of disease, which usually only affords a prognosis of a few months. Chemotherapy concurrent with surgery can be associated with a prognosis of up to 4-5 months, but this path is considered too aggressive by many owners who will elect to euthanize rather than pursue surgery/chemotherapy.

Anesthesia considerations:

Anesthesia should be avoided until any signs of CHF and pericardial effusion have resolved. If anesthesia is necessary after that time, then alpha-2 agonists, ketamine, high dose acepromazine, and Telazol should be avoided. Fluid therapy is often necessary in the setting of cardiac tamponade to improve venous return. A shorter anesthetic duration will reduce the risk of complications. Pre-oxygenation is advised. Premedication with an opioid (e.g., butorphanol, hydromorphone, oxymorphone) with or without a benzodiazepine is generally the safest protocol. An induction agent such as Propofol, alfaxalone, or diazepam/etomidate can be used to effect. Maintenance of anesthesia with isoflurane or sevoflurane is reasonable.

Diet:

No special considerations are necessary. Any high-quality food from Hills, Royal Canin, Science Diet, Eukanuba, Iams, or Purina is reasonable.

Activity:

Moderate physical activity (meandering walks, exploring the back yard, playing with toys inside, getting excited when family gets home, etc.) is encouraged, but periods of strenuous aerobic activity (jogging, strenuous outdoor ball play, prolonged play at the dog park, etc.) should be avoided, especially during periods of high heat (> 80 F) and humidity. Dogs with heart disease tend to tolerate cool and cold temperatures much better than high temperatures. Avoid sudden increases in activity (e.g. 2 block walks during the week but 2 mile walks followed by 30 minutes at the dog park on the weekends) as this may be difficult for the cardiovascular system to deal with.



PATIENT

Riley Jensen

SPECIES

Canine

BREED

Vizsla

SEX

Spayed female

AGE

13 years

WEIGHT

27.2 kg

INTERPRETED BY

Bradley Harris, DVM,
 DACVECC, DACVIM
 (cardiology)

IMAGING PERFORMED BY

Kelly Reschny

HOSPITAL NAME

Joshua Creek AH

REFERRING VET


Al-Zubaidy

INVOICE

75403

DATE

5/13/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Bradley Harris, DVM, DACVECC, DACVIM (cardiology)

info@SonoPath.com