

PATIENT

Bruiser Fehnel

SPECIES

Feline

BREED

DSH

SEX

Neutered Male

AGE

8 Years 6 Months

WEIGHT

13

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Jessica Green

HOSPITAL NAME

Stanglein Veterinary Clinic

REFERRING VET

Dr. Erin Rothrock

INVOICE

72551

DATE

1/28/26

PRESENTING CLINICAL SIGNS

Patient has a 4-5 day history of hyporexia and lethargy. His physical exam was overall fairly unremarkable, other than a newly diagnosed, low grade heart murmur.

Abnormal PE/Chem/CBC/UA Results: Moderate anemia (HCT 23.3%), moderate neutrophilic leukocytosis (neuts 18.62 U/L, WBC 19.45 K/UL), mild lymphopenia (0.47 K/uL), mild thrombocytopenia (116 K/uL); BUN mildly decreased at 13 mg/dL, moderate hyperglobulinemia (6.3 g/dL), very mild hyperbilirubinemia (1.1 mg/dL)

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended. It has a normal uniform wall thickness. Contents include primarily anechoic fluid with occasional echogenic non-shadowing debris, most consistent with incidental suspended lipid in a cat, possibly combined with exfoliated cells, mucous and/or small blood clots. Both sterile inflammation as well as urinary tract infection can also present with echogenic debris. No masses or definitive cystoliths are observed. The trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

The right kidney is normal is size (4.5 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

The left kidney is normal is size (4.47 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

The right adrenal gland is normal in size (0.40 cm), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.40 cm), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively large in size (1.1 cm thick at the hilus) with a mildly swollen but smooth capsule. Parenchyma is normal and homogenous in echogenicity and echotexture. No focal nodules or masses are observed. Splenic vasculature appears normal.

Liver

Liver is subjectively enlarged (swollen contour). Mild parenchymal remodeling with diffusely mildly coarse architecture and increased portal markings is present. No focal nodules or masses are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.



PATIENT

Bruiser Fehnel

SPECIES

Feline

BREED

DSH

SEX

Neutered Male

AGE

8 Years 6 Months

WEIGHT

13

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Jessica Green

HOSPITAL NAME

Stanglein Veterinary Clinic

REFERRING VET

Dr. Erin Rothrock

INVOICE

72551

DATE

1/28/26

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is mildly distended with a small to moderate amount of echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

The visible small intestine demonstrates areas of moderately thick muscularis layer relative to mucosa (disruption of the normal 1:3 muscularis:mucosa ratio). Small intestinal submucosa is slightly irregular, thick and hyperechoic, without evident loss of layering appreciated. The lumen of the small intestine is empty with no evidence of obstruction or foreign material.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Pancreas

Pancreas is prominent (enlarged) in size, hypoechoic to surrounding tissue and has a mildly irregular undulating contour. Parenchyma is coarse with mixed echogenic remodeling noted. No pancreatic duct dilation is noted.

Free Abdomen

There is a small amount of anechoic free fluid in these images as well as diffusely enhanced hyperechoic fat and mesentery.

Mesenteric lymph nodes are prominent in size with swollen capsular contour. Normal elongated shape (length to width ratio) is maintained. There is no loss of parenchymal detail.

PRIMARY FINDINGS

- Hypoechoic hepatomegaly – This appearance is consistent with an acute hepatopathy or acute cholangiohepatitis. Infiltrative neoplasia (round cell neoplasia) should also be considered.
- Splenomegaly– can be associated with congestion caused by sedation (if sedated) but can also be associated with diffuse infiltrative disease. Both benign conditions such as extramedullary hematopoiesis, lymphoid hyperplasia, amyloidosis as well as infiltrative neoplastic diseases such as round cell neoplasia should be considered.
- Moderate inflammatory bowel disease (IBD) pattern – Thick muscularis has been reported with infiltrative bowel disease including both benign inflammatory disease as well as infiltrative neoplasia such as lymphoma. No loss of layering or distinct characteristics of malignancy are present. Therefore, differentials cannot be further ranked without tissue sampling.
- Concurrent chronic low-grade smoldering pancreatitis can't be ruled out and should be suspected in the face of appropriate clinical signs.
- Moderate reactive mesenteric lymph nodes – infiltrative neoplastic disease cannot be ruled out but is considered less likely.

PATIENT

Bruiser Fehnel

SPECIES

Feline

BREED

DSH

SEX

Neutered Male

AGE

8 Years 6 Months

WEIGHT

13

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Jessica Green

HOSPITAL NAME

Stanglein Veterinary Clinic

REFERRING VET

Dr. Erin Rothrock

INVOICE

72551

DATE

1/28/26

- The free fluid and diffusely enhanced hyperechoic fat and mesentery are non-specific findings and could be related to a diffuse inflammatory process/steatitis associated with infiltrative bowel, hepatic, splenic disease, pancreatitis versus other.

SECONDARY FINDINGS

- Mild amount of echogenic urinary bladder debris.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

If not recently evaluated, a urinalysis and, if indicated based on urinalysis results, urine culture is recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ratio is recommended.

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

Especially given patient's concurrent hyperglobulinemia, infiltrative neoplastic disease affecting the spleen, liver +/- bowel is a consideration, and therefore tissue sampling is recommended, beginning with fine needle aspirates of the liver and spleen if patient's coagulation status is appropriate. Other differentials include infectious disease, and pending results of above, comprehensive infectious disease evaluation could be considered.

Other than supportive/symptomatic medical management of clinical signs, further treatment recommendations are largely dependent on results of the above.



SonoPath
FOSTERING THE ART OF VETERINARY MEDICINE™

SonoPath.com info@sonopath.com 1.800.838.4268



Clinical Sonography & Telecytology
Educational Teleconsultation Services™

PATIENT

Bruiser Fehnel

SPECIES

Feline

BREED

DSH

SEX

Neutered Male

AGE

8 Years 6 Months

WEIGHT

13

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Jessica Green

HOSPITAL NAME

Stanglein Veterinary Clinic

REFERRING VET

Dr. Erin Rothrock

INVOICE

72551

DATE

1/28/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
info@sonopath.com