



<b>PATIENT</b>	<b>PRESENTING CLINICAL SIGNS</b>
Ginger Transue	P presented 1/3 for not eating well and being PUPD. She had a dental performed 12/15 with no complications.
<b>SPECIES</b>	Abnormal PE/Chem/CBC/UA Results: ALT 145(H), ALP 177(H), UA 2+ protein... Rads: Soft tissue opacity just cranial to R kidney with displacement of the pylorus of the stomach to the left side of the abdomen.
Canine	
<b>BREED</b>	<b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>
Mixed	<b>Urinary System</b>
<b>SEX</b>	Urinary bladder is adequately distended. It has a normal uniform wall thickness. Contents include primarily anechoic fluid with occasional echogenic non-shadowing debris, most consistent with exfoliated cells, mucous and/or small blood clots. Both sterile inflammation as well as urinary tract infection can also present with echogenic debris. No masses or cystoliths are observed. The trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.
Spayed Female	
<b>AGE</b>	The right kidney is normal in size (5.04 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.
11 Years 11 Months	
<b>WEIGHT</b>	The left kidney is normal in size (4.38 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.
13.3 Pounds	
<b>INTERPRETED BY</b>	<b>Adrenal Glands</b>
Beth Johnson, DVM DACVIM	Adrenal glands are plump/swollen in size. Normal shape and contour are maintained without evidence of capsular invasion. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal. The right adrenal gland measures 2.54 cm long x 1.2 cm at the cranial pole and 0.67 cm at the caudal pol. A hyperechoic nodule is noted in the cranial pole of the right adrenal gland. Nodule does not disrupt normal shape and/or architecture. The left adrenal gland measures 2.17 cm long x 0.77 cm at the cranial pole and 0.80 cm at the caudal pole.
<b>IMAGING PERFORMED BY</b>	<b>Spleen</b>
Jessica Green	The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.
<b>HOSPITAL NAME</b>	<b>Liver</b>
Stanglein Vet Clinic	Liver is subjectively enlarged (swollen contour) without disruption of architecture. It has a normal homogenous echotexture. Parenchyma is diffusely hyperechoic characterized by less prominent than normal portal vein walls and increased echogenicity relative to the spleen and falciform fat. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.
<b>REFERRING VET</b>	
Dr. Erin Rothrock	
<b>INVOICE</b>	
43966	Gallbladder is moderately distended with anechoic bile as well as mild to moderate suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.
<b>DATE</b>	<b>Gastrointestinal</b>
1/5/23	



<b>PATIENT</b>	The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.
Ginger Transue	
<b>SPECIES</b>	The visible small intestines are normal in wall thickness and layering (canine duodenum < 0.5 cm and feline duodenum < 0.4 cm; other < 0.3 cm). Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.
Canine	
<b>BREED</b>	The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.
Mixed	
	<b>Pancreas</b>
<b>SEX</b>	The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.
Spayed Female	
	<b>Free Abdomen</b>
<b>AGE</b>	There is no evidence of free peritoneal effusion noted in these images.
11 Years 11 Months	There is no apparent lymphadenopathy noted in these images.
	<b>PRIMARY FINDINGS</b>
<b>WEIGHT</b>	<ul style="list-style-type: none"> <li><b>Hyperechoic hepatomegaly</b> - This appearance is non-specific and most consistent with a benign steroid (endocrine) or vacuolar hepatopathy or reactive or idiopathic hepatopathy. Inflammatory and/or infiltrative disease (such as round cell neoplasia) are also possible, but considered less likely.</li> </ul>
13.3 Pounds	
<b>INTERPRETED BY</b>	<ul style="list-style-type: none"> <li><b>Mild to moderate gallbladder debris</b> - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.</li> </ul>
Beth Johnson, DVM DACVIM	
<b>IMAGING PERFORMED BY</b>	<ul style="list-style-type: none"> <li><b>Bilateral adrenomegaly</b> - consistent with adrenal hyperplasia secondary to pituitary dependent hyperadrenocorticism vs stress or normal variant. Interpret in combination with clinical signs of hyperadrenocorticism.</li> </ul>
Jessica Green	
<b>HOSPITAL NAME</b>	<ul style="list-style-type: none"> <li><b>Hyperechoic adrenal nodule (cranial pole right adrenal)</b> - Differentials include primary adrenal cortical adenoma or adenocarcinoma, pheochromocytoma, myelolipoma, adrenal hyperplasia secondary to pituitary disease or metastatic disease. Ultrasound alone cannot differentiate between functional and non-functional nodules and/or between benign and malignant disease. Small nodules without other evidence of abdominal disease (to suggest metastatic disease) and/or clinical signs (to suggest adrenal disease) are most often incidental and should be monitored.</li> </ul>
Stanglein Vet Clinic	
<b>REFERRING VET</b>	
Dr. Erin Rothrock	
<b>INVOICE</b>	
43966	
<b>DATE</b>	<ul style="list-style-type: none"> <li>Urinary bladder debris</li> </ul>
1/5/23	
	<b>SECONDARY FINDINGS</b>



**PATIENT**

Ginger Transue

\*\*The soft tissue opacity cranial to the right kidney described in the radiographs is presumably the hepatomegaly.

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

**SPECIES**

Canine

The described adrenal gland, liver and gallbladder changes are all suggestive of hyperadrenocorticism. Given the reported clinical signs of hyperadrenocorticism, including polyuria, polydipsia, testing for hyperadrenocorticism with a LDDS test is warranted. If a LDDS test has been evaluated with a normal result, investigation of possible atypical hyperadrenocorticism with a full ACTH stimulation adrenal panel to the University of Tennessee could be considered. If clinical signs are not present, monitoring is recommended with testing pursued when/if clinical signs develop. If not recently evaluated, blood pressure is recommended. If not recently evaluated, a urinalysis and, if indicated based on urinalysis results, urine culture are also recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ratio is recommended.

**BREED**

Mixed

**SEX**

Spayed Female

Hyperadrenocorticism, however, does not typically result in a decreased appetite, except for the rare cases of pituitary dependent disease caused by a macroadenoma. This could be further investigated with a CT scan of the pituitary gland.

**AGE**

11 Years 11 Months

Additionally, given this patient's history of a recent dental, potentially oral pain could be contributing to the decreased appetite.

**WEIGHT**

13.3 Pounds

Also, given the appearance of the gallbladder, if clinical signs are consistent with cholangitis or an emerging mucocele, especially cranial abdominal pain, etc., the gallbladder could be contributing to the decreased appetite.

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM


If the clinical picture is consistent with abdominal pain/gallbladder disease and/or oral pain, etc., further testing for hyperadrenocorticism should be delayed until that other problem is addressed, as false positive diagnostic results for hyperadrenocorticism can be present in the face of concurrent disease.

**IMAGING PERFORMED BY**

Jessica Green

**HOSPITAL NAME**

Stanglein Vet Clinic



**REFERRING VET**

Dr. Erin Rothrock

**INVOICE**

43966

**DATE**

1/5/23



**PATIENT**

Ginger Transue

**SPECIES**

Canine

**BREED**

Mixed

**SEX**

Spayed Female

**AGE**

11 Years 11 Months

**WEIGHT**

13.3 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Jessica Green

**HOSPITAL NAME**

Stanglein Vet Clinic

**REFERRING VET**


Dr. Erin Rothrock

**INVOICE**

43966

**DATE**

1/5/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
Beth.Johnson@sonopath.com