



PATIENT

Stella Fleck

SPECIES

Canine

BREED

Pitbull Terrier

SEX

Spayed Female

AGE

12 Years

WEIGHT

31 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Renee Trionfetti VMD

HOSPITAL NAME

East Bradford
Veterinary Hospital

REFERRING VET

Dr. Meghan McGrath,
DVM

INVOICE

12975

DATE

01/05/2026

PRESENTING CLINICAL SIGNS

AUS to further evaluate vomiting and decreased appetite over several days. No hx of indiscriminate eating. Did attempt to eat, ate about 1/3 of food, then vomited 1 hours later (yellow bile, mucous, food). O reported the patient was improving when Stella was seen on Friday Jan 2nd. However, Stella was brought back in for a medical exam on Saturday Jan 3rd due to lethargy and no appetite. PHM: OA, multiple dermal masses. Noted some improvements on outpatient supportive care. Meds: Ondansetron 8mg, Entyce 30mg/ml, Cerenia 60mg, Librela 20mg/ml, Carprovet 100mg- client was instructed to hold off on giving carprovet, Gabapentin 300mg, Trazodone 100mg

Abnormal PE/Chem/CBC/UA Results: CBC: Hct 59%, Plts 314, remainder NSF - Chem: normal LES and renal values, Alb 3.1-n, NSF

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Left kidney is normal in size (5.91 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Right kidney is normal in size (6.31 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

Left adrenal gland is normal in size (0.44 cm at cranial pole and 0.65 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Right adrenal gland is normal in size (0.87 cm at cranial pole and 0.59 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal. The spleen measured 1.7 cm thick at the hilus.

Liver

Liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.



PATIENT

Stella Fleck

Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.

SPECIES

Canine

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is mildly distended with a small to moderate amount of echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

BREED

Pitbull Terrier

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

SEX

Spayed Female

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

AGE

12 Years

Pancreas

Pancreas is prominent (enlarged) in size and mildly irregular in shape with a slightly undulating contour. Parenchyma is coarse in echotexture and heterogenous to hypoechoic in echogenicity. There is some very subtle nonspecific enhanced hyperechoic mesentery in the cranial abdomen.

WEIGHT

31 kg

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

There is no apparent pathologic lymphadenopathy noted in these images.

ULTRASONOGRAPHIC FINDINGS

IMAGING PERFORMED BY

Renee Trionfetti VMD

- Mild gallbladder debris- Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.
- The pancreatic parenchymal changes are very mild/subtle but given the mildly nonspecific surrounding enhanced fat could indicate some chronic low-grade smoldering pancreatitis or potentially a resolving episode of acute pancreatitis. Additionally, and/or alternatively, there may be some focal bowel or stomach inflammation/gastroenteritis contributing to the changes.

HOSPITAL NAME

East Bradford
Veterinary Hospital

REFERRING VET

Dr. Meghan McGrath,
DVM

INVOICE

12975

DATE

01/05/2026

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

- A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.
- A baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.



PATIENT

Stella Fleck

SPECIES

Canine

BREED

Pitbull Terrier

SEX

Spayed Female

AGE

12 Years

WEIGHT

31 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Renee Trionfetti VMD

HOSPITAL NAME

East Bradford
Veterinary Hospital

REFERRING VET

Dr. Meghan McGrath,
DVM


INVOICE

12975

DATE

01/05/2026

- In the meantime, Supportive/symptomatic medical management of clinical signs is recommended, including anti-emetics, gastroprotectants (+/- sucralfate, especially with any history of hematemesis), an appetite stimulant and fluid therapy if indicated, etc.
- Additionally, empirical deworming with a 5-day course of Panacur is recommended as is a full course of empirical Helicobacter triple therapy.
- Finally, if tolerated, a transition in diet could be considered, based on trial-and-error response with some options to consider including a gastrointestinal biome diet vs a hydrolyzed protein diet (sometimes several trials with different brands are necessary) vs an easy to digest, bland or low-fat diet vs other.





PATIENT

Stella Fleck

SPECIES

Canine

BREED

Pitbull Terrier

SEX

Spayed Female

AGE

12 Years

WEIGHT

31 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Renee Trionfetti VMD

HOSPITAL NAME

East Bradford
Veterinary Hospital

REFERRING VET

Dr. Meghan McGrath,
DVM

INVOICE

12975

DATE

01/05/2026



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Beth Johnson, DVM DACVIM

info@sonopath.com