

PATIENT

Lucy Adshead

SPECIES

Canine

BREED

Maltese X

SEX

Spayed Female

AGE

13 Years

WEIGHT

9.3 kg

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Crystal Hill

HOSPITAL NAME

Simcoe AH

REFERRING VET

Dr. Kennedy/Dr. Lancashire

INVOICE

17577

DATE

10/3/22

PRESENTING CLINICAL SIGNS

History: inappropriate elimination in home -weight loss -abd round and tense -some diarrhea with frank blood. Has been on Metronidazole

Abnormal PE/Chem/CBC/UA Results: none run, besides UTI which was relatively normal, potential low grade infection

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended with primarily anechoic contents and occasional echogenic non-shadowing debris. Apical urinary bladder wall is diffusely thick (0.65 cm). Mucosa is hyperechoic and irregular. No masses or cystoliths are observed. The trigone and visible pelvic urethra are normal thickness with a smooth mucosal surface.

The left kidney is uniformly enlarged/swollen (4.25 cm long) with an overall hyperechoic echogenicity and slight loss of corticomedullary definition. Normal smooth peripheral margination and shape are maintained. The renal pelvis is dilated with anechoic fluid and hyperechoic thickened pelvic fat. No overt evidence of neoplasia or mineral is observed. The perinephric area is enhanced by hyperechoic fat and mesentery.

Right kidney is normal is size (4.73 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

Left adrenal gland is normal in size (1.75 cm long x 0.52 cm at cranial pole and 0.5 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Right adrenal gland is normal in size (1.24 cm long x 0.91 cm at cranial pole and 0.58 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.

Liver

Liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.

Gastrointestinal



PATIENT

Lucy Adshead

SPECIES

Canine

BREED

Maltese X

SEX

Spayed Female

AGE

13 Years

WEIGHT

9.3 kg

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Crystal Hill

HOSPITAL NAME

Simcoe AH

REFERRING VET

Dr. Kennedy/Dr. Lancashire

INVOICE

17577

DATE

10/3/22

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

The visible small intestine demonstrates areas of thick muscularis layer relative to mucosa (disruption of the normal 1:3 muscularis:mucosa ratio). Small intestinal submucosa is slightly irregular, thick and hyperechoic, without evident loss of layering appreciated. The lumen of the small intestine is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness and layering. Contents are consistent with normal formed feces and gas.

Pancreas

The observed pancreas appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

Free Abdomen

There is no evidence of peritoneal effusion. There is no apparent lymphadenopathy.

ULTRASONOGRAPHIC FINDINGS

Primary Findings

- Inflammatory bowel disease (IBD) pattern – Thick muscularis has been reported with infiltrative bowel disease including both benign inflammatory disease as well as infiltrative neoplasia such as lymphoma. No aggressive lymphadenopathy, loss of layering, etc. is noted to make lymphoma more probable, but lymphoma cannot be definitively ruled out without tissue sampling.

- Pyelonephritis – These changes are most consistent with chronic pyelonephritis. Chronic scarring and fibrosis and/or chronic nephrolith passage can also result in these pelvic dilation changes. Early infiltrative disease cannot be ruled out but is considered less likely.

- Chronic Cystitis - Urinary bladder wall changes are most consistent with chronic cystitis. Infiltrative neoplasia cannot be ruled out but is considered less likely give the location and diffuse nature of the changes.

Secondary Findings

- Gallbladder debris- Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Urinalysis and, if indicated based on urinalysis results, urine culture are recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ration is recommended.



PATIENT

Lucy Adshead

SPECIES

Canine

BREED

Maltese X

SEX

Spayed Female

AGE

13 Years

WEIGHT

9.3 kg

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Crystal Hill

HOSPITAL NAME

Simcoe AH

REFERRING VET

Dr. Kennedy/Dr. Lancashire

INVOICE

17577

DATE

10/3/22

Following urine culture, recommendations include treating any infection present as a long term complicated urinary tract infection, which means a 4-6 week course of antibiotics, based on culture and sensitivity results, including a second culture a week to 10 days into therapy to assure full clearance, followed by a final culture a week to 10 days after finishing antibiotics. If patient is currently on antibiotics or just recently finished them, it is recommended to wait at least a week to 10 days away from antibiotics to get an accurate culture result.

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

Given this patients reported hematochezia, a fecal exam is recommended if not recently evaluated, as is a fecal enteropathogen PCR panel to Texas A&M GI Laboratory could be considered for further evaluation of possible infectious disease.

In the meantime, in addition to managing the urinary signs, as described above, other therapeutic recommendations include empirical deworming with a 5-day course of Panacur, as well as a probiotic, such as Proviable or Visbiome, and potentially a transition to a hydrolyzed protein diet could be considered.

Ultimately, if GI signs persist, upper and lower endoscopy/colonoscopy may be necessary to biopsy and definitively diagnose the suspected infiltrative bowel disease.







PATIENT

Lucy Adshead

SPECIES

Canine

BREED

Maltese X

SEX

Spayed Female

AGE

13 Years

WEIGHT

9.3 kg

INTERPRETED BY

Beth Johnson, DVM DACVIM

IMAGING PERFORMED BY

Crystal Hill

HOSPITAL NAME

Simcoe AH

REFERRING VET

Dr. Kennedy/Dr. Lancashire

INVOICE

17577

DATE

10/3/22



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM DACVIM

Beth.Johnson@SonoPath.com