



PATIENT

Oreo Tondra

SPECIES

Canine

BREED

English Bulldog

SEX

Neutered Male

AGE

5 Years

WEIGHT

53.4 Pounds

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

James Hornbuckle

HOSPITAL NAME

Golden Isles AH

REFERRING VET

James Hornbuckle

INVOICE

17930

DATE

11/1/22

PRESENTING CLINICAL SIGNS

History: P came in 10-27-22 for vomiting for a few days, has history of eating rocks and possible foreign objects. As of today 11-01-22 p is not vomiting, nml bm., and light eating. Was lethargic past few days but acting more of himself now.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Prostate is normal in size, echotexture and echogenicity for a neutered male.

Left kidney is normal is size (5.8 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Right kidney is normal is size (5.4 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

Left adrenal gland is normal in size (0.4 cm at cranial pole and 0.4 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Right adrenal gland is normal in size (0.7 cm at caudal pole, the cranial pole is not able to be fully visualized), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.


Liver

Liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is moderately distended with echogenic nonshadowing luminal contents and gas, consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. Soft cloth fluid absorbing foreign material is considered less likely but cannot be definitively ruled out. This finding should be interpreted in combination with when patient last ate.



PATIENT

Oreo Tondra

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease.

SPECIES

Canine

The visible colon is normal in wall thickness and layering. Contents are consistent with normal formed feces and gas.

Pancreas

BREED

English Bulldog

The observed pancreas appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

SEX

Neutered Male

Free Abdomen

There is no evidence of peritoneal effusion. There is no apparent lymphadenopathy.

AGE

5 Years

- The stomach is full, however, the bowel also contains what appears to be consistent with normal chyme, indicating that this is likely a postprandial study. If these findings are not consistent with a non-fully-fasted patient, then foreign material, again, while considered less likely, is possible.

WEIGHT


53.4 Pounds

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

It sounds like patients gastrointestinal signs are resolving and patient is clinically improving, however, if clinical signs return, recheck imaging following a 24-hour fast should be considered.

INTERPRETED BY

Beth Johnson, DVM
DACVIM



IMAGING PERFORMED BY

James Hornbuckle

HOSPITAL NAME

Golden Isles AH

REFERRING VET


James Hornbuckle

INVOICE

17930

DATE

11/1/22



PATIENT

Oreo Tondra

SPECIES

Canine

BREED

English Bulldog

SEX


Neutered Male

AGE

5 Years

WEIGHT

53.4 Pounds



INTERPRETED BY

Beth Johnson, DVM
DACVIM

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

IMAGING PERFORMED BY

James Hornbuckle

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

HOSPITAL NAME

Beth Johnson, DVM DACVIM

Golden Isles AH

Beth.Johnson@SonoPath.com

REFERRING VET

James Hornbuckle

INVOICE

17930

DATE

11/1/22