



PATIENT

Mary Bradshaw

SPECIES

Canine

BREED

Beagle Mix

SEX

Spayed Female

AGE

13

WEIGHT

45

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Christensen

HOSPITAL NAME

Tranquility Veterinary
Clinic

REFERRING VET

Dr. Peng

INVOICE

12467

DATE

11/24/25

PRESENTING CLINICAL SIGNS

Coughing. X-rays showed rounded heart with mild pleural effusion, broncho interstitial pattern and mass on tail of spleen.

Abnormal PE/Chem/CBC/UA Results: Currently on probiotics and Cosequin. Grade 3/6 systolic murmur. BNP= 734. Chem/cbc pending.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Kidneys are overall normal in size and shape with smooth peripheral margination. A normal 1:3 cortex to medulla ratio is maintained. The medulla and cortices are uniform in texture with some mild increased cortical echogenicity and mild loss of corticomedullary distinction, expected in this age patient. There is no evidence of pyelectasia, mineral or infarcts observed. The left kidney measures 6.6 cm. The right kidney measures 6.2 cm.

Adrenal Glands

Adrenal glands are plump/swollen in size. Normal shape and contour are maintained without evidence of capsular invasion. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal. The left adrenal gland measures 1.2 cm at the cranial pole and 0.87 cm at the caudal pole. The right adrenal gland measures 0.89 cm at the cranial pole and 1.2 cm at the caudal pole. A hyperechoic nodule is noted in the cranial pole of the left adrenal gland. Nodule does not disrupt normal shape and/or architecture.

Spleen

Spleen is generally normal in size and shape with a smooth capsular contour. Parenchyma is diffusely nodular in appearance characterized by small discrete hypoechoic nodules. Splenic vasculature appears normal. Focally, off the medial aspect of the spleen is an approximately 5.5 cm x 7.6 cm mixed heterogenous largely hyperechoic mass.

Liver

Liver is subjectively enlarged with mildly irregular margins. Parenchyma is moderately heterogenous characterized by multiple poorly defined hypoechoic nodules within otherwise hyperechoic liver parenchyma. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The stomach is moderately distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. If patient was appropriately fasted,



PATIENT

Mary Bradshaw

delayed gastric emptying could be considered. Non-shadowing foreign material is considered less likely but cannot be definitively ruled out. If clinical signs are consistent (vomiting, etc.), recommendations include supportive medical care, 24 hours fasting and re-image.

SPECIES

Canine

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta/chyme. There is no evidence of obstruction, foreign material or infiltrative disease.

BREED

Beagle Mix

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

SEX

Spayed Female

Pancreas

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

AGE

13

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

WEIGHT

45

There is no apparent pathologic lymphadenopathy noted in these images.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

ULTRASONOGRAPHIC FINDINGS

Primary Findings

- Diffusely splenic micronodular hyperplasia pattern – This nodular change is often associated with benign aging nodular hyperplasia. Infiltrative neoplasia, however, including both early hemangiosarcoma as well as round cell neoplasia cannot be ruled out. Focally, the mass could represent fibrosis or calcification of an old hematoma or infarct, chronic inflammation, granulomatous disease, myelolipomas or infiltrative neoplasia such as sarcoma versus other cannot be ruled out without tissue sampling.
- Moderately heterogenous liver – These changes are most consistent with benign processes such as nodular hyperplasia, steroid (vacuolar) hepatopathy, extramedullary hematopoiesis or possibly chronic inflammatory disease and less commonly infiltrative round cell or metastatic neoplasia.
- Moderate gallbladder debris- Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.

IMAGING PERFORMED BY

Christensen

HOSPITAL NAME

Tranquility Veterinary
Clinic

REFERRING VET

Dr. Peng

INVOICE

12467

DATE

11/24/25

Secondary Findings

- Bilateral adrenomegaly – In a patient diagnosed with hyperadrenocorticism, this finding is most consistent with adrenal hyperplasia secondary to pituitary dependent hyperadrenocorticism. This finding can also be seen with stress and/or normal patient variant.



PATIENT

Mary Bradshaw

SPECIES

Canine

BREED

Beagle Mix

SEX

Spayed Female

AGE

13

WEIGHT

45

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Christensen

HOSPITAL NAME

Tranquility Veterinary
Clinic

REFERRING VET

Dr. Peng

INVOICE

12467

DATE


11/24/25

Interpret in combination with clinical signs of hyperadrenocorticism and/or other adrenal disease.

- Hyperechoic adrenal nodule in the cranial left adrenal gland – Differentials include primary adrenal cortical adenoma or adenocarcinoma, pheochromocytoma, myelolipoma, adrenal hyperplasia secondary to pituitary disease or metastatic disease. Ultrasound alone cannot differentiate between functional and non-functional nodules and/or between benign and malignant disease. Small nodules without other evidence of abdominal disease (to suggest metastatic disease) and/or clinical signs (to suggest adrenal disease) are most often incidental and should be monitored.
- Age-related kidney changes.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

- As is reportedly already in place, an echocardiogram is recommended.
- Fine needle aspirates of the splenic mass are recommended if patient's coagulation status is appropriate.
- Pending results of above or if a cytologic diagnosis is unable to be obtained, an exploratory laparotomy for planned splenectomy could be considered.
- Further evaluation or recommendations regarding the adrenomegaly is largely dependent on patient's clinical history. Having said that, a blood pressure is recommended if not recently evaluated.





PATIENT

Mary Bradshaw

SPECIES

Canine

BREED

Beagle Mix

SEX

Spayed Female

AGE

13

WEIGHT

45

INTERPRETED BY

Beth Johnson, DVM
DACVIM

**IMAGING
PERFORMED BY**

Christensen

HOSPITAL NAME

Tranquility Veterinary
Clinic

REFERRING VET


Dr. Peng

INVOICE

12467

DATE

11/24/25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Beth Johnson, DVM DACVIM

info@sonopath.com