



**PATIENT**

Jasper Virtue

**SPECIES**

Canine

**BREED**

Catahoula

**SEX**

Neutered Male

**AGE**

10 Years

**WEIGHT**

50.4 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Yvonna Aranda

**HOSPITAL NAME**

VCA Vitality AH

**REFERRING VET**

Dr. Clawson

**INVOICE**

35390

**DATE**

11/3/25

**PRESENTING CLINICAL SIGNS**

History: Clinical Exam Findings: Maintaining weight, good appetite but Will Vomit Intermittently--- either at night, or when walking O has recently moved so O thinks that Jasper is vomiting because he's eating more plants/grasses when he goes outside.

Abnormal PE/Chem/CBC/UA Results: ABNORMAL Labwork Values Bw/u/a/fecal WNL HW neg, Lyme's neg, ehrlichia neg, anaplasma neg will email you a copy of results Current Medications Simparica Trio. Also recommended o start him on Provable or Fortiflora.

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

Urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Prostate is normal in size, echotexture and echogenicity for a neutered male.

Left kidney is normal in size (6.03 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Right kidney is normal in size (6.25 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

**Adrenal Glands**

Left adrenal gland is normal in size (0.68 cm at cranial pole and 0.63 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Right adrenal gland is normal in size (0.7 cm at cranial pole and 0.54 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.


**Spleen**

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.

**Liver**

Liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.



**PATIENT**

Jasper Virtue

**SPECIES**

Canine

**BREED**

Catahoula

**SEX**

Neutered Male

**AGE**

10 Years

**WEIGHT**

50.4 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Yvonna Aranda

**HOSPITAL NAME**

VCA Vitality AH

**REFERRING VET**

Dr. Clawson

**INVOICE**

35390

**DATE**

11/3/25

***Gastrointestinal***

The visible stomach wall is normal in thickness and layering. The stomach is moderately distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. If patient was appropriately fasted, delayed gastric emptying could be considered. Non-shadowing foreign material is considered less likely but cannot be definitively ruled out. If clinical signs are consistent (vomiting, etc.), recommendations include supportive medical care, 24 hours fasting and re-image.

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta/chyme. There is no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

***Pancreas***

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

***Free Abdomen***

There is no visible free peritoneal effusion noted in these images.

There is no apparent pathologic lymphadenopathy noted in these images.

**ULTRASONOGRAPHIC FINDINGS**

- There is not a definitive ultrasonographically visible intraabdominal explanation for the patient's reported vomiting, as this is largely an unremarkable/normal postprandial study. Having said that, while the gastrointestinal contents are most consistent with normal ingesta/chyme, especially given the patient's meal history, foreign material, while thought much less likely, can't be definitively ruled out.


**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Especially given the patient's reported eosinophilia, a baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.

A routine fecal/Giardia exam is recommended if not recently evaluated.

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

A fecal enteropathogen PCR panel to Texas A&M GI Laboratory could be considered for further evaluation of possible infectious disease. Contact lab for recommendations on how long to discontinue antibiotics (if indicated) prior to obtaining a stool sample for submission.



**PATIENT**

Jasper Virtue

**SPECIES**

Canine

**BREED**

Catahoula

**SEX**

Neutered Male

**AGE**

10 Years

**WEIGHT**

50.4 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Yvonna Aranda

**HOSPITAL NAME**

VCA Vitality AH

**REFERRING VET**

Dr. Clawson

**INVOICE**

35390


**DATE**

11/3/25

In the meantime, supportive/symptomatic medical management of clinical signs is recommended, including anti-emetics, gastroprotectants (+/- sucralfate, especially with any history of hematemesis), an appetite stimulant and fluid therapy if indicated, etc.

Additionally, empirical deworming with a 5-day course of Panacur is recommended as is a full course of empirical Helicobacter triple therapy.

Finally, if tolerated, a transition in diet could be considered, based on trial-and-error response with some options to consider including a gastrointestinal biome diet vs a hydrolyzed protein diet (sometimes several trials with different brands are necessary) vs an easy to digest, bland or low-fat diet vs other.



**The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.**

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.



**PATIENT**

**Beth Johnson, DVM DACVIM**

Jasper Virtue

[info@sonopath.com](mailto:info@sonopath.com)

**SPECIES**

Canine

**BREED**

Catahoula

**SEX**

Neutered Male

**AGE**

10 Years

**WEIGHT**

50.4 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING  
PERFORMED BY**

Yvonna Aranda

**HOSPITAL NAME**

VCA Vitality AH

**REFERRING VET**

Dr. Clawson

**INVOICE**

35390

**DATE**

11/3/25