



**PATIENT**

Kodiak Samuelson

**SPECIES**

Canine

**BREED**

Pomeranian X

**SEX**

Spayed Female

**AGE**

7 Years

**WEIGHT**

25 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Brian Klug

**HOSPITAL NAME**

Sondel Family Vet  
Clinic

**REFERRING VET**

Dr. Kara Wallisch

**INVOICE**

42688

**DATE**

11/9/22

**PRESENTING CLINICAL SIGNS**

Suspect Cushingoid, check adrenals/kidneys  
Abnormal PE/Chem/CBC/UA Results: BCS 9/9 ALP 2127, rest of chem wnl Urine Cort Cr Ratio: 43 (high)

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

The right kidney is normal in size (4.85 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A hyperechoic band parallel to the corticomedullary border is present. Non-obstructive linear multifocal hyperechoic diverticular foci with acoustic shadowing are noted.

The left kidney is normal in size (4.74 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A hyperechoic band parallel to the corticomedullary border is present. Non-obstructive linear multifocal hyperechoic diverticular foci with acoustic shadowing are noted.

**Adrenal Glands**

The right adrenal gland is normal in size (0.50 cm at the cranial pole and 0.48 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.50 cm at the cranial pole and 0.41 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

**Spleen**

The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.

**Liver**

Liver is subjectively enlarged (swollen contour) without disruption of architecture. It has a normal homogenous echotexture. Parenchyma is diffusely hyperechoic characterized by less prominent than normal portal vein walls and increased echogenicity relative to the spleen and falciform fat. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

**Gastrointestinal**



<b>PATIENT</b>	The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.
Kodiak Samuelson	
<b>SPECIES</b>	The visible small intestines are normal in wall thickness and layering (canine duodenum < 0.5 cm and feline duodenum < 0.4 cm; other < 0.3 cm). Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.
Canine	
<b>BREED</b>	The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.
Pomeranian X	
<b>SEX</b>	<b>Pancreas</b> The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.
Spayed Female	
<b>AGE</b>	<b>Free Abdomen</b> There is no evidence of free peritoneal effusion noted in these images. There is no apparent lymphadenopathy noted in these images.
7 Years	
<b>WEIGHT</b>	<b>PRIMARY FINDINGS</b>
25 Pounds	<ul style="list-style-type: none"> <li><b>Hyperechoic hepatomegaly</b> - This appearance is non-specific and most consistent with a benign steroid (endocrine) or vacuolar hepatopathy or reactive or idiopathic hepatopathy. Inflammatory and/or infiltrative disease (such as round cell neoplasia) are also possible, but considered less likely.</li> </ul>
<b>INTERPRETED BY</b>	<b>SECONDARY FINDINGS</b>
Beth Johnson, DVM DACVIM	<ul style="list-style-type: none"> <li><b>Bilateral medullary rim sign</b> - This finding is of unknown clinical significance and can be a normal variant, often idiopathic. Medullary rim sign can be present with renal disease including FIP, lymphoma, hypercalcemic nephropathy, Leptospirosis, tubular disease, other and should be interpreted in combination with other more specific indications of kidney disease such as isosthenuria, proteinuria, azotemia, etc. This is a common incidental finding in patients with diabetes mellitus.</li> <li>Non-obstructive dystrophic mineralization in both kidneys</li> <li><b>Gallbladder debris</b> - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.</li> </ul>
<b>IMAGING PERFORMED BY</b>	
Brian Klug	
<b>HOSPITAL NAME</b>	
Sondel Family Vet Clinic	
<b>REFERRING VET</b>	
Dr. Kara Wallisch	
<b>INVOICE</b>	<b>INTERPRETATION OF THE FINDINGS &amp; FURTHER RECOMMENDATIONS</b>
42688	Normal appearing adrenal glands does not rule out hyperadrenocorticism and can be present with pituitary dependent hyperadrenocorticism. Therefore, if clinical signs are consistent, then testing in the form of a low-dose Dexamethasone suppression test is recommended.
<b>DATE</b>	Additionally, since clinical signs of diabetes mellitus can mimic hyperadrenocorticism, given this patient's medullary rim sign, ruling out diabetes is recommended.
11/9/22	



**PATIENT**

Kodiak Samuelson

If not recently evaluated, Urinalysis and, if indicated based on urinalysis results, urine culture are recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ration is recommended.

**SPECIES**

Canine

If there is suspicion of hyperadrenocorticism based on history, a blood pressure would also be recommended.

**BREED**

Pomeranian X

**SEX**

Spayed Female

**AGE**

7 Years

**WEIGHT**

25 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Brian Klug

**HOSPITAL NAME**

Sondel Family Vet  
Clinic

**REFERRING VET**


Dr. Kara Wallisch

**INVOICE**

42688

**DATE**

11/9/22





**PATIENT**

Kodiak Samuelson

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

**SPECIES**

Canine

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**BREED**

Pomeranian X

**Beth Johnson, DVM, DACVIM**  
Beth.Johnson@sonopath.com

**SEX**

Spayed Female

**AGE**

7 Years

**WEIGHT**

25 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING  
PERFORMED BY**

Brian Klug

**HOSPITAL NAME**

Sondel Family Vet  
Clinic

**REFERRING VET**

Dr. Kara Wallisch

**INVOICE**

42688

**DATE**

11/9/22