



PATIENT

Molly Miller

SPECIES

Canine

BREED

Mini Dachshund

SEX

Spayed Female

AGE

14

WEIGHT

9.8

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Wehmer

HOSPITAL NAME

Aevendale-Blue Pet
Hospital

REFERRING VET

Dr. Stephanie Wehmer

INVOICE

11255

DATE

2/5/2026

PRESENTING CLINICAL SIGNS

- Historically healthy.
- Isolated seizure episode 11/2025.
- Slight increase in thirst.

Abnormal PE/Chem/CBC/UA Results: CBC/Chem/UA/T4 @ that time USG 1.028 PIT 522k.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Kidneys are overall normal in size and shape with smooth peripheral margination. A normal 1:3 cortex to medulla ratio is maintained. The medulla and cortices are uniform in texture with some mild increased cortical echogenicity and mild loss of corticomedullary distinction, expected in this age patient. There is no evidence of pyelectasia, mineral or infarcts observed. Left kidney measures 3.65 cm, and the right kidney measures 4.12 cm.

Adrenal Glands

The right adrenal gland is normal in size (1.0 cm at cranial pole and 0.58 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.56 cm at cranial pole and 0.48 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). A 1.5 cm x 1.2 cm mildly heterogenous, primarily hypo- to anechoic, non-capsular disrupting nodule mid spleen. Splenic vasculature appears normal.

Liver

Liver is relatively normal in size and contour. Parenchyma is mildly heterogenous and coarse with mild likely age-related parenchymal remodeling noted. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic with some echogenic debris noted. There is no evidence of cystic or common bile duct dilation.

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.



PATIENT

Molly Miller

SPECIES

Canine

BREED

Mini Dachshund

SEX

Spayed Female

AGE

14

WEIGHT

9.8

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Wehmer

HOSPITAL NAME

Aevendale-Blue Pet
Hospital

REFERRING VET

Dr. Stephanie Wehmer

INVOICE

11255

DATE

2/5/2026

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Pancreas

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

There is no apparent pathologic lymphadenopathy noted in these images.

PRIMARY FINDINGS

- Hypo to anechoic splenic nodule - likely represents a benign lesion such as a cyst, hematoma, nodular hyperplasia, extramedullary hematopoiesis, etc., however while considered less likely, infiltrative neoplasia can mimic benign lesions and cannot be ruled out.

SECONDARY FINDINGS


- Mild to moderate age-related kidney changes.
- Mild age-related hepatobiliary changes.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

While the appearance of the spleen trends largely toward benign, fine needle aspirates of the nodule could be considered if patient's coagulation status is appropriate.

Three view thoracic radiographs are recommended for further assessment of cardio-pulmonary status as well as to further evaluate for any evidence of metastatic disease, if not recently evaluated.

Otherwise, further workup is largely dependent on patient's full clinical status. If hyperadrenocorticism is suspected clinically, largely normal appearing adrenal glands does not rule it out.





PATIENT

Molly Miller

SPECIES

Canine

BREED

Mini Dachshund

SEX

Spayed Female

AGE

14

WEIGHT

9.8

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Wehmer

HOSPITAL NAME

Aevendale-Blue Pet
Hospital

REFERRING VET


Dr. Stephanie Wehmer

INVOICE

11255

DATE

2/5/2026



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
info@sonopath.com