



PATIENT

Jasmine Muller

SPECIES

Feline

BREED

Siamese

SEX

Spayed Female

AGE

11 Years

WEIGHT

3.5 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Callihan/Pacific
Crest Mobile VS

HOSPITAL NAME

Pacific Crest Mobile VS

REFERRING VET

Dr. Foot/Village VH

INVOICE

45005

DATE

2/9/23

PRESENTING CLINICAL SIGNS

Ultrasound requested to evaluate for ongoing diarrhea. Jasmine had abdominal ultrasound in September (SonoPath Interp by Dr. Beth Johnson) which showed coarse splenic and liver parenchyma, as well as disruption of normal 3:1 mucosa:muscularis ratio in SI; We did do aspirates of liver and spleen at that time, no evidence of diffuse round cell neoplasia. Pt had been on pred for a while and showed improvement Current diet is Hills z/d

Abnormal PE/Chem/CBC/UA Results: Has history liver enzyme elevations, recheck today is pending

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended. It has a normal uniform wall thickness. Contents include primarily anechoic fluid with occasional echogenic non-shadowing debris, most consistent with incidental suspended lipid in a cat, possibly combined with exfoliated cells, mucous and/or small blood clots. Both sterile inflammation as well as urinary tract infection can also present with echogenic debris. No masses or cystoliths are observed. The trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

The right kidney is normal in size (4.0 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

The left kidney is normal in size (3.87 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

The right adrenal gland is normal in size (0.39 cm), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.33 cm), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively large in size (1.1 cm thick) with subtly scalloped or undulating capsular contour. Parenchyma is normal in echogenicity with a mildly coarse/heterogenous echotexture. No focal nodules or masses are observed. Splenic vasculature appears normal.

Liver

The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

Gastrointestinal



PATIENT

Jasmine Muller

The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

SPECIES

Feline

The visible small intestine demonstrates areas of thick muscularis layer relative to mucosa (disruption of the normal 1:3 muscularis:mucosa ratio). Small intestinal submucosa is slightly irregular, thick and hyperechoic, without evident loss of layering appreciated. The lumen is empty with no evidence of obstruction or foreign material.

BREED

Siamese

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

SEX

Spayed Female

Pancreas

The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

AGE

11 Years

Free Abdomen

There is no evidence of free peritoneal effusion noted in these images.

WEIGHT

3.5 kg

The mesenteric lymph nodes are prominent in size with swollen capsular contour. Normal elongated shape (length to width ratio) is maintained. There is no loss of parenchymal detail.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

PRIMARY FINDINGS

- **Inflammatory bowel disease (IBD) pattern** – Thick muscularis has been reported with infiltrative bowel disease including both benign inflammatory disease as well as infiltrative neoplasia such as lymphoma. No aggressive lymphadenopathy, loss of layering, etc. is noted to make lymphoma more probable, but lymphoma cannot be definitively ruled out without tissue sampling.
- **Reactive mesenteric lymph nodes** – infiltrative neoplastic disease cannot be ruled out but is considered less likely.
- **Scalloped spleen** – can be associated with benign or malignant infiltrative disease. Common causes include a reactive spleen secondary to immune stimulus or early infiltrative round cell neoplasia such as lymphoma or mast cell tumor.

IMAGING PERFORMED BY

Dr. Callihan/Pacific
Crest Mobile VS

HOSPITAL NAME

Pacific Crest Mobile VS

REFERRING VET

Dr. Foot/Village VH

SECONDARY FINDINGS

- Urinary bladder debris

INVOICE

45005

DATE

2/9/23

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The appearance of this study is overall generally improved from the previous scan, likely due to the reported Prednisone therapy. However, splenic pathology and evidence of infiltrative bowel disease are both still present. If this patient was on Prednisone at the time of the previous splenic aspirate, a splenic aspirate is recommended off of the Prednisone if patient's coagulation status is appropriate to look for evidence of infiltrative round cell neoplasia such as lymphoma. Otherwise, biopsies of the GI tract, being sure to include ileum, if possible, are the next diagnostic steps.

Additionally, if a gastrointestinal malabsorption panel was not previously evaluated, a gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.



PATIENT

Jasmine Muller

SPECIES

Feline

BREED

Siamese

SEX

Spayed Female

AGE

11 Years

WEIGHT

3.5 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Callihan/Pacific
Crest Mobile VS

HOSPITAL NAME

Pacific Crest Mobile VS

REFERRING VET


Dr. Foot/Village VH

INVOICE

45005

DATE

2/9/23





PATIENT

Jasmine Muller

SPECIES

Feline

BREED

Siamese

SEX

Spayed Female

AGE

11 Years

WEIGHT

3.5 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Callihan/Pacific
Crest Mobile VS

HOSPITAL NAME

Pacific Crest Mobile VS

REFERRING VET


Dr. Foot/Village VH

INVOICE

45005

DATE

2/9/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
Beth.Johnson@sonopath.com