



## PATIENT

Bebe Yip

## SPECIES

Canine

## BREED

Yorkie

## SEX

Intact Male

## AGE

7 years 7 months

## WEIGHT

6.6 kg

## INTERPRETED BY

Beth Johnson, DVM  
DACVIM

## IMAGING PERFORMED BY

Dr. Jill Rankin

## HOSPITAL NAME

Argo Vet

## REFERRING VET

Dr. Alan Laung

## INVOICE

11467

## DATE

3/11/2026

## PRESENTING CLINICAL SIGNS

- Laboratory abnormalities noted on March 2 and 3, 2026, primarily indicating hepatobiliary disease, hypercholesterolemia, and potential hypothyroidism.
- Elevated ALP 885 U/L; 0-140. Elevated TCHO >11.64mmol/L; 3.10-8.02.
- Elevated GGT 20 U/L; 0-14 & Elevated ALP.
- Low T4 15.2 nmol/L; 16.9-58.5.

## ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

### Urinary System

The urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Prostate is symmetrically enlarged with smooth margins that are well differentiated from surrounding tissue. Normal bilobed shape is maintained. Parenchyma is heterogenous with scattered hyperechoic foci present. No mineral or cysts are noted. Prostate measures 2.5 cm wide in the sagittal view.

The right kidney is normal in size (4.9 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

The left kidney is normal in size (4.8 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

### Adrenal Glands

The right adrenal gland is normal in size (0.56 cm at cranial pole and 0.52 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.34 cm at cranial pole and 0.53 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

### Spleen

The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal lesions except for a discrete, homogenous, hypo- to anechoic non-capsular disrupting nodule in the mid spleen. Splenic vasculature appears normal.

### Liver

Liver is subjectively enlarged with mildly irregular margins. Parenchyma is mildly heterogenous characterized by multiple poorly defined hypoechoic nodules within otherwise hyperechoic liver parenchyma. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is mildly overdistended with a moderate amount of non-dependent, mildly aggregated/inspissated sludge. Hypo to anechoic cystic areas are noted between the gallbladder



## PATIENT

Bebe Yip

## SPECIES

Canine

## BREED

Yorkie

## SEX

Intact Male

## AGE

7 years 7 months

## WEIGHT

6.6 kg

## INTERPRETED BY

Beth Johnson, DVM  
DACVIM

## IMAGING PERFORMED BY

Dr. Jill Rankin

## HOSPITAL NAME

Argo Vet

## REFERRING VET

Dr. Alan Laung

## INVOICE

11467

## DATE

3/11/2026

sludge and luminal wall. The wall is otherwise smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion.

### *Gastrointestinal*

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

### *Pancreas*

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

### *Free Abdomen*

There is no visible free peritoneal effusion noted in these images.

There is no apparent pathologic lymphadenopathy noted in these images.

## PRIMARY FINDINGS

- Emerging mucocele – Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. The non-dependent nature of this sludge combined with the cystic areas are suggestive, however, of possible emerging cystic mucosal hyperplasia or early gallbladder mucocele.
- Mildly heterogenous liver – These changes are most consistent with benign processes such as nodular hyperplasia, steroid (vacuolar) hepatopathy, extramedullary hematopoiesis or possibly chronic inflammatory disease and less commonly infiltrative round cell or metastatic neoplasia.

## SECONDARY FINDINGS

- Hypo to anechoic splenic nodule – likely represents a benign lesion such as a cyst, hematoma, nodular hyperplasia, extramedullary hematopoiesis, etc., however while considered less likely, infiltrative neoplasia can mimic benign lesions, and cannot be ruled out.
- Benign prostatic hyperplasia – Prostatic findings are most consistent with Benign Prostatic Hyperplasia (BPH) and hyperechoic foci consistent with increased vascularity and fibrosis often associated with BPH. Active prostatitis cannot be ruled out. Infiltrative neoplasia cannot be ruled out but is considered less likely.



## PATIENT

Bebe Yip

## SPECIES

Canine

## BREED

Yorkie

## SEX

Intact Male

## AGE

7 years 7 months

## WEIGHT

6.6 kg

## INTERPRETED BY

Beth Johnson, DVM  
DACVIM

## IMAGING PERFORMED BY

Dr. Jill Rankin

## HOSPITAL NAME

Argo Vet

## REFERRING VET

Dr. Alan Laung

## INVOICE


11467

## DATE

3/11/2026

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Patient's reportedly increased alkaline phosphatase is likely at least in part due to the gallbladder changes described above and further recommendations are largely dependent on clinical signs. In an asymptomatic patient, empirical hepatic nutraceuticals including ursodiol could be considered while monitoring for improvement.





## PATIENT

Bebe Yip

## SPECIES

Canine

## BREED

Yorkie

## SEX

Intact Male

## AGE

7 years 7 months

## WEIGHT

6.6 kg

## INTERPRETED BY

Beth Johnson, DVM  
DACVIM

## IMAGING PERFORMED BY

Dr. Jill Rankin

## HOSPITAL NAME

Argo Vet

## REFERRING VET

Dr. Alan Laung

## INVOICE

11467

## DATE

3/11/2026



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
info@sonopath.com