



PATIENT

Alvin Zbeetneff

SPECIES

Canine

BREED

Alaskan Husky Mix

SEX

NM

AGE

11

WEIGHT

33 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Cory

HOSPITAL NAME

Brighton Veterinary
Clinic P.C. Inc.

REFERRING VET

Dr. Brittany
Johnsgaard

INVOICE

11508

DATE

3/17/2026

PRESENTING CLINICAL SIGNS

- Hx of Cushing's disease diagnosed in 2024, has been on treatment with si miao sen.
- Seen today for evaluation of smaller stools lately. Has lost 7 kg since last year. Moderate muscle loss all over body.

Abnormal PE/Chem/CBC/UA Results: CBC- borderline anemia Chem-Mild increased ALP.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is only mildly distended. Visible contents are anechoic. Urinary bladder wall is unable to be fully assessed for pathology without further distension. No visible masses or definitive cystoliths are observed. The trigone and visible pelvic urethra are normal thickness with a smooth mucosal surface. In the face of urinary signs and/or suspected urinary bladder pathology, reassessment after complete filling is recommended.

Prostate is normal in size, echotexture, and echogenicity for a neutered male.

The right kidney is normal is size (7.12 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

The left kidney is normal is size (6.33 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

The areas of the adrenal glands are examined without evident adrenal gland pathology. The full left gland is difficult to fully visualize/isolate for measurement with one pole measuring 0.54 cm, and overall normal appearance. Similarly, the right adrenal gland has a largely subjectively normal appearance, but the cranial pole is difficult to fully visualize/isolate for measurement. The caudal pole measures 0.58 cm.

Spleen

Spleen is subjectively large in size with a mildly swollen but smooth capsule. Parenchyma is normal and homogenous in echogenicity and echotexture. No focal nodules or masses are observed. Splenic vasculature appears normal.

Liver

Liver is subjectively enlarged with mildly irregular margins. Parenchyma is mildly heterogenous characterized by multiple poorly defined hypoechoic nodules within otherwise hyperechoic liver parenchyma. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.



PATIENT

Alvin Zbeetneff

SPECIES

Canine

BREED

Alaskan Husky Mix

SEX

NM

AGE

11

WEIGHT

33 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Cory

HOSPITAL NAME

Brighton Veterinary
Clinic P.C. Inc.

REFERRING VET

Dr. Brittany
Johnsgaard

INVOICE

11508

DATE

3/17/2026

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Pancreas

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

Medial iliac lymph nodes are prominent in size with swollen capsular contour. Normal elongated shape (length to width ratio) is maintained. There is no loss of parenchymal detail.

ULTRASONOGRAPHIC FINDINGS

- Mildly heterogenous liver – These changes are most consistent with benign processes such as nodular hyperplasia, steroid (vacuolar) hepatopathy, extramedullary hematopoiesis or possibly chronic inflammatory disease and less commonly infiltrative round cell or metastatic neoplasia.
- Mild gallbladder debris - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.
- Subjective splenomegaly– can be associated with congestion caused by sedation (if sedated) but can also be associated with diffuse infiltrative disease. Both benign conditions such as extramedullary hematopoiesis, lymphoid hyperplasia, as well as infiltrative neoplastic diseases such as round cell neoplasia should be considered.
- Mildly reactive medial iliac lymph nodes – infiltrative neoplastic disease cannot be ruled out but is considered less likely.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Further recommendations given patient's reported weight loss are largely dependent on caloric intake. If not already evaluated, a thorough evaluation of daily caloric intake is recommended to assure an



PATIENT

Alvin Zbeetneff

SPECIES

Canine

BREED

Alaskan Husky Mix

SEX

NM

AGE

11

WEIGHT

33 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Cory

HOSPITAL NAME

Brighton Veterinary
Clinic P.C. Inc.

REFERRING VET

Dr. Brittany
Johnsgaard

INVOICE

11508

DATE


3/17/2026

adequate daily caloric intake is occurring vs an inadvertent reduction in calories due to change in diet and/or feeding schedule, competitive eating environment, etc.

Urinalysis and, if indicated based on urinalysis results, urine culture is recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ration is recommended.

Especially if patient's daily caloric intake is normal or even increased, further evaluation of digestion and absorption is recommended via a gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

Otherwise, further evaluation for possible pain (dental, orthopedic, other), upper respiratory disease or oropharyngeal disease, cardiac disease and/or neurologic disease vs other as possible causes for decreased appetite and/or unintentional weight loss is also recommended. Additionally, cortisol levels may be warranted.





PATIENT

Alvin Zbeetneff

SPECIES

Canine

BREED

Alaskan Husky Mix

SEX

NM

AGE

11

WEIGHT

33 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Stephanie Cory

HOSPITAL NAME

Brighton Veterinary
Clinic P.C. Inc.

REFERRING VET


Dr. Brittany
Johnsgaard

INVOICE

11508

DATE

3/17/2026



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
info@sonopath.com