

## PATIENT

Mimi Woods

## SPECIES

Canine

## BREED

Yorkie Mix

## SEX

FS

## AGE

11Y, 8M

## WEIGHT

12lbs

## INTERPRETED BY

Beth Johnson, DVM,  
DACVIM (SAIM)

## IMAGING PERFORMED BY

Vincent Ravancho,  
CVT

## HOSPITAL NAME

Legacy Animal Hospital

## REFERRING VET

Dr. Potenzzone

## INVOICE

74299

## DATE

3-23-26

## PRESENTING CLINICAL SIGNS

- Chronic D+
- Chronic on off V+, responsive to cerenia
- No weight loss
- Current medications - Cerenia, Sucralfate, Bland Diet

Abnormal PE/Chem/CBC/UA Results: ALT 135, GGT 13

## ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

### *Urinary System*

The urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Kidneys are overall normal in size (Left 4.75 cm; Right 4.78 cm) and shape with smooth peripheral margination. A normal 1:3 cortex to medulla ratio is maintained. The medulla and cortices are uniform in texture with some mild increased cortical echogenicity and mild loss of corticomedullary distinction, expected in this age patient. There is no evidence of pyelectasia, mineral or infarcts observed.

### *Adrenal Glands*

The left adrenal gland is normal in size (0.48 cm at cranial pole and 0.4 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The right adrenal gland is normal in size (1.0 cm at cranial pole and 0.64 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

### *Spleen*

The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.

### *Liver*

The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

### *Gastrointestinal*

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is moderately overdistended with primarily fluid as well as some echogenic non-shadowing luminal contents and gas consistent with normal chyme. There is no evidence of obstruction, foreign material, or infiltrative disease. Pyloric outflow tract appears patent.



## PATIENT

Mimi Woods

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

## SPECIES

Canine

The colon is mildly thick measuring 0.4 cm thick primarily in the descending colon with normal intact layering and the lumen is empty.

## BREED

Yorkie Mix

### *Pancreas*

Pancreas is prominent (enlarged) in size and mildly irregular in shape with a slightly undulating contour. Parenchyma is coarse in echotexture and heterogenous to hypoechoic in echogenicity.

## SEX

FS

### *Free Abdomen*

There is no visible free peritoneal effusion noted in these images.

## AGE

11Y, 8M

There is no apparent pathologic lymphadenopathy noted in these images.

## ULTRASONOGRAPHIC FINDINGS

### Primary

- The appearance of the colon trends toward a benign colitis with differentials including parasitic, infectious, dietary related, other benign inflammatory with infiltrative neoplasia unable to be ruled out but considered less likely.
- The gastric distension is most likely functional ileus or delayed gastric emptying secondary to underlying bowel or other metabolic disease with no visible evidence of obstruction.
- Chronic low grade smoldering pancreatitis cannot be ruled out.

### Secondary

- Age related kidney changes.

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

A routine fecal/giardia exam is recommended if not recently evaluated.

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

A fecal enteropathogen PCR panel to Texas A&M GI Laboratory could be considered for further evaluation of possible infectious disease. Contact lab for recommendations on how long to discontinue antibiotics (if indicated) prior to obtaining a stool sample for submission.

A baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.

In the meantime:

- Supportive/symptomatic medical management of clinical signs is recommended, including anti-emetics, gastroprotectants (+/- sucralfate, especially with any history of hematemesis), an appetite stimulant and fluid therapy if indicated, etc.
- Additionally, empirical deworming with a 5-day course of Panacur is recommended.

## INVOICE

74299

## DATE

3-23-26

## INTERPRETED BY

Beth Johnson, DVM,  
DACVIM (SAIM)

## IMAGING PERFORMED BY


Vincent Ravancho,  
CVT

## HOSPITAL NAME

Legacy Animal Hospital

## REFERRING VET

Dr. Potenzzone



**PATIENT**

Mimi Woods

**SPECIES**

Canine

**BREED**

Yorkie Mix

**SEX**

FS

**AGE**

11Y, 8M

**WEIGHT**

12lbs

**INTERPRETED BY**

Beth Johnson, DVM,  
DACVIM (SAIM)

**IMAGING PERFORMED BY**

Vincent Ravancho,  
CVT

**HOSPITAL NAME**

Legacy Animal Hospital

**REFERRING VET**

Dr. Potenzone


**INVOICE**

74299

**DATE**

3-23-26

- A full course of empirical Helicobacter triple therapy could be considered.
- A probiotic, such a visbiome or proviable, may be helpful.
- Finally, if tolerated, a transition in diet could be considered, based on trial-and-error response with some options to consider including a gastrointestinal biome diet vs a hydrolyzed protein diet (sometimes several trials with different brands are necessary) vs an easy to digest, bland or low-fat diet vs other.





## PATIENT

Mimi Woods

## SPECIES

Canine

## BREED

Yorkie Mix

## SEX

FS

## AGE

11Y, 8M

## WEIGHT

12lbs

## INTERPRETED BY

Beth Johnson, DVM,  
DACVIM (SAIM)

## IMAGING PERFORMED BY

Vincent Ravancho,  
CVT

## HOSPITAL NAME

Legacy Animal Hospital

## REFERRING VET

Dr. Potenzzone

## INVOICE

74299

## DATE

3-23-26

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
info@sonopath.com