



PATIENT

Harriette Ahearn

SPECIES

Canine

BREED

Corgi Pembroke Welsh

SEX

Spayed Female

AGE

6 Years 4 Months

WEIGHT

16.4 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Jill Rankin

HOSPITAL NAME

VCA Canada Coach Hill
Animal Hospital

REFERRING VET

Dr. Barker

INVOICE

74095

DATE

3/31/26

PRESENTING CLINICAL SIGNS

The patient has a history of proteinuria, which has shown improvement, and a previous finding of splenomegaly. In early December 2025, an ultrasound was done (see previous report for details). On December 24, 2025, blood work revealed slightly elevated triglycerides at 4.23, but the CBC, chemistry panel, precision PSL, and T4 were otherwise unremarkable. A urinalysis from the same date showed a specific gravity of 1.039, 3+ proteinuria, a pH of 6, 1+ bilirubin, and 2 to 3 calcium oxalate dihydrate crystals per high power field. The corresponding urine protein-to-creatinine (UPC) ratio was elevated at 1.5. A follow-up UPC was performed on January 29, which showed a decrease to 0.5. A urinalysis and UPC are being rechecked today.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

The right kidney is normal is size (6.0 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

The left kidney is normal is size (5.5 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

The right adrenal gland is normal in size (0.44 cm at cranial pole and 0.51 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.57 cm at cranial pole and 0.58 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively large in size (2.2 cm thick at the hilus) with normal smooth margins. Parenchyma is normal in echogenicity with a diffusely coarse/heterogenous echotexture. No discrete sizable focal nodules or masses are observed. Splenic vasculature appears normal.


Liver

Liver is subjectively enlarged with mildly irregular margins. Parenchyma is mildly heterogenous characterized by multiple poorly defined hypoechoic nodules within otherwise hyperechoic liver parenchyma. Visible vasculature and biliary tree appear normal without distension or congestion.

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.



PATIENT

Harriette Ahearn

SPECIES

Canine

BREED

Corgi Pembroke Welsh

SEX

Spayed Female

AGE

6 Years 4 Months

WEIGHT

16.4 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Jill Rankin

HOSPITAL NAME

VCA Canada Coach Hill
Animal Hospital

REFERRING VET

Dr. Barker

INVOICE

74095

DATE

3/31/26

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Pancreas

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

There is no apparent pathologic lymphadenopathy noted in these images.

ULTRASONOGRAPHIC FINDINGS

- Mildly heterogenous liver – These changes are most consistent with benign processes such as nodular hyperplasia, steroid (vacuolar) hepatopathy, extramedullary hematopoiesis or possibly chronic inflammatory disease and less commonly infiltrative round cell or metastatic neoplasia.
- Coarse splenomegaly – can be associated with congestion caused by sedation (if sedated) but can also be associated with diffuse infiltrative disease. Both benign conditions such as extramedullary hematopoiesis, lymphoid hyperplasia, as well as infiltrative neoplastic diseases such as round cell neoplasia should be considered.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The appearance of the spleen is static to mildly progressive in size. Therefore, if not already evaluated, a fine needle aspirates of the spleen +/- liver could be considered if patient's coagulation status is appropriate.

Otherwise, given patient's reported improvement, and as is reportedly already planned, monitoring of the proteinuria for complete normalization is recommended to help determine future workup, treatment, etc. if indicated.





PATIENT

Harriette Ahearn

SPECIES

Canine

BREED

Corgi Pembroke Welsh

SEX

Spayed Female

AGE

6 Years 4 Months

WEIGHT

16.4 kg

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Jill Rankin

HOSPITAL NAME

VCA Canada Coach Hill
Animal Hospital

REFERRING VET


Dr. Barker

INVOICE

74095

DATE

3/31/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
info@sonopath.com