



**PATIENT**

Ned Stark Haydock

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Neutered Male

**AGE**

9 Years

**WEIGHT**

4.6 kg

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Callihan

**HOSPITAL NAME**

Animal Emergency  
Care

**REFERRING VET**

Dr. Johnson

**INVOICE**

46871

**DATE**

4/22/23

**PRESENTING CLINICAL SIGNS**

Ultrasound requested as part of workup for severe nonregenerative anemia in indoor only cat presenting as transfer to ER. Also weight loss

Abnormal PE/Chem/CBC/UA Results: On PE mildly febrile, thin topline with somewhat pendulous abd consistent with recent weight loss. Chem 10 showed low BUN and Cr PCV 12% Most recent FeLV/FIV test 2021

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Kidneys are large in size with increased cortical echogenicity. Normal smooth peripheral margination and shape are maintained. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. The left kidney measures 4.78 cm. The right kidney measures 4.66 cm.

**Adrenal Glands**

The right adrenal gland is normal in size (0.32 cm), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.40 cm), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

**Spleen**

Spleen is subjectively large in size (1.24 cm thick) with a mildly swollen but smooth capsule. Parenchyma is normal and homogenous in echogenicity and echotexture. No focal nodules or masses are observed. Splenic vasculature appears normal.

**Liver**

Liver is subjectively enlarged (swollen contour). Mild parenchymal remodeling with diffusely mildly coarse architecture and increased portal markings is present. No focal nodules or masses are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.

**Gastrointestinal**

The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

The visible small intestine demonstrates areas of thick muscularis layer relative to mucosa (disruption of the normal 1:3 muscularis:mucosa ratio). Small intestinal submucosa is slightly irregular, thick and hyperechoic, without evident loss of layering appreciated. The lumen is empty with no evidence of obstruction or foreign material.



<b>PATIENT</b>	The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.
Ned Stark Haydock	
<b>SPECIES</b>	<b>Pancreas</b>
Feline	The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.
<b>BREED</b>	<b>Free Abdomen</b>
DSH	There is no evidence of free peritoneal effusion noted in these images.
<b>SEX</b>	There is no apparent lymphadenopathy noted in these images.
Neutered Male	
<b>AGE</b>	<ul style="list-style-type: none"> <li><b>Hypersplenism</b> – can be associated with congestion caused by sedation (if sedated) but can also be associated with diffuse infiltrative disease. Both benign conditions such as extramedullary hematopoiesis, lymphoid hyperplasia, amyloidosis as well as infiltrative neoplastic diseases such as round cell neoplasia should be considered.</li> </ul>
9 Years	
<b>WEIGHT</b>	<ul style="list-style-type: none"> <li><b>Hypoechoic hepatomegaly</b> – This appearance is consistent with an acute hepatopathy or acute cholangiohepatitis. Infiltrative neoplasia (round cell neoplasia) should also be considered.</li> </ul>
4.6 kg	
<b>INTERPRETED BY</b>	<ul style="list-style-type: none"> <li><b>Inflammatory bowel disease (IBD) pattern</b> – Thick muscularis has been reported with infiltrative bowel disease including both benign inflammatory disease as well as infiltrative neoplasia such as lymphoma. No aggressive lymphadenopathy, loss of layering, etc. is noted to make lymphoma more probable, but lymphoma cannot be definitively ruled out without tissue sampling.</li> </ul>
Beth Johnson, DVM DACVIM	
<b>IMAGING PERFORMED BY</b>	<ul style="list-style-type: none"> <li><b>Feline renomegaly</b> – These renal changes can be seen with glomerular or interstitial nephritis, FIP, amyloidosis, acute tubular necrosis or infiltrative neoplasia such as lymphoma. Normal variant due to fat deposition cannot be ruled out but is less common in an enlarged kidney.</li> </ul>
Dr. Callihan	
<b>HOSPITAL NAME</b>	<b>SECONDARY FINDINGS</b>
Animal Emergency Care	<ul style="list-style-type: none"> <li><b>Mild gallbladder debris</b> - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness, however, it can also be associated with hepatobiliary disease in cats and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.</li> </ul>
<b>REFERRING VET</b>	
Dr. Johnson	
<b>INVOICE</b>	<b>INTERPRETATION OF THE FINDINGS &amp; FURTHER RECOMMENDATIONS</b>
46871	Differentials for this patient’s reported severe non-regenerative anemia include either early hemorrhage or hemolysis without time yet for regeneration to develop, possibly secondary to infectious disease versus infiltrative neoplasia or autoimmune versus anemia of chronic disease or bone marrow disease versus other.
<b>DATE</b>	Recommendations include fine needle aspirates of the spleen +/- liver is patient’s coagulation status is appropriate.
4/22/23	Pending results of the aspirates (if performed), and given the concurrent weight loss the following could be considered:



**PATIENT**

Ned Stark Haydock

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

**SPECIES**

Feline

Ideally, biopsies of the GI tract, being sure to include ileum if possible, are recommended to definitively diagnose and therefore manage the infiltrative bowel disease.

**BREED**

DSH

If biopsies cannot be obtained, empirical therapies could include diet change, empirical deworming with a 5 day course of Panacur, cobalamin supplementation (unless cobalamin level is evaluated and supplementation is not warranted) and prednisolone (if not contraindicated based on patient contraindications, co-morbidities, etc.). Other supportive therapeutic considerations could include fiber supplementation, especially with large bowel diarrhea and/or a probiotic.

**SEX**

Neutered Male

**AGE**

9 Years

**WEIGHT**

4.6 kg

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Callihan

**HOSPITAL NAME**

Animal Emergency Care

**REFERRING VET**


Dr. Johnson

**INVOICE**

46871

**DATE**

4/22/23





**PATIENT**

Ned Stark Haydock

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Neutered Male

**AGE**

9 Years

**WEIGHT**

4.6 kg

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Callihan

**HOSPITAL NAME**

Animal Emergency  
Care

**REFERRING VET**


Dr. Johnson

**INVOICE**

46871

**DATE**

4/22/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
Beth.Johnson@sonopath.com