



<b>PATIENT</b>	<b>PRESENTING CLINICAL SIGNS</b>
Blarney McNally	P presented for US due to suspected PLE. P initially presented on 4/13/26 ADR with diarrhea and weight loss. P previous over 100#. rdvm fast scan minimal ascites seen. P started on Prednisone, Vit B12, and Clopidogrel.
<b>SPECIES</b>	
Canine	Abnormal PE/Chem/CBC/UA Results: Urinalysis usg >1.050, Pro Negative Fecal Negative Chem TP 3.4, Alb 1.3, Glob 2.2, Chol 107
<b>BREED</b>	<b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>
Bernese Mountain Dog	<b><i>Urinary System</i></b>
<b>SEX</b>	The urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.
Spayed Female	
<b>AGE</b>	The right kidney is normal is size (6.77 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.
5 Years 5 Months	
<b>WEIGHT</b>	The left kidney is normal is size (6.72 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.
91.2 lbs	
<b>INTERPRETED BY</b>	<b><i>Adrenal Glands</i></b>
Beth Johnson, DVM DACVIM	Adrenal glands are small (flattened contour). Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal. Left measures 0.36 cm at the cranial pole and 0.49 cm at the caudal pole. Right measures 0.72 cm at the cranial pole and 0.65 cm at the caudal pole.
<b>IMAGING PERFORMED BY</b>	<b><i>Spleen</i></b>
Kathleen Byrnes	The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.
<b>HOSPITAL NAME</b>	<b><i>Liver</i></b>
Pet Care Clinic of the High Country	The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.
<b>REFERRING VET</b>	
Dr. Sturgill	
<b>INVOICE</b>	The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.
74688	<b><i>Gastrointestinal</i></b>
<b>DATE</b>	
4/22/26	The visible stomach wall is normal in thickness and layering. The stomach is moderately distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. If patient was appropriately fasted,



**PATIENT**

Blarney McNally

delayed gastric emptying could be considered. Non-shadowing foreign material is considered less likely but cannot be definitively ruled out.

**SPECIES**

Canine

If clinical signs are consistent (vomiting, etc.), recommendations include supportive medical care, 24 hours fasting and re-image.

**BREED**

Bernese Mountain Dog

The visible small intestines are normal in wall thickness and layering. Hyperechoic mucosal fogging or speckling is noted. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction or foreign material.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

**SEX**

Spayed Female

**Pancreas**

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

**AGE**

5 Years 5 Months

**Free Abdomen**

There is no visible free peritoneal effusion noted in these images.

**WEIGHT**

91.2 lbs

There is no apparent pathologic lymphadenopathy noted in these images.

**INTERPRETED BY**

Beth Johnson, DVM  
 DACVIM

**ULTRASONOGRAPHIC FINDINGS**

- Mild to moderate mucosal speckling – Mucosal speckling is often present with inflammatory bowel disease (IBD). It is not specific for type or severity of disease. Mild speckling change can occur as a normal patient variant in the post-prandial state.
- Flat adrenal glands – This can be a normal patient variant and/or a sign of exogenous cortisol administration. If exogenous steroids are not being administered, hypoadrenocorticism (either relative or absolute) should be considered.

**IMAGING PERFORMED BY**

Kathleen Byrnes

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

**HOSPITAL NAME**

Pet Care Clinic of the  
 High Country

If a pre-steroid sample is available, submission of sample for a baseline cortisol level is recommended. If not, if or when patient is able to come off steroids, a baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.

**REFERRING VET**

Dr. Sturgill

In the meantime, additionally a gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

**INVOICE**

74688

A fecal enteropathogen PCR panel to Texas A&M GI Laboratory could be considered for further evaluation of possible infectious disease. Contact lab for recommendations on how long to discontinue antibiotics (if indicated) prior to obtaining a stool sample for submission.

**DATE**

4/22/26

If hypoadrenocorticism is ruled out, especially given the mucosal speckling, and bowel disease is determined to be the primary cause of patient's clinical signs and hypoalbuminemia, ideally biopsies of the GI tract will be recommended to definitively diagnose and therefore manage the process.



**PATIENT**

Blarney McNally

**SPECIES**

Canine

**BREED**

Bernese Mountain Dog

**SEX**

Spayed Female

**AGE**

5 Years 5 Months

**WEIGHT**

91.2 lbs

**INTERPRETED BY**

Beth Johnson, DVM  
 DACVIM

**IMAGING PERFORMED BY**

Kathleen Byrnes

**HOSPITAL NAME**

Pet Care Clinic of the  
 High Country

**REFERRING VET**

Dr. Sturgill

**INVOICE**

74688


**DATE**

4/22/26

If biopsies cannot be obtained safely due to low albumin or patient stability, etc., empirical therapies could include diet change to an ultra-low-fat diet, empirical deworming with a 5-day course of Panacur, cobalamin supplementation (unless cobalamin level is evaluated and supplementation is not warranted) a probiotic and prednisolone (if not contraindicated based on patient contraindications, co-morbidities, etc.).

Calcium monitoring, and supplementation, if necessary, is also recommended.

Additionally, if patient's coagulation status is otherwise appropriate, anti-thrombotics such as clopidogrel or low dose aspirin may also be warranted.





**PATIENT**

Blarney McNally

**SPECIES**

Canine

**BREED**

Bernese Mountain Dog

**SEX**

Spayed Female

**AGE**

5 Years 5 Months

**WEIGHT**

91.2 lbs

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING  
PERFORMED BY**

Kathleen Byrnes

**HOSPITAL NAME**

Pet Care Clinic of the  
High Country

**REFERRING VET**

Dr. Sturgill

**INVOICE**

74688

**DATE**

4/22/26

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
[info@sonopath.com](mailto:info@sonopath.com)