

PATIENT

Dallas Thornton

SPECIES

Canine

BREED

Boxer Mix

SEX

Spayed Female

AGE

11

WEIGHT

53

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. John Sampson

HOSPITAL NAME

Richboro Veterinary
Hospital

REFERRING VET

Dr. John Sampson

INVOICE

15584

DATE

04/30/26

PRESENTING CLINICAL SIGNS

Presented 4/20 for 2 weeks history of PUPD, diarrhea, and lethargy. PE was unremarkable AUS to look for reasons for hypoalbuminemia

Abnormal PE/Chem/CBC/UA Results: CBC - hct 41.7 CHEM - ALB 1.5, TP 4.5, Ca 7.5 (due to hypoalbuminemia) tt4 - 0.7 UA - usg 1025, no protein

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Left kidney is normal in size (7.69 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Right kidney is normal in size (6.94 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

The adrenal glands were unable to be well visualized in these images.

Spleen

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). Multifocal well-demarcated hyperechoic homogenous nodules are noted. Splenic vasculature appears normal.


Liver

Liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The stomach is moderately distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease. If patient was appropriately fasted, delayed gastric emptying could be considered. Non-shadowing foreign material is considered less likely but cannot be definitively ruled out. If clinical signs are consistent (vomiting, etc.), recommendations include supportive medical care, 24 hours fasting and re-image.



PATIENT

Dallas Thornton

SPECIES

Canine

BREED

Boxer Mix

SEX

Spayed Female

AGE

11

WEIGHT

53

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. John Sampson

HOSPITAL NAME

Richboro Veterinary
Hospital

REFERRING VET

Dr. John Sampson

INVOICE

15584

DATE

04/30/26

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta/chyme. There is no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Pancreas

The area of the pancreas contains irregular hyperechoic pancreatic remodeling.

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

There is no apparent pathologic lymphadenopathy noted in these images.

There is subjectively, diffusely enhanced hyperechoic mesentery and fat throughout these images.

ULTRASONOGRAPHIC FINDINGS

- Hyperechoic pancreas- This finding is suggestive of pancreatic fibrosis, possibly secondary to chronic pancreatitis. A TLI is recommended to rule out exocrine pancreatic insufficiency (EPI), especially if clinical signs (weight loss, diarrhea, etc.) are present.
- Diffusely enhanced/hyperechoic mesentery and fat- consistent with possible diffuse inflammation secondary to bowel disease +/- concurrent chronic low-grade smoldering pancreatitis versus other.
- Hyperechoic splenic nodules- Monitor splenic nodules for changes in size and/or appearance, as myelolipomas typically remain unchanged, or fine needle aspirate of the spleen if patient's coagulation status is appropriate.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS


A baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

A fecal enteropathogen PCR panel to Texas A&M GI Laboratory could be considered for further evaluation of possible infectious disease. Contact lab for recommendations on how long to discontinue antibiotics (if indicated) prior to obtaining a stool sample for submission.

Additionally, a routine fecal/Giardia exam is recommended if not recently evaluated.

In the meantime, supportive/symptomatic medical management of clinical signs is recommended, including a probiotic (such as visbiome or proviable), empirical deworming with a 5-day course of Panacur and, if tolerated, a transition in diet, based on trial-and-error response, beginning possibly with a gastrointestinal biome diet vs a hydrolyzed protein diet vs other. Some patients respond to one brand/version of a hydrolyzed protein diet better than another brand, so several brand attempts may be required.



PATIENT

Dallas Thornton

SPECIES

Canine

BREED

Boxer Mix

SEX

Spayed Female

AGE

11

WEIGHT

53

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. John Sampson

HOSPITAL NAME

Richboro Veterinary
Hospital

REFERRING VET

Dr. John Sampson


INVOICE

15584

DATE

04/30/26

Calcium monitoring and supplementation if necessary is also recommended. Additionally, if patient's coagulation status is otherwise appropriate, antithrombotics such as clopidogrel or low-dose aspirin may also be warranted.



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Beth Johnson, DVM DACVIM

info@sonopath.com