



PATIENT

Willow Turville

SPECIES

Canine

BREED

Cockapoo

SEX

Spayed Female

AGE

10 Years

WEIGHT

10 kg

INTERPRETED BY

Beth Johnson, DVM
 DACVIM

IMAGING PERFORMED BY

Amanda Stewart

HOSPITAL NAME

Aldershot Animal Hospital

REFERRING VET

Dr. Patton

INVOICE

75335

DATE

5/21/26

PRESENTING CLINICAL SIGNS

Urinary leakage. Current Medications: Van-P, Surolan BID AU, TRIZedta

Abnormal PE/Chem/CBC/UA Results: The main abnormality is a marked increase in ALP at 600, with ALT within normal range. In the context of Willow's dilute urine and urinary leakage, this raises concern for possible hormone-related disease such as hyperadrenocorticism/Cushing's disease, although this result is not diagnostic on its own. Other possibilities include liver/gallbladder changes, adrenal changes, steroid/enzyme induction, or age-related hepatobiliary changes. Primary Question to Be Answered in This Exam r/o hyperadrenocorticism

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

Urinary bladder is under distended with primarily anechoic contents and occasional echogenic non-shadowing debris. Apical urinary bladder wall is diffusely thick (0.50 cm). Mucosa is hyperechoic and irregular. The under distended state of the urinary bladder could be in part exacerbating the appearance of the thick, irregular wall appearance. No masses or cystoliths are observed. The trigone and visible pelvic urethra are normal thickness with a smooth mucosal surface.

Kidneys are overall normal in size and shape with smooth peripheral margination. A normal 1:3 cortex to medulla ratio is maintained. The medulla and cortices are uniform in texture with some mild increased cortical echogenicity and mild loss of corticomedullary distinction, expected in this age patient. There is no evidence of pyelectasia, mineral or infarcts observed. Left kidney measured 4.87 cm. Right kidney measured 5.56 cm.

Adrenal Glands

The right adrenal gland is normal in size (1.0 cm at cranial pole and 0.40 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.39 cm at cranial pole and 0.51 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). Multifocal well-demarcated hyperechoic homogenous nodules are noted. Splenic vasculature appears normal.

Liver

Liver is subjectively enlarged with mildly irregular margins. Parenchyma is mildly heterogenous characterized by multiple poorly defined hypoechoic nodules within otherwise hyperechoic liver parenchyma. Visible vasculature and biliary tree appear normal without distension or congestion

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.



PATIENT

Willow Turville

SPECIES

Canine

BREED

Cockapoo

SEX

Spayed Female

AGE

10 Years

WEIGHT

10 kg

INTERPRETED BY

Beth Johnson, DVM
 DACVIM

IMAGING PERFORMED BY

Amanda Stewart

HOSPITAL NAME

Aldershot Animal
 Hospital

REFERRING VET

Dr. Patton

INVOICE

75335

DATE

5/21/26

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen is mildly distended with primarily fluid as well as some echogenic non-shadowing luminal contents and gas consistent with normal chyme. There is no evidence of obstruction, foreign material, or infiltrative disease. Pyloric outflow tract appears patent.

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Pancreas

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

There is no apparent pathologic lymphadenopathy noted in these images.

PRIMARY FINDINGS

- Mildly heterogenous liver – These changes are most consistent with benign processes such as nodular hyperplasia, steroid (vacuolar) hepatopathy, extramedullary hematopoiesis or possibly chronic inflammatory disease and less commonly infiltrative round cell or metastatic neoplasia.

SECONDARY FINDINGS

- Hyperechoic splenic nodules – most consistent with benign myelolipomas. Other differentials such as fibrosis or calcification caused by old hematomas or infarcts, chronic inflammation, granulomatous disease or metastatic disease cannot be ruled out, but are considered less likely.
- Mild to moderate age related kidney changes.
- Possible chronic cystitis - Urinary bladder wall changes are most consistent with chronic cystitis. Infiltrative neoplasia cannot be ruled out but is considered less likely give the location and diffuse nature of the changes.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

If not recently evaluated, a urinalysis and, if indicated based on urinalysis results, urine culture is recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ratio is recommended.

A blood pressure is also recommended.

While hyperadrenocorticism cannot be ruled out based on the appearance of normal adrenal glands,



PATIENT	there is not a definitive ultrasonographically visible intraabdominal explanation for patient's reported urinary leakage. If patient is PU/PD, then that could be exacerbating previously non-clinical urinary incontinence due to urethral sphincter mechanism incompetence (spay incontinence) versus other.
Willow Turville	
SPECIES	Differentials for PU/PD are vast and include, but are not limited to:
Canine	Primary polyuria caused by chronic kidney disease, pyelonephritis, liver disease, diabetes mellitus, hyperthyroidism, hypercalcemia, hyperadrenocorticism, hypoadrenocorticism, E.coli infections ie)
BREED	pyometra in females, polycythemia, central diabetes insipidus or primary nephrogenic diabetes insipidus.
Cockapoo	
SEX	Primary polydipsia caused by psychogenic polydipsia, fever, pain, or central nervous system disease.
Spayed Female	Most causes of PU/PD can be diagnosed with a comprehensive history and physical exam, a first AM urine specific gravity to see if urine concentration is possible (as most animals naturally consume less water overnight) followed by a comprehensive CBC, serum chemistry panel, electrolytes, and urinalysis.
AGE	If not, next step(s) may include a urine culture, low dose dexamethasone suppression test, T4, bile acids, Leptospirosis testing and/or an empirical course of antibiotics.
10 Years	
WEIGHT	If a diagnosis is still not obtained, a more advanced work-up is indicated and consultation with an internist may be warranted.
10 kg	
INTERPRETED BY	Regarding patient's reportedly increased ALP, differentials for a primary cholestatic liver enzyme pattern (increased ALP) are vast and non-specific. Differentials include, but are not limited to, benign nodular hyperplasia which occurs in 70% of older dogs and often does not result in an abnormal ultrasound, reactive or idiopathic/vacuolar hepatopathy, cholestasis and/or hyperadrenocorticism as well as many chronic non-hepatobiliary diseases such as chronic infections/inflammation from dental disease, IBD, neoplasia, hyperlipidemia, hypothyroidism, chronic pancreatitis, chronic stress, etc.
Beth Johnson, DVM DACVIM	
IMAGING PERFORMED BY	<ul style="list-style-type: none"> - Adrenocortical testing such as a low dose dexamethasone suppression test could be considered if clinical signs of hyperadrenocorticism are present. - Ursodiol could be considered if gallbladder sludge is noted as a finding.
Amanda Stewart	
HOSPITAL NAME	<ul style="list-style-type: none"> - A fine needle aspirate of the liver could be considered if patient's coagulation status is appropriate. - Otherwise, recommendations include addressing any other concurrent disease and monitoring. If values are progressive, recheck imaging is recommended.
Aldershot Animal Hospital	
REFERRING VET	
Dr. Patton	


INVOICE

75335

DATE

5/21/26





PATIENT

Willow Turville

SPECIES

Canine

BREED

Cockapoo

SEX

Spayed Female

AGE

10 Years

WEIGHT

10 kg

INTERPRETED BY

Beth Johnson, DVM
 DACVIM

IMAGING PERFORMED BY

Amanda Stewart

HOSPITAL NAME

Aldershot Animal
 Hospital

REFERRING VET


Dr. Patton

INVOICE

75335

DATE

5/21/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
 info@sonopath.com