



PATIENT

Teketa Holmes

SPECIES

Canine

BREED

Bichon Frise

SEX

Spayed Female

AGE

13 Years

WEIGHT

22.8 lbs

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Sara Hansen

HOSPITAL NAME

Forest Valley
Veterinary Clinic

REFERRING VET

Dr. Atkinson

INVOICE

75020

DATE

5/7/26

PRESENTING CLINICAL SIGNS

Bloodwork, and in house ultrasound suspect enlarged liver. Had Animal Sounds ultrasound and found liver nodules, FNA was then requested. ABNORMAL Labwork Values: ALK PHOS 456, Elevated platelets. Current Medications: Clindamycin 75mg, Carprofen 75mg

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is adequately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Kidneys are overall normal in size and shape with smooth peripheral margination. A normal 1:3 cortex to medulla ratio is maintained. The medulla and cortices are uniform in texture with some mild increased cortical echogenicity and mild loss of corticomedullary distinction, expected in this age patient. There is no evidence of pyelectasia, mineral or infarcts observed. Left kidney measures 4.63 cm. Right kidney measures 5.1 cm.

Adrenal Glands

The right adrenal gland is normal in size (0.80 cm at cranial pole and 0.60 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (0.70 cm at cranial pole and 0.60 cm at caudal pole), shape and overall architecture, echogenicity and echotexture. Visible surrounding vasculature appears normal.

Spleen

Spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). Multifocal well-demarcated hyperechoic homogenous nodules are noted. Splenic vasculature appears normal.

Liver

The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. However, in the mid liver, adjacent to the gallbladder is an approximately 2.7 cm x 3.6 cm (or in another view measuring 1.7 cm x 5.2 cm) homogeneous, hypoechoic nodule/mass. Visible vasculature and biliary tree appear normal without distension or congestion.

Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.

Gastrointestinal

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.



PATIENT

Teketa Holmes

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

SPECIES

Canine

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

BREED

Bichon Frise

Pancreas

The pancreas that is observed appears appropriately isoechoic to surrounding omental fat. Visible capsule is smooth and normal in contour. Visible pancreatic parenchyma is homogenous and unremarkable. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

SEX

Spayed Female

Free Abdomen

There is no visible free peritoneal effusion noted in these images.

AGE

13 Years

There is no apparent pathologic lymphadenopathy noted in these images.

WEIGHT

22.8 lbs

PRIMARY FINDINGS

- Mild gallbladder debris - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.
- The appearance of the liver nodule/mass is static with unchanged differentials. Fine needle aspirates were performed at this time of this recheck without ultrasonographically visible evidence of complication.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

SECONDARY FINDINGS

- Hyperechoic splenic nodules – most consistent with benign myelolipomas. Other differentials such as fibrosis or calcification caused by old hematomas or infarcts, chronic inflammation, granulomatous disease or metastatic disease cannot be ruled out, but are considered less likely.
- Mild age related kidney changes.

IMAGING PERFORMED BY

Sara Hansen

HOSPITAL NAME

Forest Valley
Veterinary Clinic

REFERRING VET

Dr. Atkinson

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Further recommendations are largely dependent on results of the fine needle aspirate.

INVOICE

75020

DATE

5/7/26



PATIENT

Teketa Holmes

SPECIES

Canine

BREED

Bichon Frise

SEX

Spayed Female

AGE

13 Years

WEIGHT

22.8 lbs

INTERPRETED BY

Beth Johnson, DVM
 DACVIM

IMAGING PERFORMED BY

Sara Hansen

HOSPITAL NAME

Forest Valley
 Veterinary Clinic

REFERRING VET


Dr. Atkinson

INVOICE

75020

DATE

5/7/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
 info@sonopath.com