



<b>PATIENT</b>	<b>PRESENTING CLINICAL SIGNS</b>
Sprite Dupal-Demers	Elevated liver enzymes. No clinical signs.
<b>SPECIES</b>	Abnormal PE/Chem/CBC/UA Results: Mild ALP 430, moderate ALT elevation 434, mild AST elevation 148.
Canine	
<b>BREED</b>	<b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>
Bearded Collie	<b>Urinary System</b>
<b>SEX</b>	Urinary bladder is adequately distended with primarily anechoic contents and occasional echogenic non-shadowing debris. Apical urinary bladder wall is diffusely thick (0.34 cm thick). Mucosa is hyperechoic and irregular. No masses or cystoliths are observed. The trigone and visible pelvic urethra are normal thickness with a smooth mucosal surface.
Spayed Female	
<b>AGE</b>	The right kidney is normal in size (4.95 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A hyperechoic band parallel to the corticomedullary border is present.
13 Years	
<b>WEIGHT</b>	The left kidney is normal in size (4.49 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A hyperechoic band parallel to the corticomedullary border is present.
12.4 kg	
<b>INTERPRETED BY</b>	<b>Adrenal Glands</b>
Beth Johnson, DVM DACVIM	The right adrenal gland is normal in size (0.77 cm at the cranial pole and 0.38 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.
<b>IMAGING PERFORMED BY</b>	The left adrenal gland is normal in size (0.40 cm at the cranial pole and 0.50 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.
Dr. Sarah Barthelémy	
<b>HOSPITAL NAME</b>	<b>Spleen</b>
South Pointe Pet Hospital	The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.
<b>REFERRING VET</b>	<b>Liver</b>
Dr. James	The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.
<b>INVOICE</b>	
43538	
<b>DATE</b>	Gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.
6/28/23	<b>Gastrointestinal</b>
	The visible stomach wall is normal in thickness and layering. The lumen of the stomach is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta.



<b>PATIENT</b>	There is no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.
Sprite Dupal-Demers	
<b>SPECIES</b>	The visible small intestines are normal in wall thickness and layering (canine duodenum < 0.5 cm and feline duodenum < 0.4 cm; other < 0.3 cm). Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.
Canine	
<b>BREED</b>	The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.
Bearded Collie	
	<b><i>Pancreas</i></b>
<b>SEX</b>	The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.
Spayed Female	
<b>AGE</b>	<b><i>Free Abdomen</i></b>
13 Years	There is no evidence of free peritoneal effusion noted in these images.
<b>WEIGHT</b>	There is no apparent lymphadenopathy noted in these images.
12.4 kg	
<b>INTERPRETED BY</b>	
Beth Johnson, DVM DACVIM	<ul style="list-style-type: none"> <li>• <b>Chronic Cystitis</b> - Urinary bladder wall changes are most consistent with chronic cystitis. Infiltrative neoplasia cannot be ruled out but is considered less likely give the location and diffuse nature of the changes.</li> <li>• <b>Bilateral medullary rim sign</b> - This finding is of unknown clinical significance and can be a normal variant, often idiopathic. Medullary rim sign can be present with renal disease including FIP, lymphoma, hypercalcemic nephropathy, Leptospirosis, tubular disease, other and should be interpreted in combination with other more specific indications of kidney disease such as isosthenuria, proteinuria, azotemia, etc. This is a common incidental finding in patients with diabetes mellitus.</li> </ul>
<b>IMAGING PERFORMED BY</b>	
Dr. Sarah Barthelmy	
<b>HOSPITAL NAME</b>	<ul style="list-style-type: none"> <li>• <b>Mild gallbladder debris</b> - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness. Cholecystic debris is not necessarily related to hepatobiliary disease. Echogenic bile is most commonly an incidental finding in dogs and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.</li> </ul>
South Pointe Pet Hospital	
<b>REFERRING VET</b>	<ul style="list-style-type: none"> <li>• An obvious cause for the reported increased liver enzymes is not identified in these images. Microscopic disease such as Leptospirosis, bacterial cholangiohepatitis, chronic active hepatitis, copper-associated hepatotoxicity, other hepatotoxicity, infiltrative neoplasia (considered unlikely), etc. cannot be definitively ruled out.</li> </ul>
Dr. James	
<b>INVOICE</b>	
43538	
<b>DATE</b>	
6/28/23	

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Recommendations include an "antigen search" for sources of reactive hepatopathy (including testing for Leptospirosis), followed by a course of empirical antibiotics and hepatic nutraceuticals, with monitoring of ALT for improvement. If improvement is not noted and/or enzyme increase progresses, a liver biopsy may be warranted.



**PATIENT**

Sprite Dupal-Demers

Additionally, if not recently evaluated, a urinalysis and, if indicated based on urinalysis results, urine culture are recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ratio is recommended.

**SPECIES**

Canine

**BREED**

Bearded Collie

**SEX**

Spayed Female

**AGE**

13 Years

**WEIGHT**

12.4 kg

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Sarah Barthelemy

**HOSPITAL NAME**

South Pointe Pet  
Hospital

**REFERRING VET**


Dr. James

**INVOICE**

43538

**DATE**

6/28/23





**PATIENT**

Sprite Dupal-Demers

**SPECIES**

Canine

**BREED**

Bearded Collie

**SEX**

Spayed Female

**AGE**

13 Years

**WEIGHT**

12.4 kg

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Sarah Barthelemy

**HOSPITAL NAME**

South Pointe Pet  
Hospital

**REFERRING VET**

Dr. James

**INVOICE**

43538

**DATE**

6/28/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
info@sonopath.com