



PATIENT PRESENTING CLINICAL SIGNS

Luna Iruegas Weight loss. Current meds: Mirataz Transdermal Ointment

SPECIES ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Feline Urinary System

BREED Urinary bladder is moderately distended. It has a normal uniform wall thickness (<0.2 cm). Contents include primarily anechoic fluid combined with suspended echogenic non-shadowing debris within the fluid. Dependent mineral density with acoustic shadowing is noted, consistent with a cystolith, measuring approximately 0.70 cm. No masses noted. The trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

SEX The right kidney is normal in size (3.86 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Spayed Female

AGE The left kidney is normal in size (3.8 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

6 Years

Adrenal Glands

WEIGHT The right adrenal gland is normal in size (0.36 cm), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

11 Pounds

The left adrenal gland is normal in size (0.35 cm), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

Spleen

The spleen is generally normal in size and shape with a smooth capsular contour. Parenchyma is diffusely nodular in appearance characterized by small discrete hypoechoic nodules. Splenic vasculature appears normal.

IMAGING PERFORMED BY

Jessica Miller

Liver

The liver is subjectively enlarged (swollen contour). Mild parenchymal remodeling with diffusely mildly coarse architecture and increased portal markings is present. No focal nodules or masses are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

HOSPITAL NAME

Willowbrook AC

The gallbladder is moderately distended with anechoic bile as well as suspended and gravity dependent echogenic debris. The wall is smooth without visible thickening. There is no evidence of cystic or CBD dilation. There is no evidence of effusion or inflammation.

REFERRING VET

Dr. Palescandolo

Gastrointestinal

INVOICE

39101

The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is empty with no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

DATE

6/29/22

The visible small intestine demonstrates areas of thick muscularis layer relative to mucosa (disruption of the normal 1:3 muscularis:mucosa ratio). Small intestinal submucosa is slightly irregular, thick and hyperechoic, without evident loss of layering appreciated.



PATIENT	The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.
Luna Iruegas	
	Pancreas
SPECIES	The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.
Feline	
	Free Abdomen
BREED	A scant amount of anechoic free fluid is noted between liver lobes.
DSH	
	Mesenteric lymph nodes are enlarged with swollen irregular capsular contour and loss of normal length to width ratio (rounded in shape). Nodes are hypoechoic with loss of normal parenchymal detail.
SEX	
Spayed Female	
	PRIMARY FINDINGS
AGE	<ul style="list-style-type: none"> Hypoechoic hepatomegaly – This appearance is consistent with an acute hepatopathy or acute cholangiohepatitis. Infiltrative neoplasia (round cell neoplasia) should also be considered. Splenic micronodular hyperplasia – This nodular change is often associated with benign aging nodular hyperplasia. Infiltrative neoplasia, however, including both early hemangiosarcoma as well as round cell neoplasia cannot be ruled out. Aggressive mesenteric lymph nodes – most consistent with infiltrative round cell or metastatic neoplasia. A benign aggressive inflammatory response cannot be ruled out without tissue sampling +/- culture. Inflammatory bowel disease (IBD) pattern - This finding has been reported with infiltrative bowel disease including both benign inflammatory disease as well as infiltrative neoplasia such as lymphoma.
6 Years	
WEIGHT	
11 Pounds	
INTERPRETED BY	
Beth Johnson, DVM DACVIM	
	SECONDARY FINDINGS
IMAGING PERFORMED BY	<ul style="list-style-type: none"> Urinary bladder debris with cystoliths noted. Gallbladder debris - Cholecystic debris is of unknown clinical significance. It can be seen with biliary stasis from fasting or illness, however, it can also be associated with hepatobiliary disease in cats and should be interpreted in combination with clinical signs such as nausea, inappetence, cranial abdominal discomfort and/or laboratory changes such as increased ALP and/or increased Tbili.
Jessica Miller	
HOSPITAL NAME	
Willowbrook AC	
REFERRING VET	INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS
Dr. Palescandolo	Given the diffuse changes involving the bowel, liver, potentially the spleen, and lymph nodes, infiltrative round cell neoplasia such as lymphoma or less likely mast cell tumor is a concern. Benign inflammatory bowel disease with reactive lymph nodes, etc. cannot be ruled out, but is considered less likely. Recommendations include:
INVOICE	<ul style="list-style-type: none"> Fine needle aspirate of the liver, spleen, and enlarged lymph nodes, if patient's coagulation status is appropriate.
39101	
DATE	
6/29/22	



PATIENT

Luna Iruegas

- A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

SPECIES

Feline

- If round cell neoplasia is not diagnosed via cytology, biopsies of the GI tract, being sure to include ileum, if possible, may be necessary to definitively diagnose and therefore manage the suspected infiltrative bowel disease as a cause for weight loss.


BREED

DSH

- Urinalysis and, if indicated based on urinalysis results, urine culture are recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ration is recommended.

SEX

Spayed Female




AGE

6 Years

WEIGHT

11 Pounds



INTERPRETED BY


Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

Willowbrook AC



REFERRING VET

Dr. Palescandolo

INVOICE

39101

DATE

6/29/22



PATIENT

Luna Iruegas

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

6 Years

WEIGHT

11 Pounds

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

Willowbrook AC

REFERRING VET



Dr. Palescandolo

INVOICE

39101

DATE

6/29/22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
Beth.Johnson@sonopath.com