



PATIENT PRESENTING CLINICAL SIGNS

Finn Reiss Intermittent vomiting and inappetence, drinks large amount of pool salt water, weight loss.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

SPECIES *Urinary System*

Canine The urinary bladder is moderately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

BREED

Irish Setter Prostate is normal in size, echotexture and echogenicity for a neutered male.

SEX

Neutered Male The right kidney is normal in size (8.19 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

AGE

2018 The left kidney is normal in size (8.49 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

WEIGHT

103.8 **Adrenal Glands**
 Adrenal glands are small (flattened contour). Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal. The right adrenal gland measures 0.87 cm at the cranial pole and 0.46 cm at the caudal pole. The left adrenal gland measures 0.49 cm at the cranial pole and 0.38 cm at the caudal pole.

Spleen

INTERPRETED BY

Beth Johnson, DVM DACVIM Spleen is generally normal in size and shape with a smooth capsular contour. Parenchyma is diffusely nodular in appearance characterized by small discrete hypoechoic nodules. Splenic vasculature appears normal.

IMAGING PERFORMED BY

Liver

Rebekah Jakum, CVT ARDMS/RVT The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

HOSPITAL NAME

Stanglein Vet Clinic The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.

REFERRING VET

Gastrointestinal

Dr. Green The visible stomach wall is normal in thickness and layering. The lumen of the stomach is mildly distended with very echogenic reverberation artifact from intraluminal gas. There is no evidence of obstruction, foreign material or infiltrative disease; however, complete visualization of far wall is partially inhibited by gas. Pyloric outflow tract appears patent.

INVOICE

43835

DATE

7/13/23

The visible small intestines are normal in wall thickness and layering (canine duodenum < 0.5 cm and feline duodenum < 0.4 cm; other < 0.3 cm). Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.



PATIENT The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

Finn Reiss

Pancreas

SPECIES The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

Canine

Free Abdomen

BREED

There is no evidence of free peritoneal effusion noted in these images.

Irish Setter

There is no apparent lymphadenopathy noted in these images.

SEX

ULTRASONOGRAPHIC FINDINGS

Neutered Male

- **Flat adrenal glands** – This can be a normal patient variant and/or a sign of exogenous cortisol administration. If exogenous steroids are not being administered, hypoadrenocorticism (either relative or absolute) should be considered.

AGE

- **Splenic micronodular hyperplasia pattern** – This nodular change is often associated with benign aging nodular hyperplasia. Infiltrative neoplasia, however, including both early hemangiosarcoma as well as round cell neoplasia cannot be ruled out.

2018

WEIGHT

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

103.8

If not recently evaluated, a general metabolic health screen is recommended beginning with a CBC/Chem panel, electrolytes, a urinalysis and, if indicated based on urinalysis results, urine culture. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ratio is recommended.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

A baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.

IMAGING PERFORMED BY

Pending results of the above, a fine needle aspirate of the spleen could be considered if patient's coagulation status is appropriate.

Rebekah Jakum, CVT
ARDMS/RVT

Additionally, again pending the above results, a gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

HOSPITAL NAME

Stanglein Vet Clinic

In the meantime, empirical deworming with a 5-day course of Panacur is recommended, as is an empirical course of therapy for helicobacter. Additionally, a transition in diet could be considered based on trial and error response, with some trial options including a bland easy to digest or low-fat diet, or potentially a hydrolyzed protein diet, etc.

REFERRING VET

Dr. Green

INVOICE

43835

DATE

7/13/23



PATIENT

Finn Reiss

SPECIES

Canine

BREED

Irish Setter

SEX

Neutered Male

AGE

2018

WEIGHT

103.8

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Rebekah Jakum, CVT
ARDMS/RVT

HOSPITAL NAME

Stanglein Vet Clinic

REFERRING VET


Dr. Green

INVOICE

43835

DATE

7/13/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
info@sonopath.com