



<b>PATIENT</b>	<b>PRESENTING CLINICAL SIGNS</b>
Peaches Morrell	Increased LES found on wellness lab work ALT 404 (previously 177), alkp 249 (previously normal) Abnormal PE/Chem/CBC/UA Results: NR PE, urine pending, rest of mini chem NR. no clinical signs reported at home. no azotemia.
<b>SPECIES</b>	
Canine	<b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>
<b>BREED</b>	<b>Urinary System</b>
Labradoodle	Urinary bladder is adequately distended. It has a normal uniform wall thickness. Contents include primarily anechoic fluid with a large amount of echogenic non-shadowing debris, most consistent with exfoliated cells, mucous and/or small blood clots. Both sterile inflammation as well as urinary tract infection can also present with echogenic debris. No masses or cystoliths are observed. The trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.
<b>SEX</b>	
Spayed Female	The right kidney is normal in size (8.1 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.
<b>AGE</b>	
6.5 Years	The left kidney measures 4.9 cm but is measured in an oblique view and suspected to be of normal size, shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.
<b>WEIGHT</b>	
81.5	
<b>INTERPRETED BY</b>	<b>Adrenal Glands</b>
Beth Johnson, DVM DACVIM	The caudal pole of the right adrenal gland is normal in size (0.80 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal. The cranial pole is unable to be well visualized in these images.
<b>IMAGING PERFORMED BY</b>	
Dr. Arms	The caudal pole of the left adrenal gland is normal in size (0.58 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal. The cranial pole is unable to be well visualized in these images.
<b>HOSPITAL NAME</b>	<b>Spleen</b>
Gilberstville VH	The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). A 1.6 cm round, mildly heterogeneous, primarily hypoechoic nodule is noted in the mid spleen, resulting in mild capsular bulge. Splenic vasculature appears normal.
<b>REFERRING VET</b>	<b>Liver</b>
Dr. Arms	Liver is subjectively enlarged with mildly irregular margins. Parenchyma is heterogenous characterized by multiple poorly defined hypoechoic nodules within otherwise hyperechoic liver parenchyma. Visible vasculature and biliary tree appear normal without distension or congestion.
<b>INVOICE</b>	
43801	
<b>DATE</b>	
7/5/23	The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.



**PATIENT** *Gastrointestinal*

Peaches Morrell

The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is mildly distended with fluid, as well as echogenic nonshadowing luminal contents and gas consistent with normal ingesta/chyme. There is no evidence of obstruction, foreign material or infiltrative disease. Pyloric outflow tract appears patent.

**SPECIES**

Canine

**BREED**

Labradoodle

The visible small intestines are normal in wall thickness and layering (canine duodenum < 0.5 cm and feline duodenum < 0.4 cm; other < 0.3 cm). Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

**SEX**

Spayed Female

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

**AGE**

6.5 Years

**Pancreas**

The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

**WEIGHT**

81.5

**Free Abdomen**

There is no evidence of free peritoneal effusion noted in these images.

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

There is no apparent lymphadenopathy noted in these images.

**ULTRASONOGRAPHIC FINDINGS**

**IMAGING PERFORMED BY**

Dr. Arms

- **Heterogenous Liver** – These changes are most consistent with benign processes such as nodular hyperplasia, steroid (vacuolar) hepatopathy, extramedullary hematopoiesis or possibly chronic inflammatory disease and less commonly infiltrative round cell or metastatic neoplasia.
- **Hypo to anechoic splenic nodule** – likely represents a benign lesion such as a cyst, hematoma, nodular hyperplasia, extramedullary hematopoiesis, etc., however while considered less likely, infiltrative neoplasia can mimic benign lesions, and cannot be ruled out.
- Urinary bladder debris

**HOSPITAL NAME**

Gilberstville VH

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

**REFERRING VET**

Dr. Arms

Recommendations include an “antigen search” for sources of reactive hepatopathy (including testing for Leptospirosis), followed by a course of empirical antibiotics and hepatic nutraceuticals, with monitoring of ALT for improvement. If improvement is noted, antibiotics should be continued until liver enzymes either normalize or plateau (recheck every 2-3 weeks); however, if improvement is not noted and/or enzyme increase progresses, antibiotics should not be continued long term and sampling, beginning with a FNA of the liver if patient’s coagulation status is appropriate or progressing to a liver biopsy (including copper level assessment) may ultimately be warranted.

**INVOICE**

43801

**DATE**

7/5/23

Additionally, a fine needle aspirate of the splenic nodule could be considered if patient’s coagulation status is appropriate.



**PATIENT**

Peaches Morrell

If not recently evaluated, a urinalysis and, if indicated based on urinalysis results, urine culture are recommended. If protein is present in an otherwise quiet sediment, protein quantification with a urine protein to creatinine ration is recommended.

**SPECIES**

Canine

The liver enzyme increases are an atypical pattern for typical hyperadrenocorticism. Additionally, testing for hyperadrenocorticism is typically not recommended in an asymptomatic patient (i.e., no PU/PD, etc.).

**BREED**

Labradoodle

Therefore, in this patient, the pending low-dose Dexamethasone suppression test should be interpreted cautiously to avoid overinterpreting a false positive result.

**SEX**

Spayed Female

**AGE**

6.5 Years

**WEIGHT**

81.5

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Arms

**HOSPITAL NAME**

Gilberstville VH

**REFERRING VET**


Dr. Arms

**INVOICE**

43801

**DATE**

7/5/23





**PATIENT**

Peaches Morrell

**SPECIES**

Canine

**BREED**

Labradoodle

**SEX**

Spayed Female

**AGE**

6.5 Years

**WEIGHT**

81.5

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Dr. Arms

**HOSPITAL NAME**

Gilberstville VH

**REFERRING VET**


Dr. Arms

**INVOICE**

43801

**DATE**

7/5/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
info@sonopath.com