



**PATIENT PRESENTING CLINICAL SIGNS**

Paolo Venturini Intermittent v+, decreased appetite and diarrhea over last several months. Current meds: Intermittent cerenia, pepsid, metronidazole

**SPECIES**

Abnormal PE/Chem/CBC/UA Results: Lymph 4838, Mono 1416

Canine

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**BREED**

**Urinary System**

Whoodle

The urinary bladder is moderately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

**SEX**

Neutered Male

The right kidney is normal in size (4.68 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A hyperechoic band parallel to the corticomedullary border is present.

**AGE**

2 Years

The left kidney is normal in size (4.89 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A hyperechoic band parallel to the corticomedullary border is present.

**WEIGHT**

24 Pounds

**INTERPRETED BY**

**Adrenal Glands**

Beth Johnson, DVM  
DACVIM

The right adrenal gland is normal in size (1.0 cm at the cranial pole and 0.46 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

**IMAGING PERFORMED BY**

The left adrenal gland is normal in size (0.42 cm at the cranial pole and 0.59 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

Jessica Miller, RDMS

**Spleen**

**HOSPITAL NAME**

The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.

The Venturing Vet

**REFERRING VET**

**Liver**

Dr. Herzog

The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

**INVOICE**

44525

**DATE**

8/1/23

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.



**PATIENT**

***Gastrointestinal***

Paolo Venturini

The stomach wall is normal in thickness (canine < 0.5 cm and feline < 0.4 cm) and layering. The lumen of the stomach is mildly distended with a moderate amount of echogenic material, some of which demonstrates distally progressive acoustic shadow.

**SPECIES**

Canine

The visible small intestines are normal in wall thickness and layering. Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is mildly distended with echogenic non-shadowing luminal contents and gas consistent with normal ingesta. There is no evidence of obstruction, foreign material or infiltrative disease.

**BREED**

Whoodle

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

**SEX**

Neutered Male

***Pancreas***

The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

**AGE**

2 Years

***Free Abdomen***

**WEIGHT**

24 Pounds

There is no evidence of free peritoneal effusion noted in these images.

There is no apparent lymphadenopathy noted in these images.

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**ULTRASONOGRAPHIC FINDINGS**

- **Bilateral medullary rim sign** - This finding is of unknown clinical significance and can be a normal variant, often idiopathic. Medullary rim sign can be present with renal disease including FIP, lymphoma, hypercalcemic nephropathy, Leptospirosis, tubular disease, other and should be interpreted in combination with other more specific indications of kidney disease such as isosthenuria, proteinuria, azotemia, etc. This is a common incidental finding in patients with diabetes mellitus.
- The appearance of the gastrointestinal tract in these images is most consistent with a post-prandial abdomen with ingesta and gas in the stomach. However, given the acoustic shadowing in the stomach in some images, partially obstructive or non-obstructive foreign material cannot be definitively ruled out.

**IMAGING PERFORMED BY**

Jessica Miller, RDMS

**HOSPITAL NAME**

The Venturing Vet

**REFERRING VET**

Dr. Herzog

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Given this patient's clinical signs and reported lymphocytosis, a baseline cortisol is recommended. If baseline cortisol is less than 2, a full ACTH stimulation test is recommended to rule out hypoadrenocorticism.

**INVOICE**

44525

A gastrointestinal malabsorption panel (including cobalamin, folate, TLI and PLI) to Texas A&M GI Laboratory is recommended for further evaluation of GI and pancreatic function.

**DATE**

8/1/23

A fecal enteropathogen PCR panel to Texas A&M GI Laboratory could be considered for further evaluation of possible infectious disease.



**PATIENT**

Contact lab for recommendations on how long to discontinue Metronidazole prior to obtaining a stool sample for submission.

Paolo Venturini

**SPECIES**

In the meantime, while awaiting results, supportive/symptomatic medical management is recommended in the form of antiemetics, gastroprotectants, a probiotic such as Visbiome or Provable, and empirical deworming with a 5-day course of Panacur, as well as fluid therapy, etc. if clinically indicated.

Canine

**BREED**

Additionally, an additional 12-24 hours of fasting followed by recheck imaging of the GI tract could be considered to further evaluate gastric contents, especially if empirical management is instituted and vomiting persists.

Whoodle

**SEX**

Neutered Male

**AGE**

2 Years

**WEIGHT**

24 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Jessica Miller, RDMS

**HOSPITAL NAME**

The Venturing Vet

**REFERRING VET**

Dr. Herzog

**INVOICE**

44525

**DATE**

8/1/23





**PATIENT**

Paolo Venturini

**SPECIES**

Canine

**BREED**

Whoodle

**SEX**

Neutered Male

**AGE**


2 Years

**WEIGHT**

24 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM



**IMAGING PERFORMED BY**

Jessica Miller, RDMS

**HOSPITAL NAME**

The Venturing Vet

**REFERRING VET**

Dr. Herzog

**INVOICE**

44525

**DATE**

8/1/23

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Beth Johnson, DVM, DACVIM**  
info@sonopath.com