



PATIENT

Duke LaPenta

SPECIES

Canine

BREED

American Bulldog

SEX

Neutered Male

AGE

11 Years

WEIGHT

89 Pounds

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Arch Gordon

HOSPITAL NAME

Coral Ridge AH

REFERRING VET

Dr. Arch Gordon

INVOICE

41321

DATE

9/14/22

PRESENTING CLINICAL SIGNS

Panting/Painful? Improved after Sucralfate administered Abdomen slight distention- no palpable pain
Mild liver enzyme elevation after CCS administered for acute moist dermatitis in June- now normal
Multiple areas of spondylosis including LS JCT noted on radiographs

Abnormal PE/Chem/CBC/UA Results: T4 low - 0.7 - 1.0- 4.0 -full profile pending Amylase low at 202 (337-1469) urine protein 1 +

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic contents. No masses, inflammatory changes, echogenic sediment or cystoliths are observed. The urinary bladder, trigone and visible pelvic urethra are normal in thickness with a smooth mucosal surface.

Prostate is normal in size, echotexture and echogenicity for a neutered male.

The right kidney is normal in size (6.84 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed. A small cortical cyst is noted.

The left kidney is normal in size (6.47 cm), shape and echogenicity. It has smooth peripheral margination. There is a normal 1:3 cortex to medulla ratio with appropriate corticomedullary distinction. There is no evidence of pyelectasia, mineral or infarcts observed.

Adrenal Glands

The right adrenal gland is normal in size (1.96 cm long x 1.0 cm at the cranial pole and 0.54 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

The left adrenal gland is normal in size (1.56 cm long x 0.37 cm at the cranial pole and 0.53 cm at the caudal pole), shape and contour. Corticomedullary structure is unremarkable. Visible surrounding vasculature appears normal.

Spleen

The spleen is subjectively normal in size with a normal smooth capsular contour. Parenchyma is appropriately finely textured and homogenous with normal echogenicity relative to surrounding tissue (hyperechoic to liver). No focal nodules or masses are observed. Splenic vasculature appears normal.

Liver

The liver is subjectively normal in size with normal smooth curvilinear peripheral contour. Parenchyma is appropriately hypoechoic to the spleen in echogenicity and appropriately mildly coarse and homogenous in echotexture. No focal lesions are observed. Visible vasculature and biliary tree appear normal without distension or congestion.

The gallbladder is non-distended in size. The wall is smooth without visible thickening. Luminal contents are primarily anechoic. There is no evidence of cystic or common bile duct dilation.



PATIENT

Gastrointestinal

Duke LaPenta

The visible stomach wall is normal in thickness and layering. The lumen of the stomach is mildly distended with very echogenic reverberation artifact from intraluminal gas. There is no evidence of obstruction, foreign material or infiltrative disease; however, complete visualization of far wall is partially inhibited by gas. Pyloric outflow tract appears patent.

SPECIES

Canine

The visible small intestines are normal in wall thickness and layering (canine duodenum < 0.5 cm and feline duodenum < 0.4 cm; other < 0.3 cm). Small intestinal motility appears adequate (1-3 contractions per min). The lumen of the small intestine is empty with no evidence of obstruction, foreign material or infiltrative disease.

BREED

American Bulldog

The visible colon is normal in wall thickness (< 0.2 cm) and layering. Contents are consistent with normal formed feces and gas.

SEX

Neutered Male

Pancreas

The pancreatic parenchyma is appropriately isoechoic to surrounding tissue. Visible capsule is smooth and normal in contour. There is no visible pancreatic duct dilation. There is no evidence of active peripancreatic inflammation.

AGE

11 Years

Free Abdomen

There is no evidence of free peritoneal effusion noted in these images.

WEIGHT

89 Pounds

There is no apparent lymphadenopathy noted in these images.

ULTRASONOGRAPHIC FINDINGS

- Unremarkable/normal abdomen without evident pathology to explain this patient's reported panting/pain.

INTERPRETED BY

Beth Johnson, DVM
DACVIM

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Recommendations are to follow up with suspected spinal pain/spinal lesions to see if managing that pain helps to control the panting, and if not, further evaluation of the cardiopulmonary system in the form of radiographs +/- an echocardiogram may be warranted. Blood pressure is also recommended if not recently evaluated.

IMAGING PERFORMED BY

Dr. Arch Gordon

HOSPITAL NAME

Coral Ridge AH

REFERRING VET


Dr. Arch Gordon

INVOICE

41321

DATE

9/14/22





PATIENT

Duke LaPenta

SPECIES

Canine

BREED

American Bulldog

SEX

Neutered Male

AGE

11 Years

WEIGHT

89 Pounds

INTERPRETED BY

Beth Johnson, DVM
DACVIM

IMAGING PERFORMED BY

Dr. Arch Gordon

HOSPITAL NAME

Coral Ridge AH

REFERRING VET


Dr. Arch Gordon

INVOICE

41321

DATE

9/14/22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Beth Johnson, DVM, DACVIM
Beth.Johnson@sonopath.com