



PATIENT

Maggie Larder

SPECIES

Canine

BREED

Labradoodle

SEX

Spayed female

AGE

5 years

WEIGHT

17 kg

INTERPRETED BY

Dr Brittany Sinclair,
BVSc(hons), DACVECC

IMAGING PERFORMED BY

Dr. Belan

HOSPITAL NAME

Healing Traditions VC

REFERRING VET

Dr. Vockeroth

INVOICE

43107

DATE

3/6/23

PRESENTING CLINICAL SIGNS

History: Intermittent vomiting for last 11 months Has been on multiple food residences as well as antibiotics, probiotics
Abnormal PE/Chem/CBC/UA Results: Non diagnostic blood work but Heliobacter was cultured from the stool.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder, trigone, and visible pelvic urethra were of normal thickness. The ureters were not visible which is normal. There was normal wall layering with no masses, uroliths or abnormal thickening visualized. Urine was anechoic. No evidence of inflammatory or neoplastic changes were noted.

The kidneys have a smooth capsule and with mild hazing of corticomedullary definition with approximate maintenance of normal ratio (cortex 1/3 of medulla). No evidence of pelvic dilation was present. Left kidney contains spherical anechoic fluid accumulation consistent with cortical cyst. The left kidney measured 4.74 cm and the right kidney measured 4.84 cm.

Adrenal Glands

Both adrenal glands were visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were unremarkable. Capsule, cortex, and medullary definition were normal for this age patient. The left adrenal gland measured 2.51 cm in length and 0.48 cm at the cranial pole and 0.41 cm at the caudal pole. The right adrenal gland measured 2.18 cm in length and 0.81 cm at the cranial pole and 0.41 cm at the caudal pole.

Spleen


The spleen was normal with a smooth homogeneous parenchyma hyperechoic to liver and renal cortical parenchyma and smooth capsule, with normal splenic vasculature with no signs of congestion or thrombosis. No sonographic evidence of acute or chronic inflammatory, neoplastic, or infarct changes were noted.

Liver

The liver is subjectively normal in size with normal contours and structure. There is appropriate echogenicity and echotexture. No overt structural evidence of inflammatory, infiltrative or regenerative pathology is evident. Vascular and biliary tracts are of normal volume with no evidence of congestion. No pathological hepatic lymphadenopathy observed. The gall bladder is moderately distended with anechoic fluid, with hyperechoic non-shadowing debris present. There is no surrounding free fluid or signs of active inflammation.

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and



PATIENT

Maggie Larder

SPECIES

Canine

BREED

Labradoodle

SEX

Spayed female

AGE

5 years

WEIGHT

17 kg

INTERPRETED BY

Dr Brittany Sinclair,
BVSc(hons), DACVECC

IMAGING PERFORMED BY

Dr. Belan

HOSPITAL NAME

Healing Traditions VC

REFERRING VET

Dr. Vockeroth

INVOICE

43107

DATE

3/6/23

there is no impression of reduced peristaltic activity. No masses or focal lesions were observed. The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed. The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The base and limbs of the pancreas were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour and parenchyma were normal. No overt evidence of active inflammatory or neoplastic disease was noted.

Lymph Nodes

No clinically significant lymphadenopathy or abnormalities noted.

Free Abdomen

No masses or free fluid were noted.

ULTRASONOGRAPHIC FINDINGS

Primary Findings

1. Normal GI tract
2. Normal pancreas
3. Left renal cyst, mild bilateral degenerative renal changes
4. Gall bladder debris

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is no ultrasonographically evident cause of reported vomiting, decreased appetite and weight loss in this abdominal study. Pancreas and GI tract are within normal limits. Consideration for dietary indiscretion, food sensitivity/allergy or mild inflammatory bowel disease is reasonable. While not sonographically evident, pancreatitis cannot be completely ruled out. Empiric treatment for GI signs including anti-nausea, appetite stimulant and fluid support as clinically indicated is warranted. A diet trial with hydrolyzed protein or select protein diet could be considered if food sensitivity is suspected clinically. If signs are persistent or recurrent, additional diagnostics to be considered include GI panel (TLI/PLI/cobalamin/folate), baseline cortisol +/- ACTH stimulation test, fecal pathogen panel, thyroid testing, bile acid profile, and thoracic radiographs to rule out occult neoplasia, cardiac disease and esophageal disease as potential causes. Ultimately GI biopsy may be required for more definitive diagnosis if the patient is not responsive to medical treatment.

Renal changes are mild and are likely age related degenerative changes. Correlate clinical significance with blood work/urinalysis findings and clinical signs.



PATIENT

Maggie Larder

Gall bladder debris is likely an incidental finding and is often subclinical and often does not warrant specific treatment or further investigation. Correlate clinical significance with bloodwork findings and clinical signs. Serial imaging for monitoring could be considered especially if liver enzymes subsequently become elevated. If otherwise clinically indicated, investigation for endocrinopathy such as hyperadrenocorticism or hypothyroidism could be considered as an underlying cause predisposing to gall bladder debris accumulation.

SPECIES

Canine

BREED

Labradoodle

SEX

Spayed female

AGE

5 years

WEIGHT

17 kg

INTERPRETED BY

Dr Brittany Sinclair,
BVSc(hons), DACVECC

IMAGING PERFORMED BY



Dr. Belan

HOSPITAL NAME

Healing Traditions VC

REFERRING VET

Dr. Vockeroth




INVOICE

43107

DATE

3/6/23



PATIENT

Maggie Larder

SPECIES

Canine

BREED

Labradoodle

SEX

Spayed female

AGE

5 years

WEIGHT

17 kg

INTERPRETED BY

Dr Brittany Sinclair,
BVSc(hons), DACVECC

**IMAGING
PERFORMED BY**

Dr. Belan

HOSPITAL NAME

Healing Traditions VC

REFERRING VET


Dr. Vockeroth

INVOICE

43107

DATE

3/6/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Dr Brittany Sinclair, BVSc(hons), DACVECC
info@SonoPath.com