

PATIENT

Chabot Coombs

SPECIES

Canine

BREED

Labrador Retriever Mix

SEX

Neutered Male

AGE

13 Years

WEIGHT

88.5 pounds

INTERPRETED BY

Eric Lindquist, DMV,
DABVP(CFM), Cert.
IVUSS

IMAGING PERFORMED BY

Dr. Chrissy Krell DVM

HOSPITAL NAME

Taylor Emergency
Hospital

REFERRING VET

Dr. Chrissy Krell DVM

INVOICE

13099

DATE

01/12/26

PRESENTING CLINICAL SIGNS

Acute onset of diarrhea became very hemorrhagic over the past 72 hours. History of a sarcoma being removed from his lip in December. Patient is having significant difficulty walking today. Did eat some homemade mac and cheese with garlic prior to the onset of the GI upset.

PE: patient paretic, prefers to lay (difficulty standing). Passed notable yellow brown diarrhea with blood clots, abdomen slightly distended and mildly uncomfortable. HR 130 bpm. Noted multiple SQ masses over the chest and abdomen. CBC: mild non-regenerative anemia (likely historic) Chem: SDMA 16 TT4: 0.8 UA pending

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The **urinary bladder** presented a relatively uniform thickening of the cranioventral and craniodorsal mucosae with micropolypoid mucosal changes without involvement of the submucosae. The urine presented some echogenicity consistent with a minor amount of sand. No evidence of urethral pathology was present. This presentation is most consistent with chronic cystitis. Technically transitional cell carcinoma cannot be ruled out without histopathological review but is not overtly suspected based on this pattern. Cystocentesis and urine culture +/- pathological review of urine cytology would be warranted. No overt calculi were present at this time. The pelvic urethra was normal in structure and tone to a depth of 2.0 cm.

The **kidneys** revealed largely normal size and structure, corticomedullary definition and ratio (cortex 1/3 of medulla) were essentially maintained with some minor age-related loss of curvilinear patterns regarding the capsule and C/M junction. The cortices presented largely uniform texture with some increased echogenicity expected for his age patient. Medullary structure differed distinctly from that of the cortex and no evidence of pelvic dilation was present. The left kidney measured 7.16 cm in length. The right kidney measured 6.9 cm in length.


Adrenal Glands

The bilateral **adrenal glands** were slightly enlarged in size. No evidence of focal capsular expansion or invasion into the phrenic veins was noted. No overt suspicion of neoplasia was noted. This is considered likely a hyperplastic change associated with stress or adrenal endocrinopathy (PDH). If isosthenuria is persistently present and the patient morphologically suggests Cushing's disease then ACTH testing would be indicated. The left adrenal gland measured 2.7 cm x 1.03 cm width at the caudal pole and 0.73 cm width at the cranial pole. The right adrenal gland measured 2.13 cm x 1.69 cm width at the cranial pole and 0.74 cm width at the caudal pole.

Spleen

The **spleen** presented a smooth homogeneous parenchyma hyperechoic to liver and renal cortical parenchyma. The capsule was smooth without noticeable expansion or deviation from within the spleen or adjacent pathology. The splenic vasculature demonstrated normal volume without signs of congestion or thrombosis. No sonographic evidence of acute or chronic inflammatory, neoplastic, or infarctual changes were noted.

Liver



PATIENT

Chabot Coombs

SPECIES

Canine

BREED

Labrador Retriever Mix

SEX

Neutered Male

AGE

13 Years

WEIGHT

88.5 pounds

INTERPRETED BY

Eric Lindquist, DMV,
DABVP(CFM), Cert.
IVUSS

IMAGING PERFORMED BY

Dr. Chrissy Krell DVM

HOSPITAL NAME

Taylor Emergency
Hospital

REFERRING VET

Dr. Chrissy Krell DVM

INVOICE

13099

DATE

01/12/26

The **liver** images from right and left intercostal as well as subcostal views revealed subjectively normal liver size, contour, and structure. Some minor age-related parenchymal remodeling was noted but likely not clinically significant at this time. Vascular and biliary tracts were of normal volume and no evidence of congestion was noted. The gallbladder presented some dependent debris with essentially normal contour. The cystic and common bile ducts were normal. No overt evidence of active inflammatory, infiltrative or regenerative pathology was noted but should be paired with current or past LE elevations regarding any clinical significance to this presentation. The hepatic lymph nodes were unremarkable.

Gastrointestinal

The **gastrointestinal tract** presented considerable gastric artifact due to the presence of ingesta. This did not permit thorough evaluation of portions of the gastric and upper intestinal structure. No overt abnormality was seen in the visualized tissue, however. This is consistent with a post-prandial presentation within a few hours of mealtime. If the prandial temporal interval does not fit the case history, and the patient presents a history of post-prandial vomiting, this could indicate a delayed upper gastrointestinal outflow due to primary or secondary pyloric hypertrophy, upper GI infiltrative disease, motor deficits, or a non-visualized foreign body. A prudent approach would be to rescan this patient at 24 hour NPO status to further review the non-visible regions if stomach primarily as well as assess any delayed outflow issue.

Pancreas


The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.

ULTRASONOGRAPHIC FINDINGS

- Bilateral adrenal hypertrophy.
- Age-related renal/hepatic changes.
- Chronic cystitis pattern.
- Full stomach.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Differentials for diarrhea include occult parasitism, dietary indiscretion, dietary intolerance, antibiotic responsive colitis, intestinal dysbiosis and occult Addison's should all be considered as causes of diarrhea in this patient. A hydrolyzed diet trial may be in this patient's best interest +/- probiotics. 24-hour NPO and reintroduction of bland diet indicated. I recommend a baseline cortisol or ACTH stimulation test, a fresh fecal smear and fecal floatation analysis if not already performed. Note that recent research has shown that indiscriminate use of antibiotics may actually cause harm. Most acute cases of diarrhea will respond to probiotic therapy, fiber, and gastrointestinal diets over the next 3-5 days.



PATIENT

Chabot Coombs

SPECIES

Canine

BREED

Labrador Retriever Mix

SEX

Neutered Male

AGE

13 Years

WEIGHT

88.5 pounds

INTERPRETED BY

Eric Lindquist, DMV,
DABVP(CFM), Cert.
IVUSS

IMAGING PERFORMED BY

Dr. Chrissy Krell DVM

HOSPITAL NAME

Taylor Emergency
Hospital

REFERRING VET


Dr. Chrissy Krell DVM


INVOICE

13099

DATE

01/12/26





PATIENT

Chabot Coombs

SPECIES

Canine

BREED

Labrador Retriever Mix

SEX

Neutered Male

AGE

13 Years

WEIGHT

88.5 pounds

INTERPRETED BY

Eric Lindquist, DMV,
DABVP(CFM), Cert.
IVUSS

IMAGING PERFORMED BY

Dr. Chrissy Krell DVM

HOSPITAL NAME

Taylor Emergency
Hospital

REFERRING VET

Dr. Chrissy Krell DVM

INVOICE

13099

DATE

01/12/26

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

Eric Lindquist, DMV, DABVP(CFM), Cert. IVUSS,

CEO, Owner, Founder -- SonoPath.com

info@SonoPath.com