



**PATIENT**

Koji Bojourquez

**SPECIES**

Canine

**BREED**

Pit Bull

**SEX**

Spayed Female

**AGE**

13 Years

**WEIGHT**

60 Pounds

**INTERPRETED BY**

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Loetitia-Saint Jacques,  
LVT, RVT

**HOSPITAL NAME**

Pine Creek VC

**REFERRING VET**

Dr. Denny Nolet

**INVOICE**

26554

**DATE**

10/21/21

**PRESENTING CLINICAL SIGNS**

12lb weight loss- follow up  
Abnormal PE/Chem/CBC/UA Results: prev report attached

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The **urinary bladder**, trigone, and pelvic urethra presented normal thicknesses and normal tone. The ureters were not visible which is normal. The amount of debris has significantly diminished. No evidence of inflammatory or neoplastic changes were noted. Ureteral papillae were normal. Significantly improved compared to the prior sonogram.

The **kidneys** revealed largely normal size and structure, corticomedullary definition and ratio (cortex 1/3 of medulla) were essentially maintained with some age-related loss of curvilinear patterns regarding the capsule and C/M junction. The cortices presented largely uniform texture with some increased echogenicity expected for his age patient. Medullary structure differed distinctly from that of the cortex and no evidence of pelvic dilation was present. Slight corticomedullary mineralization noted. The left kidney measured 7.23 cm.

**Adrenal Glands**

The **right adrenal gland** was hypoechoic and mildly enlarged, more heterogeneous than on the prior sonogram with hyperechoic surrounding capsule. The right adrenal gland measured 1.07 cm at the cranial pole and 0.88 cm at the caudal pole. The **left adrenal gland** presented similar changes to the right and measured 0.9 cm at the caudal pole and 0.76 cm at the cranial pole. Slight mineralization noted in both adrenal glands.

**Spleen**

The **spleen** was normal size and relatively normal contour with multifocal hyperechoic areas of mineralization. This is a benign change; however, can be related to Cushing's disease or other endocrinopathies.

**Liver**


The **liver** presented multifocal nodular changes, similar to the prior sonogram with generalized swelling and remodeling and increased portal markings. A portion of the right liver revealed a hyperechoic area with hypoechoic caudal portion of the lobe. The area in question measured approximately 4.0 cm. Other patchy, mixed echogenic changes were noted in the liver with swollen, hypoechoic nodular changes in the caudal aspect of the left lateral lobe as well as the caudate process. These nodular changes appear more pronounced than on the prior sonogram. Ultrasound guided FNA warranted in the various echotextures throughout the liver. Minor gallbladder debris noted.

**Gastrointestinal**

Examination of the **gastrointestinal tract** revealed a stomach and intestine free of stasis, of normal wall thickness, acceptable curvilinear mural detail, and peristaltic activity. Small and large intestine demonstrated normal luminal chyme and stool consistency respectively. No obstructive or overt infiltrative disease was noted. No associated abnormal lymphatic activity was noted.

**Pancreas**

The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Some parenchymal remodeling, however, with mild deviation from curvilinear normalcy was observed.



**PATIENT**

Koji Bojourquez

Pancreatic duct and capsular irregularities were present consistent with age related changes. If pain upon imaging (+ Murphy sign) was present or if the patient is focally painful in subxyphoid palpation then low-grade smoldering chronic pancreatitis should be suspected.

**SPECIES**

Canine

- Bilateral adrenal hypertrophy – increased in size compared to the prior sonogram

**BREED**

Pit Bull

- Pronounced nodular liver changes - Concern for suppurative changes or possible neoplasia, progressive.
- Mineralizing spleen

**SEX**

Spayed Female

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Recommend assessment of urinalysis. If USG is < 1.020, then workup for PDH/Cushing's warranted now that the bladder debris has largely resolved. Blood pressure measurements warranted. FNA of the liver changes strongly recommended and culture of the hepatic aspirates.

**AGE**

13 Years

**Efficient & Accurate Cushing's Work up-Lindquist**

**Notes regarding Cushing's Clinical Presentations:**

**WEIGHT**

60 Pounds

Nearly all Cushing's dogs have SAP elevations and true PU/PD (USG < 1.025) and most are polyphagic. Cushing's dogs are > 6 years and usually > 9 years old, usually have poor skin coats, body scores > 3/5, and are usually sedentary animals.

Its important to remember that Cushing's dogs usually look and play the part and other diseases cause false + stress related cortisol spikes. On rare occasion a Cushing's dog will not follow the rules but this is truly an exception.

**INTERPRETED BY**

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

Potential Cushing's patient workups can be costly and frustrating if not definitive and, in my experience, the non-definitive patient usually has something else going on that may be contributing to some of the clinical signs a Cushing's dog will have, especially SAP elevations or PU/PD. Based on this prelude of information I came up with the following algorithm in the spirit of diagnostic efficiency.

**IMAGING PERFORMED BY**

Loetitia-Saint Jacques,  
LVT, RVT

The following suggested protocol is based on current available literature on Cushing's disease and extensive clinical-sonographic experience evaluation + Cushing's and False + LDDST & ACTH stim. cases in order to maximize the efficiency of a Cushing's workup in practice.

**HOSPITAL NAME**

Pine Creek VC

**Screen first, workup second**

1) **UA:** Repeatable (2-3 urine samples) Urine specific gravity & urine cortisol/creatinine ratio (UCCR): If **repeatable USG < 10.20 and + UCCR** move to next step 2.

**REFERRING VET**

Dr. Denny Nolet

Note: UA is inexpensive and easy to obtain and if UA criteria is not met for Cushing's then resources can be spent into other more pertinent diagnostics or left on hold until the UA criteria is met in emerging Cushing's cases.


**INVOICE**

26554

2) **Sonogram:** Does the patient **have concurrent disease** clinically or sonographically as non-Cushing's illness will influence the potential false + LDDST or even ACTH stim. The sonogram gives a global perspective of the internal health of the patient to be considered in the Cushing's workup as an assessment of concurrent disease. Is there a concurrent neoplastic process, UTI pancreatitis, mucocele....? Are the adrenals enlarged (Cushing's-PDH, stress, age related or breed variant), or atrophied (Iatrogenic Cushing's or adrenal burnout), have asymmetric enlargement (Adrenal tumor, hyperplasia, adenoma, age related variant), or is there vascular invasion (Invasive pheo with false + UA criteria or adenocarcinoma or phrenic thrombosis)? The sonogram answers these questions proactively.

**DATE**

10/21/21



**PATIENT**

Koji Bojourquez

3) **LDDST** (0.01 D-Sodium phosphate mg/kg IV) (Better screening test but plagued with false +) Use if there is potential early Cushing's or if adrenal asymmetry present on sonogram suspecting tumor. Use LDDST in cats at a higher dose (0.1 mg/kg IV).

**SPECIES**

Canine

OR

4) **ACTH stim.** (Better confirming test but can have false +) Use if the patient "looks" Cushingoid or if bilateral adrenal enlargement is present, or high normal width on sonogram, or if iatrogenic Cushing's suspected (Cortisone Tx in past).

**BREED**

Pit Bull

5) If **diabetic** then run both LDDST & ACTH stim.

**SEX**

Spayed Female

5) Run a **serial blood pressure** in a BP friendly non "white coat effect" atmosphere. Run at least 3 at different times over a few hours or when eating as the patient tends to be calm when eating or give Torbutrol when entering the facility.

**AGE**

13 Years


6) **Perform CT** of the pituitary to identify macro adenoma expansion if any lethargy or dullness or other central clinical CNS signs are minimally present.

Suggested reading:

Behrend EN, Kooistra HS, Nelson R, et al. Diagnosis of Spontaneous Canine Hyperadrenocorticism: 2012 ACVIM Consensus Statement (Small Animal). J Vet Intern Med 2013;27:1292-1304.

**WEIGHT**

60 Pounds




**INTERPRETED BY**

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Loetitia-Saint Jacques,  
LVT, RVT



**HOSPITAL NAME**

Pine Creek VC

**REFERRING VET**

Dr. Denny Nolet

**INVOICE**

26554

**DATE**

10/21/21



**PATIENT**

Koji Bojourquez

**SPECIES**

Canine

**BREED**

Pit Bull

**SEX**

Spayed Female

**AGE**

13 Years

**WEIGHT**

60 Pounds

**INTERPRETED BY**

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Loetitia-Saint Jacques,  
LVT, RVT

**HOSPITAL NAME**

Pine Creek VC

**REFERRING VET**


Dr. Denny Nolet

**INVOICE**

26554

**DATE**

10/21/21



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Eric Lindquist**, DMV, DABVP, Cert. IVUSS, CEO of SonoPath.com  
Eric.Lindquist@SonoPath.com