



PATIENT

Bella Saveli

SPECIES

Canine

BREED

Yorshire Terrier

SEX

Spayed Female

AGE

15 Years

WEIGHT

6.7 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Kelly Vazquez

HOSPITAL NAME

Ramsey AC

REFERRING VET

Dr. Bishnoi

INVOICE

17569

DATE

10/3/22

PRESENTING CLINICAL SIGNS

History: Patient presents for wobbly gait, non-ambulatory, falling over. Breathing hard, wheezing in chest. Enlarged and painful abdomen. Treated 3 days ago, routine blood work. Supportive care: Injection Dex SP 0.5 mls SQ, Injection Azium 0.3 mls IM, Injection Azium 0.3 mls IV, Injection Cerenia 0.3 mls SQ, Injection Vitamin 0.5 mls SQ. Nutrical.

Abnormal PE/Chem/CBC/UA Results: BUN 54, BUN/creat. 60, calcium 8.1, magnesium 2.8, potassium 5.7, PrecisionPSL 173. U/A: protein 2+, Urine/Protein creatinine ratio: 2.2.

ULTRASONOGRAPHIC EXAMINATION OF THE HEART & ABDOMEN

CANINE CARDIAC PARAMETERS	MR VMAX (m/s)	TR VMAX (m/s)	LA/AO (Boon method)	LA/AO (Heart Base; Swe)	FS (%)	EF (%)	EPSS (cm)
NORMAL PARAMETER	4.5-5.5	<2.7	1.3	<1.3	28-40	40-100	<0.6
PATIENT		4.0	1.1	--	63	94	NM
CANINE CARDIAC PARAMETERS	HR (BPM)	AV VMAX (m/s)	PV MAX (m/s)	BODY WEIGHT (kg)	LA 2D short axis Base view (cm)	LVIDd Avg; 2D and m-mode short axis (cm)	LVIDs Avg; 2D and m-mode short axis (cm)
NORMAL PARAMETER	50-100	0.7-1.7	0.7-1.6				
PATIENT	--			--	1.6	1.03	--

Cardiac Presentation

The right atrium was severely enlarged with volume contracted left heart. Overwhelming right sided cardiac volume overload present with severe tricuspid insufficiency and pulmonary hypertension. Respiratory artifact was abundant, obscuring some visibility. The echocardiogram was performed rapidly owing to the precarious state of the patient.

Urinary System

The **urinary bladder**, trigone, and pelvic urethra presented normal thicknesses and normal tone. The ureters were not visible which is normal. No uroliths or sediment were visualized and anechoic urine was present. No evidence of inflammatory or neoplastic changes were noted. Ureteral papillae were normal.

The **kidneys** revealed largely normal size and structure, corticomedullary definition and ratio (cortex 1/3 of medulla) were essentially maintained with some moderate age-related loss of curvilinear patterns regarding the capsule and C/M junction. The cortices presented largely uniform texture with some increased echogenicity expected for his age patient. Medullary structure differed distinctly from that of the cortex. Slight pyelectasia was noted (1.48 cm x 0.23 cm in the left kidney). The left kidney measured 3.06 cm.



PATIENT

Bella Saveli

SPECIES

Canine

BREED

Yorshire Terrier

SEX

Spayed Female

AGE

15 Years

WEIGHT

6.7 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

**IMAGING
PERFORMED BY**

Kelly Vazquez

HOSPITAL NAME

Ramsey AC

REFERRING VET

Dr. Bishnoi

INVOICE

17569

DATE

10/3/22

Adrenal Glands

Both **adrenal glands** were visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were unremarkable. Capsule, cortex, and medullary definition were normal for this age patient.

Spleen

The **spleen** presented a smooth homogeneous parenchyma hyperechoic to liver and renal cortical parenchyma. The capsule was smooth without noticeable expansion or deviation from within the spleen or adjacent pathology. The splenic vasculature demonstrated normal volume without signs of congestion or thrombosis. No sonographic evidence of acute or chronic inflammatory, neoplastic, or infarctual changes were noted.

Liver

The **liver** revealed coarse architecture. Increased portal markings were present. The gallbladder and common bile duct were unremarkable. The hepatic veins were dilated, up to 0.7 cm.

Gastrointestinal

Examination of the **gastrointestinal tract** revealed a stomach and intestine free of stasis, of normal wall thickness, acceptable curvilinear mural detail, and peristaltic activity. Small and large intestine demonstrated normal luminal chyme and stool consistency respectively. No obstructive or overt infiltrative disease was noted. No associated abnormal lymphatic activity was noted.

Pancreas

The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.

Free Abdomen


A large amount of **ascites** was noted with passive congestion pattern.

ULTRASONOGRAPHIC FINDINGS

- Geriatric abdomen with passive congestion abdominal pattern and secondary ascites owing to right sided heart failure
- Coarse architecture and increased portal markings in the liver

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Sildenafil is indicated at 1 mg/kg BID for two weeks, if the patient is able to be stabilized, increase to 1.5 mg/kg BID, Ace-Inhibitor at 0.5 mg per kg SID, progressing to BID and Spironolactone at 1-2 mg/kg BID. Prognosis is very poor long term; however, some palliative measures may prove effective.



PATIENT

Bella Savelli

SPECIES

Canine

BREED

Yorshire Terrier

SEX

Spayed Female

AGE

15 Years

WEIGHT

6.7 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Kelly Vazquez

HOSPITAL NAME

Ramsey AC

REFERRING VET


Dr. Bishnoi

INVOICE

17569

DATE

10/3/22





PATIENT

Bella Saveli

SPECIES

Canine

BREED

Yorshire Terrier

SEX

Spayed Female

AGE

15 Years

WEIGHT

6.7 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Kelly Vazquez

HOSPITAL NAME

Ramsey AC

REFERRING VET


Dr. Bishnoi

INVOICE

17569

DATE

10/3/22



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not



PATIENT visible in the image/video clips provided.

Bella Saveli Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

SPECIES

Canine **Eric Lindquist**, DMV, DABVP, Cert. IVUSS, CEO of SonoPath.com
info@SonoPath.com

BREED

Yorshire Terrier

Right Heart Disease-General Considerations

SEX

<http://www.sonopath.com/RightHeartDisease>

Spayed Female

AGE

15 Years

Description: Right heart disease is often an incidental finding, which can be either cardiogenic or secondary to respiratory or systemic disease. The coughing patient with right heart disease may present with primary respiratory disease (i.e., bronchial collapse, collapsing trachea, pneumonitis) and suffer from secondary pulmonary hypertension (PHT). Concurrent mitral valve disease and chronic left-sided congestive heart failure (CHF) might also lead to PHT. The dyspeic patient with right heart enlargement might have pulmonary hypertension due to airway disease, chronic CHF, parenchymal lung disease (e.g. pulmonic fibrosis), or a cardiac shunt with secondary PHT and shunt reversal.

WEIGHT

6.7 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

Primary cardiac causes of right heart enlargement include: tricuspid dysplasia/degeneration; pulmonic stenosis; pulmonic insufficiency; atrial or septal defects; patent ductus arteriosus; right auricular masses; and pericardial peritoneal diaphragmatic hernias. The second most common cause of right-sided enlargement is secondary PHT, which results in high-velocity tricuspid insufficiency (TR vel.>2.8 m/sec) and pulmonic insufficiency due to diseases that cause increased pulmonary vascular resistance or increased pulmonary wedge pressures. The most common cause of secondary PHT is left-sided heart failure (LHF), which presents radiographically as a more globoid-shaped heart with marked left atrial and ventricular enlargement. There are also signs of left-sided CHF as opposed to a simple prominent cranial waist or reverse D radiographic presentation.

IMAGING PERFORMED BY

Kelly Vazquez

HOSPITAL NAME

Ramsey AC

REFERRING VET

Dr. Bishnoi

Secondary, non-cardiac causes of PHT include: acute or chronic respiratory disease; pulmonary thromboembolic disease; thoracic neoplasia; excessive thoracic fat deposition (e.g. Pickwickian syndrome, which leads to chronic hypoxia); brachycephalic syndrome; high altitude disease; heartworm disease; and primary vascular disease.


INVOICE

17569

Clinical Signs: The most common presenting symptoms of right heart disease are collapse, syncope, intermittent or constant acute respiratory distress (e.g. thromboembolic disease), and exercise intolerance.

DATE

10/3/22



PATIENT

Bella Saveli

SPECIES

Canine

BREED

Yorshire Terrier

SEX

Spayed Female

AGE

15 Years

WEIGHT

6.7 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

**IMAGING
PERFORMED BY**

Kelly Vazquez

HOSPITAL NAME

Ramsey AC

REFERRING VET

Dr. Bishnoi

INVOICE

17569

DATE

10/3/22

Diagnostics: Physical examination may reveal a right-sided apical heart murmur and/or a cranial left heart murmur, a split S2, jugular distension, ascites, and signs consistent with respiratory disease (i.e., cough, wheeze, tracheal collapse, tachypnea). Radiographic findings may reveal an enlarged right atrium, right ventricle, and/or primary/secondary branches of the pulmonary artery. In cases of PHT, an enlarged or engorged pulmonary artery is often present. Tortuous arteries or those that suddenly terminate can indicate the presence of thromboembolic disease or heartworms. An interstitial pattern might indicate the presence of pulmonary parasitism or primary interstitial lung disease. Pulmonic stenosis is suspected if the pulmonic segment is enlarged. ECG findings include tall P and S waves with a right axis shift.

Treatment: Please refer to the chapter “Pulmonary Hypertension” for therapeutic recommendations.

References:

Oyama MA, Rush JE, Rozanski EA, et al. Assessment of serum N-terminal pro-B-type natriuretic peptide concentration for differentiation of congestive heart failure from primary respiratory tract disease as the cause of respiratory signs in dogs. *J Am Vet Med Assoc* 2009;235:1319-25.

Rozanski E. Interstitial lung disease in small animals. Proceedings from American College of Veterinary Internal Medicine Forum, Denver, CO, June 15-18, 2011.

Zoia A, Augusto M, Drigo M, Caldin M. Evaluation of hemostatic and fibrinolytic markers in dogs with ascites attributable to right-sided congestive heart failure. *J Am Vet Med Assoc* 2012;241:1336-43.