



## PATIENT

Dolly Manning

## SPECIES

Canine

## BREED

Australian Shepherd

## SEX

Spayed female

## AGE

7 years

## WEIGHT

43.4 lbs

## INTERPRETED BY

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

## IMAGING PERFORMED BY

Dr. Cerf

## HOSPITAL NAME

Veterinary Center of  
Hardyston

## REFERRING VET

Dr. Verf

## INVOICE

75125

## DATE

5/1/26

## PRESENTING CLINICAL SIGNS

History: R/O PLN vs Lyme nephritis vs other

Abnormal PE/Chem/CBC/UA Results: Lyme Positive Monocytes 23 - 10% LOW A/G RATIO 0.40.8 - 2.0 LOW ALBUMIN 2.1 2.7 - 4.4 g/dL LOW GLOBULIN 5.2 1.6 - 3.6 g/dL HIGH ALT (SGPT) 11 12 - 118 IU/L LOW AMYLASE 1178 290 - 1125 IU/L HIGH BUN 33 6 - 31 MG/DL HIGH

## ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

### Urinary System

The **urinary bladder**, trigone, and pelvic urethra presented normal thicknesses and normal tone. The ureters were not visible which is normal. No uroliths or sediment were visualized and anechoic urine was present. No evidence of inflammatory or neoplastic changes was noted. Ureteral papillae were normal.

The **kidneys** revealed non-specific, mild swelling with slight increased cortical echogenicity and remodeling. The left kidney measured 5.25 cm. The right kidney measured 4.7 cm.

### Adrenal Glands

The **left adrenal gland** was visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were unremarkable. Capsule, cortex, and medullary definition were normal for this age patient. The left adrenal gland measured 0.6 cm at the cranial pole.

The **right adrenal gland** was not overtly visualized. However, the region of the right adrenal gland appeared unremarkable.

### Spleen

The **spleen** revealed subtle, minor heterogenous parenchymal changes. There was no evidence of significant pathology.

### Liver

The **liver** images submitted revealed subjectively normal liver size, contour, and structure. Parenchymal echogenicity was naturally coarse and hypoechoic to the spleen. Vascular and biliary tracts were of normal volume with no evidence of congestion. The gallbladder presented acceptably thin walls with primarily anechoic content. The cystic and common bile ducts were normal. No pathological hepatic lymphadenopathy was evident. No overt structural evidence of inflammatory, infiltrative or regenerative pathology was evident.



## PATIENT

Dolly Manning

## SPECIES

Canine

## BREED

Australian Shepherd

## SEX

Spayed female

## AGE

7 years

## WEIGHT

43.4 lbs

## INTERPRETED BY

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

## IMAGING PERFORMED BY

Dr. Cerf

## HOSPITAL NAME

Veterinary Center of  
Hardyston

## REFERRING VET

Dr. Verf

## INVOICE

75125

## DATE

5/1/26

## *Gastrointestinal*

There was some residual chyme and gas was noted in the **stomach**, yet not pathological. This is consistent with post prandial presentation. Dense, acoustic shadowing material was noted and measured up to 1.3 cm, some of which appeared to be ingesta, possible medications. Transit of chyme into the small intestine was normal. Curvilinear patterns were maintained throughout the GI tract. No evidence of pathology. Small and large intestine demonstrated normal luminal chyme and stool consistency respectively. No obstructive or overt infiltrative disease was noted. No associated abnormal lymphatic activity was noted.

## *Pancreas*

The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.

## ULTRASONOGRAPHIC FINDINGS

Shadowing material noted in the stomach.

Non-specific, mild degenerative renal changes.

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The low albumin is likely deriving from protein losing nephropathy depending upon urinalysis results. However, if no significant proteinuria is present then screening for Addison's is indicated and presumably protein losing enteropathy could be occurring, yet there were no structural changes in the GI tract. Doxycycline trial and management for protein losing nephropathy is warranted if significant proteinuria is noted.

Internal medicine consult can be utilized through SonoPath.com. You can select the internal medicine drop down at <http://spa.sonopath.com/>.

One of the world's top internists & SonoPath associate Dr. Remo Lobetti BVSc, MMedVet, PhD, DECVIM can evaluate your case through SonoPath. <https://sonopath.com/resources/sonopath-services/internal-medicine-teleconsultation-services>



**PATIENT**

Dolly Manning

**SPECIES**

Canine

**BREED**

Australian Shepherd

**SEX**

Spayed female

**AGE**

7 years

**WEIGHT**

43.4 lbs

**INTERPRETED BY**

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Dr. Cerf

**HOSPITAL NAME**

Veterinary Center of  
Hardyston

**REFERRING VET**


Dr. Verf

**INVOICE**

75125

**DATE**

5/1/26





## PATIENT

Dolly Manning

## SPECIES

Canine

## BREED

Australian Shepherd

## SEX

Spayed female

## AGE

7 years

## WEIGHT

43.4 lbs

## INTERPRETED BY

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

## IMAGING PERFORMED BY

Dr. Cerf

## HOSPITAL NAME

Veterinary Center of  
Hardyston

## REFERRING VET

Dr. Verf

## INVOICE

75125

## DATE

5/1/26

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Eric Lindquist, DMV, DABVP (CFM), Cert. IVUSS, CEO of SonoPath.com

[info@SonoPath.com](mailto:info@SonoPath.com)

## Protein-Losing Nephropathy (PLN)

<http://www.sonopath.com/PLN>


**Description:** Protein-losing nephropathy (PLN) is a common form of renal disease that typically affects dogs in middle age; it occurs less commonly in cats. Glomerular causes of renal protein loss encompass two broad categories: glomerulonephritis (GN) and amyloidosis. (The causes of GN in human medicine are more specifically differentiated based on a combination of histopathology, immunofluorescence, and electron microscopy findings.) Membranoproliferative glomerulonephritis is the most common cause of GN in dogs and is associated with infectious disease with secondary immune complex deposition as well as Lyme disease. Membranous nephropathy is the second most common cause of GN in dogs and the most common cause in cats. It occurs due to primary immune complex deposition on the urinary side of the basement membrane of the glomerulus, resulting in the leakage of albumin. Amyloidosis is caused by the deposition of amyloid A proteins in a  $\beta$ -pleated sheet configuration in the glomeruli. It is a familial disease in the Shar Pei, but occurs as a reactive disease in other canine breeds. It is also inheritable in the Abyssinian cat, but the amyloidosis occurs in the medulla and is therefore not a protein-losing condition in this breed.

Glomerular lesions can be associated with:

- Infectious diseases:
  - Protozoan: *Babesia*, *Hepatozoon*, and *Leishmania*.
  - Bacterial: *Borrelia*, *Bartonella*, *Brucella*, *Ehrlichia*, *Mycoplasma*, pyometra, pyoderma, endocarditis, and pyelonephritis.
  - Viral: FeLV, FIV, and FIP.
  - Fungal
  - Helminthic: *Dirofilaria*.
- Non-infectious inflammatory diseases: pancreatitis, chronic dermatitis, inflammatory bowel disease, periodontal disease, polyarthritis, and systemic lupus erythematosus (SLE).
- Neoplasia: lymphoma, leukemia, and mast cell disease.
- Familial conditions in the soft-coated Wheaten Terrier, Shar Pei, Beagle, Cocker Spaniel, and Bernese mountain dog.



<b>PATIENT</b>	<ul style="list-style-type: none"> <li>• Idiopathic conditions.</li> </ul>
Dolly Manning	Post-glomerular causes, such as hemorrhage and inflammation, also contribute to urine protein quantification.
<b>SPECIES</b>	
Canine	<i>Proteinuria Classifications:</i> Patients can be divided into three tiers, depending on their clinical characteristics:
<b>BREED</b>	
Australian Shepherd	Tier 1A: persistent subclinical proteinuria
<b>SEX</b>	Tier 1B: persistent proteinuria with hypertension
Spayed female	
<b>AGE</b>	Tier 2A: proteinuria and hypoalbuminemia
7 years	Tier 2B: proteinuria, hypoalbuminemia, and hypertension
<b>WEIGHT</b>	
43.4 lbs	Tier 3A: proteinuria and azotemia
<b>INTERPRETED BY</b>	Tier 3B: proteinuria, azotemia, and hypertension
Eric Lindquist, DMV DABVP, Cert. IVUSS	Tier 3C: proteinuria, azotemia, hypertension, and hypoalbuminemia
<b>IMAGING PERFORMED BY</b>	
Dr. Cerf	<p><b>Diagnostics:</b> Traditionally, urine protein loss has been detected either through a qualitative test, such as a urine dipstick, or with a semi-quantitative test, such as a urine protein-creatinine (UPC) ratio. When the latter is greater than 0.5, it is considered abnormal. False positive results can occur due to contamination of urine with red blood cells, white blood cells, and bacterial protein. Thus, one must use a urine sample with inactive sediment and a negative culture for measurement purposes. A 24-hour urine protein quantification is more accurate but technically more difficult to obtain, as it requires hospitalization and 24-hour urinary catheterization with a closed collection system. Pooling urine samples can be considered in cases where urine protein loss is stable. One must obtain three different urine samples, combine 1 ml from each sample to submit for a UPC test, and ensure that inactive sediments are present in all the samples. There should be a high degree of correlation between the UPC on the pooled sample and the mean of the three samples measured independently. Research has not yet demonstrated the accuracy of pooled samples for urine samples with high protein loss (i.e., in cases where the UPC is &gt; 8).</p>
<b>HOSPITAL NAME</b>	
Veterinary Center of Hardyston	
<b>REFERRING VET</b>	
Dr. Verf	
<b>INVOICE</b>	
75125	
<b>DATE</b>	Further diagnostic tests will depend on the tier classification. Once proteinuria is documented repeatedly, additional tests can be considered to assess for potential underlying causes, and, further to that, possible sources of antigen stimulation. Depending on presentation, tests may include:
5/1/26	



## PATIENT

Dolly Manning

## SPECIES

Canine

## BREED

Australian Shepherd

## SEX

Spayed female

## AGE

7 years

## WEIGHT

43.4 lbs

## INTERPRETED BY

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

## IMAGING PERFORMED BY

Dr. Cerf

## HOSPITAL NAME

Veterinary Center of  
Hardyston

## REFERRING VET

Dr. Verf

## INVOICE

75125

## DATE

5/1/26

- CBC and biochemical profile
- Urine culture and sensitivity
- 4DX
- Blood pressure measurement
- Thoracic and abdominal radiographs
- Spinal radiographs to assess for discospondylitis
- Abdominal ultrasound to assess for evidence of underlying infection or neoplasia
- Echocardiogram to assess for vegetative endocarditis and possible effects of hypertension
- Screen for Cushing's disease, especially if hypertensive (LDDST or ACTH stimulation)
- ANA
- Expanded tick or infectious disease screen
- Renal biopsy to differentiate among specific causes of PLN

Renal biopsy should be considered if proteinuria is severe (UPC > 3.5) and hypoalbuminemia and/or hypertension have been documented. Renal biopsy is an invasive procedure and should be considered only to determine if there is an underlying disease process that would benefit from specific therapy. If the patient is debilitated, severely azotemic, or has uncontrolled hypertension or coagulation abnormalities, then the risk of the procedure and anesthesia may be too great and should not be pursued.

Tissue samples should be submitted for a combination of light microscopy (in formalin; use with special stains), immunofluorescence (in Michel's solution or frozen), and electron microscopy (in formalin with glutaraldehyde). It is imperative to request special media before obtaining the biopsy. Samples can be obtained via ultrasound guidance, laparotomy, or laparoscopy, but cortical samples must be divided so that they can be placed in the three different media. One must ensure that the pre-surgical clotting profile and platelet count are both normal. Patients should undergo pre-biopsy and post-biopsy diuresis.

**Treatment:** The main goals of therapy are to i) reduce proteinuria (i.e., UPC < 1.0); ii) prevent a thrombotic event; iii) manage hypertension; and iv) replace fluid deficits. Fluid therapy should be approached cautiously, especially in patients with nephrotic syndrome. Standard therapy for PLN includes a low-protein diet, which in itself will reduce proteinuria, and the administration of an angiotensin-converting enzyme (ACE) inhibitor, such as enalapril (0.5 mg/kg PO BID) or benazepril (0.5 mg/kg PO Q24hr). Newer proposed therapeutic protocols include increasing the ACE inhibitor dose slowly while monitoring BUN and creatinine carefully. The dose can be raised to 1 mg/kg PO BID if needed, provided creatinine has not increased more than 30% from the baseline level.



## PATIENT

Dolly Manning

## SPECIES

Canine

## BREED

Australian Shepherd

## SEX

Spayed female

## AGE

7 years

## WEIGHT

43.4 lbs

## INTERPRETED BY

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

## IMAGING PERFORMED BY

Dr. Cerf

## HOSPITAL NAME

Veterinary Center of  
Hardyston

## REFERRING VET

Dr. Verf

## INVOICE

75125

## DATE


5/1/26

Another class of drugs currently being used is angiotensin receptor blockers, such as Losartan (the dose in azotemic dogs is 0.125-0.25 mg/kg/day PO Q12-24hr and 0.5-1.0 mg/kg/day in non-azotemic patients). This can be combined with an ACE inhibitor, but it is important to monitor BUN, creatinine, and potassium levels. Spironolactone has been used in people in combination with the other two classes of drugs to further modify the renin-angiotensin-aldosterone system (RAAS) (1-2 mg/kg PO BID); however, the effect of using all three drug classes in dogs has not yet been fully investigated. All of these medications are potassium sparing; thus, monitoring for hyperkalemia is important.

Hypertension is managed with amlodipine (0.1-0.2 mg/kg PO Q12-24hr) when an ACE inhibitor is insufficient to control blood pressure. Supplementing with an anti-thrombotic agent, such as aspirin (1 mg/kg PO Q24hr), may be considered in advanced cases, especially once the patient is hypoalbuminemic. Omega-3 fatty acids can be given (0.25-0.5 g/day), but are typically increased in standard kidney diets.

The most recent controversy in the management of glomerular diseases is the use of immunosuppressive medications. Because it is possible to arrive at a more definitive diagnosis in human patients, the use of immunosuppressive agents can be useful in the management of the disease, specifically when the disease is immune-mediated in its pathogenesis, such as SLE, membranous nephropathy, and minimal change disease glomerulonephritis. The procurement of a renal biopsy is being advocated in dogs so that practitioners can identify the population of patients that may benefit most from immunosuppressive therapy. Presently, there is no evidence-based medicine to suggest that immunosuppressive therapy should definitely be incorporated into a daily protocol for canine patients; however, it could be beneficial in some cases and may even result in remission. Further investigation is warranted. Trials are currently being conducted in patients with Lyme nephritis that are treated with immunosuppressive agents in addition to standard antibiotic therapy. The IRIS Treatment of Canine Glomerular Disease Study Group has suggested the trial use of immunosuppressive therapy in severe, persistent, or progressive PLN, even without a biopsy diagnosis in specific cases that are unresponsive to standard therapy (i.e., nephrotic syndrome, progressively azotemic, hypoalbuminemic patients). One can also consider administering the following drugs: pulse steroid therapy, mycophenolate, cyclophosphamide, azathioprine, and chlorambucil. One should monitor blood work, UPC ratio, and blood pressure weekly for 2 weeks, then biweekly for 6 weeks, then monthly. If there is further deterioration, immunosuppressive therapy should be discontinued.

## References:



<b>PATIENT</b>	Goldstein R and Polzin D. Treatment of canine glomerular disease: report of the IRIS treatment of canine glomerular disease study group. Proceedings from the American College of Veterinary Internal Medicine, Denver, CO, June 15-18, 2011.
Dolly Manning	
<b>SPECIES</b>	
Canine	Grauer GF, Greco DS, Getzy DM, et al. Effects of enalapril versus placebo as a treatment for canine idiopathic glomerulonephritis. <i>J Vet Intern Med</i> 2000;14:526-33.
<b>BREED</b>	
Australian Shepherd	
<b>SEX</b>	Less GE, Cianciolo RE, and Clubb FJ. Renal biopsy and pathologic evaluation of glomerular disease. <i>Top Companion Anim Med</i> 2011;26(3):143-53.
Spayed female	
<b>AGE</b>	LeVine DA, Zhang D, Vaden SL. The use of pooled vs. serial urine samples to measure urine protein:creatinine ratios. <i>Vet Clin Pathol</i> 2010;39(1):53-56.
7 years	
<b>WEIGHT</b>	
43.4 lbs	Nabity MB, Boggess MM, Kashtan CE, et al. Day-to-day variability in the urine protein:creatinine ratio in female dogs with stable glomerular proteinuria caused by x-linked hereditary nephropathy. <i>J Vet Intern Med</i> 2007;21:425-30.
<b>INTERPRETED BY</b>	
Eric Lindquist, DMV DABVP, Cert. IVUSS	Vaden SL. Glomerular Disease. <i>Top Companion Anim Med</i> 2011;26(3):128-34.
<b>IMAGING PERFORMED BY</b>	
Dr. Cerf	Vaden SL. Glomerular Diseases. In: Ettinger SJ and Feldman EC, eds. <i>Textbook of Veterinary Internal Medicine, 7<sup>th</sup> Ed.</i> St Louis, MI: Saunders Elsevier; 2010;2021-36.
<b>HOSPITAL NAME</b>	
Veterinary Center of Hardyston	Vaden SL. Microalbuminuria: What is it and how do I interpret it. Proceedings from the American College of Veterinary Internal Medicine, Charlotte, NC, June 4-7, 2003.
<b>REFERRING VET</b>	
Dr. Verf	
<b>INVOICE</b>	Vaden SL et al. Urinary tract inflammation has a variable effect in urine albumin concentration. <i>J Vet Intern Med</i> 2002;16:378 (abstract).
75125	
<b>DATE</b>	
5/1/26	