



**PATIENT**

Penny Welsh

**SPECIES**

Canine

**BREED**

Toy Poodle

**SEX**

Female

**AGE**

4 Years

**WEIGHT**

2.7 kg

**INTERPRETED BY**

Eric Lindquist, DMV

DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Dr. Laura De Cordon

**HOSPITAL NAME**

Mason Dixon  
Animal Emergency

**REFERRING VET**

Dr. Laura De Cordon

**PRESENTING CLINICAL SIGNS**

waxing/ waning appetite and V+ for almost a week not responded to outpatient treatment radiographs and bloodwork were unremarkable P received last dose of omeprazole and sucralfate on 6/29 in the AM

Abnormal PE/Chem/CBC/UA Results: none

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The **urinary bladder**, trigone, and pelvic urethra presented normal thicknesses and normal tone. The ureters were not visible which is normal. No uroliths or sediment were visualized and anechoic urine was present. No evidence of inflammatory or neoplastic changes were noted. Ureteral papillae were normal.

The **kidneys** revealed normal size and structure, corticomedullary definition and ratio for this age. The cortices presented largely uniform texture with normal echogenic relationship to liver and spleen. Medullary structure differed distinctly from the cortex and no evidence of pelvic dilation was present. The capsules were acceptably uniform without significant irregularities. The right kidney measured 3.18 cm. The left kidney measured 3.17 cm.

**Adrenal Glands**

The **left adrenal gland** was visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were unremarkable. Capsule, cortex, and medullary definition were normal for this age patient. The left adrenal gland measured 0.50 cm.

The region of the **right adrenal gland** was imaged, no evident pathology.

**Spleen**

The **spleen** presented a smooth homogeneous parenchyma hyperechoic to liver and renal cortical parenchyma. The capsule was smooth without noticeable expansion or deviation from within the spleen or adjacent pathology. The splenic vasculature demonstrated normal volume without signs of congestion or thrombosis. No sonographic evidence of acute or chronic inflammatory, neoplastic, or infarctual changes were noted.

**Liver**

The **liver** images submitted revealed subjectively normal liver size, contour, and structure. Parenchymal echogenicity was naturally coarse and hypoechoic to the spleen. Vascular and biliary tracts were of normal volume with no evidence of congestion. The gallbladder presented acceptably thin walls with primarily anechoic content. The cystic and common bile ducts were normal. No pathological hepatic lymphadenopathy was evident. No overt structural evidence of inflammatory, infiltrative or regenerative pathology was evident.

**Gastrointestinal**

The **stomach** revealed mucosal thickening with a minor amount of luminal fluid and trace amount of chyme. Mucosal remodeling noted, most consistent with gastritis. The small intestine and colon were unremarkable. Curvilinear patterns maintained. Minor soft stool noted in the colon.

**INVOICE**

39122

**DATE**

6/30/22



**PATIENT**

**Pancreas**

Penny Welsh

The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.

**SPECIES**

Canine

**PRIMARY FINDINGS**

- Gastritis pattern

**BREED**

Toy Poodle

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

GI protectant protocol warranted +/- endoscopy. A clinical trial of the following may prove effective. No evidence of neoplasia or foreign bodies. Given that the right adrenal gland was not overtly visible, screening for underlying occult Addison's would be indicated.

**SEX**

Female

**Helicobacter/Gastritis protocol**

A clinical trial of **Zithromax** (Dogs: 5-10 mg/kg p.o. q24h. May increase dosing interval to q48h after 3-5 days of treatment), **Metronidazole** (10-20 mg/kg p.o. b.i.d.), **Pepcid** (0.5-1 mg/kg s.i.d.) and **Sucralfate** (0.5-2 g/dog PO) or **Omeprazole** (1 mg/kg p.o. s.i.d.) over the next 3 weeks along with a **novel-protein or hydrolyzed diet** with slurry feeding b.i.d./t.i.d. over the next 2-4 days and then increase to canned diet bid. Dry food should be avoided over the next 4 weeks. A recheck sonogram to assess GI improvement or progression would be ideal in 4 weeks.

**WEIGHT**

2.7 kg

**INTERPRETED BY**

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Dr. Laura De Cordon

**HOSPITAL NAME**

Mason Dixon  
Animal Emergency

**REFERRING VET**



Dr. Laura De Cordon

**INVOICE**

39122

**DATE**

6/30/22





**PATIENT**

Penny Welsh

**SPECIES**

Canine

**BREED**

Toy Poodle

**SEX**

Female

**AGE**

4 Years

**WEIGHT**

2.7 kg



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Eric Lindquist, DMV, DABVP, Cert. IVUSS, CEO of SonoPath.com

[info@SonoPath.com](mailto:info@SonoPath.com)

**INTERPRETED BY**

Eric Lindquist, DMV

DABVP, Cert. IVUSS

**IMAGING PERFORMED BY**

Dr. Laura De Cordon

**HOSPITAL NAME**

Mason Dixon  
Animal Emergency

**REFERRING VET**

Dr. Laura De Cordon

**INVOICE**

39122

**DATE**

6/30/22