



PATIENT PRESENTING CLINICAL SIGNS

Bella Warneking

Was at physical therapy for cruciate rupture, when clinician there thought she looked pot-bellied. Abdominal tap yielded blood, so was send here for workup. Recently had last of her teeth extracted. BAR, normal MM, somewhat pot-bellied. not Painful, no obvious abdominal mass. Stiff joints. 1/6 murmur. Current Medications: carprofen, gabapentin and Bravecto

SPECIES

Canine

BREED

Lab Retriever

Abnormal PE/Chem/CBC/UA Results: 8/3 Labwork (see attached) CBC WNL, Chem Alkp 230, rest WNL

SEX

Spayed Female

Urinary System

The **urinary bladder**, trigone, and pelvic urethra presented normal thicknesses and normal tone. The ureters were not visible which is normal. No uroliths or sediment were visualized and anechoic urine was present. No evidence of inflammatory or neoplastic changes were noted. Ureteral papillae were normal.

AGE

12 Years

The **kidneys** revealed largely normal size and structure, corticomedullary definition and ratio (cortex 1/3 of medulla) were essentially maintained with some age-related loss of curvilinear patterns regarding the capsule and C/M junction. The cortices presented largely uniform texture with some increased echogenicity expected for his age patient. Medullary structure differed distinctly from that of the cortex and no evidence of pelvic dilation was present. Slight anechoic cyst noted in the dorsal cortex of the left kidney. The left kidney measured 7.07 cm. The right kidney measured 7.99 cm.

WEIGHT

78 Pounds

INTERPRETED BY

Eric Lindquist, DMV

Adrenal Glands

DABVP, Cert. IVUSS

The **left adrenal gland** was visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were unremarkable. Capsule, cortex, and medullary definition were normal for this age patient. The left adrenal gland measured 3.58 cm x 0.35 cm at the cranial pole and 0.51 cm at the caudal pole.

IMAGING PERFORMED BY

Amanda Crook – SDEP
Certified Clinical
Sonographer

The **right adrenal gland** was enlarged, measuring approximately 1.8 cm at the cranial pole and 1.58 cm at the caudal pole. Vena cava appeared to be free of evident pathology.

HOSPITAL NAME

Rivers Edge PMC

Spleen

The **spleen** was folded upon itself caudally with minor heterogeneous parenchymal changes. No evidence of significant disease.

REFERRING VET

Dr. Bridget Hayes

Liver

The **liver** was uniformly swollen with minor, excessive gallbladder debris and over distension with dependent and suspended bile without evidence of overt mucocele formation. However, excessive sludge was present. The liver presented coarse architecture with mildly increased portal markings and subtle, mixed echogenic changes. This is consistent with vacuolar hepatopathy and some level of remodeling and history of inflammatory component. There was no overt suspicion of neoplasia.

INVOICE

44656

DATE

8/3/23

Gastrointestinal

There was some residual chyme and gas was noted in the **stomach**, yet not pathological. This is consistent with end post prandial presentation. Transit of chyme into the small intestine was normal.



PATIENT

Bella Warneking

Curvilinear patterns were maintained throughout the GI tract. No evidence of pathology. Small and large intestine demonstrated normal luminal chyme and stool consistency respectively. No obstructive or overt infiltrative disease was noted. No associated abnormal lymphatic activity was noted.

SPECIES

Pancreas

Canine

The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.

BREED

Lab Retriever

Free Abdomen

SEX

Rapid view of the heart revealed no evident pathology.

Spayed Female

ULTRASONOGRAPHIC FINDINGS

AGE

12 Years

- Mildly enlarged right adrenal gland
- Folded, mildly heterogeneous spleen
- Vacuolar hepatopathy liver pattern
- Age related renal changes
- Partially full stomach

WEIGHT

78 Pounds

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

INTERPRETED BY

Eric Lindquist, DMV

Further imaging under sedation of the right adrenal recommended, given its irregular contour, especially if evidence of adrenal disease is present and/or systemic hypertension is an issue. If USG is <1.020, then workup for Cushing's indicated, especially given the history of cruciate rupture.

DABVP, Cert. IVUSS

Efficient & Accurate Cushing's Work up-Lindquist

IMAGING PERFORMED BY

Amanda Crook – SDEP
Certified Clinical
Sonographer

Notes regarding Cushing's Clinical Presentations:

Nearly all Cushing's dogs have SAP elevations and true PU/PD (USG < 1.025) and most are polyphagic. Cushing's dogs are > 6 years and usually > 9 years old, usually have poor skin coats, body scores > 3/5, and are usually sedentary animals.

HOSPITAL NAME

Rivers Edge PMC

Its important to remember that Cushing's dogs usually look and play the part and other diseases cause false + stress related cortisol spikes. On rare occasion a Cushing's dog will not follow the rules but this is truly an exception.

REFERRING VET

Dr. Bridget Hayes

Potential Cushing's patient workups can be costly and frustrating if not definitive and, in my experience, the non-definitive patient usually has something else going on that may be contributing to some of the clinical signs a Cushing's dog will have, especially SAP elevations or PU/PD. Based on this prelude of information I came up with the following algorithm in the spirit of diagnostic efficiency.

INVOICE

44656

The following suggested protocol is based on current available literature on Cushing's disease and extensive clinical-sonographic experience evaluation + Cushing's and False + LDDST & ACTH stim. cases in order to maximize the efficiency of a Cushing's workup in practice.

DATE

8/3/23

Screen first, workup second

1) **UA:** Repeatable (2-3 urine samples) Urine specific gravity & urine cortisol/creatinine ratio (UCCR): If **repeatable USG < 10.20 and + UCCR** move to next step 2.



PATIENT

Bella Warneking

Note: UA is inexpensive and easy to obtain and if UA criteria is not met for Cushing's then resources can be spent into other more pertinent diagnostics or left on hold until the UA criteria is met in emerging Cushing's cases.

SPECIES

Canine

2) **Sonogram:** Does the patient **have concurrent disease** clinically or sonographically as non-Cushing's illness will influence the potential false + LDDST or even ACTH stim. The sonogram gives a global perspective of the internal health of the patient to be considered in the Cushing's workup as an assessment of concurrent disease. Is there a concurrent neoplastic process, UTI pancreatitis, mucocele...? Are the adrenals enlarged (Cushing's-PDH, stress, age related or breed variant), or atrophied (iatrogenic Cushing's or adrenal burnout), have asymmetric enlargement (Adrenal tumor, hyperplasia, adenoma, age related variant), or is there vascular invasion (Invasive pheo with false + UA criteria or adenocarcinoma or phrenic thrombosis)? The sonogram answers these questions proactively.

BREED

Lab Retriever

SEX

Spayed Female

3) **LDDST** (0.01 D-Sodium phosphate mg/kg IV) (Better screening test but plagued with false +) Use if there is potential early Cushing's or if adrenal asymmetry present on sonogram suspecting tumor. Use LDDST in cats at a higher dose (0.1 mg/kg IV).

AGE

12 Years

OR

WEIGHT

78 Pounds

4) **ACTH stim.** (Better confirming test but can have false +) Use if the patient "looks" Cushingoid or if bilateral adrenal enlargement is present, or high normal width on sonogram, or if iatrogenic Cushing's suspected (Cortisone Tx in past).

INTERPRETED BY

Eric Lindquist, DMV

5) If **diabetic** then run both LDDST & ACTH stim.

DABVP, Cert. IVUSS

5) Run a **serial blood pressure** in a BP friendly non "white coat effect" atmosphere. Run at least 3 at different times over a few hours or when eating as the patient tends to be calm when eating or give Torbutrol when entering the facility.

IMAGING PERFORMED BY

Amanda Crook - SDEP
Certified Clinical
Sonographer

6) **Perform CT** of the pituitary to identify macro adenoma expansion if any lethargy or dullness or other central clinical CNS signs are minimally present.

Suggested reading:

HOSPITAL NAME

Rivers Edge PMC

Behrend EN, Kooistra HS, Nelson R, et al. Diagnosis of Spontaneous Canine Hyperadrenocorticism: 2012 ACVIM Consensus Statement (Small Animal). J Vet Intern Med 2013;27:1292-1304.

REFERRING VET


Dr. Bridget Hayes

INVOICE

44656

DATE

8/3/23





PATIENT

Bella Warneking

SPECIES

Canine

BREED

Lab Retriever

SEX

Spayed Female

AGE

12 Years

WEIGHT

78 Pounds

INTERPRETED BY

Eric Lindquist, DMV

DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Amanda Crook - SDEP
Certified Clinical
Sonographer

HOSPITAL NAME

Rivers Edge PMC

REFERRING VET


Dr. Bridget Hayes

INVOICE

44656

DATE

8/3/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Eric Lindquist, DMV, DABVP, Cert. IVUSS, CEO of SonoPath.com

info@SonoPath.com