


PATIENT

One Two Bernabe

PRESENTING CLINICAL SIGNS

 Not eating, vomiting
 Abnormal PE/Chem/CBC/UA Results: ALKP 295, ALT 176, Lymps 0.89, EOS 0.05, MCV 57.0, RETIC-HGB 20.3

SPECIES

Canine

ULTRASONOGRAPHIC EXAMINATION OF THE HEART & ABDOMEN
BREED

French Bulldog

SEX

Neutered Male

AGE

12 Years

WEIGHT

30.9 Pounds

CANINE CARDIAC PARAMETERS	MR VMAX (m/s)	TR VMAX (m/s)	LA/AO (Boon method)	LA/AO (Heart Base; Swe)	FS (%)	EF (%)	EPSS (cm)
NORMAL PARAMETER	4.5-5.5	<2.7	1.3	<1.6	28-40	40-100	<0.6
PATIENT		2.29	NM	1.53	24	48	NM
CANINE CARDIAC PARAMETERS	HR (BPM)	AV VMAX (m/s)	PV MAX (m/s)	BODY WEIGHT (kg)	LA 2D short axis Base view (cm)	LVIDd Avg; 2D and m-mode short axis (cm)	LVIDs Avg; 2D and m-mode short axis (cm)
NORMAL PARAMETER	50-100	0.7-1.7	0.7-1.6				
PATIENT	140	0.96	0.63		3.82	3.65	

INTERPRETED BY

 Eric Lindquist, DMV
 DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

 Summit Dog & Cat
 Hospital

REFERRING VET

Dr. Levitian

INVOICE

25852

DATE

9/29/21

Cardiac Presentation

Mitral insufficiency noted, not clinically significant. Tricuspid insufficiency noted. The heart base revealed an infiltrative mass obstructing the vena cava inflow into the right heart. The echotexture would suggest chemodectoma/aortic body tumor. However, hemangiosarcoma or other neoplasia could not be ruled out. The portal vein was deviated, and the heart base and inflow from the vena cava were obstructed. The mass is not resectable and is extensive, measuring 8.4 cm in its maximum width with irregular infiltrative contour. Tachyarrhythmia noted during the exam.

Urinary System

The **urinary bladder** revealed a minor amount of calculi and slight sand, a grouping of which measured 7.0 mm. The largest calculus measured approximately 3.0 mm. The residual prostate was uniform at 5.0 mm. The pelvic urethra was imaged 3.0 cm beyond the cystourethral junction.

The **kidneys** revealed largely normal size and structure, corticomedullary definition and ratio (cortex 1/3 of medulla) were essentially maintained with some age-related loss of curvilinear patterns regarding the capsule and C/M junction. The cortices presented largely uniform texture with some increased echogenicity expected for his age patient. Medullary structure differed distinctly from that of the cortex and no evidence of pelvic dilation was present. The right kidney measured 5.28 cm. The left kidney measured 5.5 cm.

Adrenal Glands

The **right adrenal gland** was visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were



PATIENT

One Two Bernabe

unremarkable. Capsule, cortex, and medullary definition were normal for this age patient. The right adrenal gland measured 2.58 cm x 0.8 cm at the caudal pole and 0.87 cm at the cranial pole.

SPECIES

Canine

The **left adrenal gland** was heterogeneous and mildly enlarged, measuring 2.54 cm x 0.81 cm at the caudal pole and 0.65 cm at the cranial pole.

Spleen

The **spleen** was slightly enlarged, uniform, with slight scalloping contour.

BREED

French Bulldog

Liver

The **liver** revealed hepatic vein dilation, passive congestion and ascites. The gallbladder was edematous owing to the ascites.

SEX

Neutered Male

Gastrointestinal

Examination of the **gastrointestinal tract** revealed a stomach and intestine free of stasis, of normal wall thickness, acceptable curvilinear mural detail, and peristaltic activity. Small and large intestine demonstrated normal luminal chyme and stool consistency respectively. No obstructive or overt infiltrative disease was noted. No associated abnormal lymphatic activity was noted.

AGE

12 Years

Pancreas

The base and limbs of the **pancreas** were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.

WEIGHT

30.9 Pounds

ULTRASONOGRAPHIC FINDINGS

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

- Obstructive heart base mass – suspect chemodectoma or less likely hemangiosarcoma
- Mitral and tricuspid insufficiency
- Passive congestion liver and secondary ascites owing to thoracic disease
- Minor renal mineralization

IMAGING PERFORMED BY


Jessica Miller

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No evidence of clinically significant primary abdominal disease in this patient. Blood pressures and EKG warranted and palliative care. Prognosis is poor. The patient is at high risk for sudden death.

HOSPITAL NAME

Summit Dog & Cat
Hospital



REFERRING VET

Dr. Levitian

INVOICE

25852

DATE

9/29/21



PATIENT

One Two Bernabe

SPECIES

Canine

BREED

French Bulldog

SEX

Neutered Male

AGE

12 Years

WEIGHT

30.9 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

Summit Dog & Cat
Hospital

REFERRING VET




Dr. Levitian

INVOICE

25852

DATE

9/29/21





PATIENT

One Two Bernabe

SPECIES

Canine

BREED

French Bulldog

SEX

Neutered Male

AGE

12 Years

WEIGHT

30.9 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

Summit Dog & Cat
Hospital

REFERRING VET


Dr. Levitian

INVOICE

25852

DATE

9/29/21





PATIENT


One Two Bernabe

SPECIES

Canine

BREED

French Bulldog



SEX

Neutered Male

AGE

12 Years

WEIGHT

30.9 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

Summit Dog & Cat
Hospital

REFERRING VET

Dr. Levitian

INVOICE

25852

DATE

9/29/21

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.


Eric Lindquist, DMV, DABVP, Cert. IVUSS, CEO of SonoPath.com
info@SonoPath.com

[Pericardial Effusion and Cardiac Neoplasia](#)

<http://www.sonopath.com/CardiacNeoplasiaEffusion>

Description: The pericardium is a fibrous sac that encloses the heart and the great vessels—aorta, pulmonary artery, proximal pulmonary veins, and vena cava—located at the heart’s base. It is attached caudally to the diaphragm and under normal circumstances contains 1-15 mL of fluid. The latter is comprised of phospholipids that lubricate the heart and allow it to expand and contract without generating friction. The pericardium also fixes the heart, prevents excess motion, and links the diastolic distensibility of the ventricles, thus limiting the degree to which either the left or the right ventricle will distend during diastole. When there are acute changes in venous return (i.e., during exercise), the pericardium plays a critical role in limiting ventricular filling. In cases of chronic cardiac enlargement, the pericardium also becomes distended, and its ability to limit ventricular filling, especially when the heart is at rest, becomes compromised. Pericardial tamponade occurs when there is a rapid accumulation of fluid and the pressure inside the pericardium increases significantly. With tamponade, ventricular filling is restricted and cardiac output is decreased. The right atrium and ventricle are the most vulnerable to this condition as these compartments have thinner walls and a lower pressure.

Etiology: Causes of pericardial effusion include:



PATIENT

One Two Bernabe

SPECIES

Canine

BREED

French Bulldog

SEX

Neutered Male

AGE

12 Years

WEIGHT

30.9 Pounds

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

IMAGING PERFORMED BY

Jessica Miller

HOSPITAL NAME

Summit Dog & Cat
Hospital

REFERRING VET

Dr. Levitian

INVOICE

25852

DATE

9/29/21

- Neoplasia
 - Right atrial (RA) hemangiosarcoma
 - Heart base (aortic body) tumors
 - Mesothelioma
 - Rhabdomyosarcoma
 - Ectopic thyroid carcinoma
 - Metastatic neoplasia
- Idiopathic
- Congestive heart failure
- Peritoneal-pericardial diaphragmatic hernia
- Pericardial cyst
- Hypoalbuminemia
- Infectious pericarditis (bacterial, *Coccidioides immitus*)
- Feline infectious peritonitis
- Left atrial tear secondary to valvular disease
- Coagulopathy

The majority of neoplastic masses consist of hemangiosarcoma and heart-based tumors (chemodectomas or ectopic thyroid adenocarcinoma). Idiopathic pericardial effusion is a diagnosis of exclusion; the effusion is typically hemorrhagic. Approximately 50% of dogs will be cured with a single pericardiocentesis, while some dogs will require multiple pericardiocenteses as well as surgery. A peritoneal-pericardial diaphragmatic hernia is a congenital hernia seen in dogs and cats in which the abdominal contents (i.e., liver, small intestine, spleen, stomach) herniate into the pericardial sac. Constrictive pericarditis is an uncommon condition in which a non-distensible, thickened, fibrotic pericardium develops over time.

Clinical Signs: One will observe the following clinical signs, which often present in combination: ascites, lethargy, exercise intolerance, pale mucous membranes, weak pulses, *pulsus paradoxus*, and respiratory distress.

Diagnostics: Survey radiographs will reveal hepatomegaly, cardiomegaly (generalized or sectorial globoid), and small pulmonary vessels. Pulmonary edema is typically not found, although one may discover concurrent pulmonary metastatic disease. An ECG will show electrical alternans or small complexes, but often the changes are very subtle and difficult to detect.

Echocardiography is usually considered the gold standard for diagnosing pericardial effusion. Findings include:

- Anechoic space between the heart and the pericardium.
- Abnormal side-to-side cardiac motion.
- Decreased chamber size (right ventricle [RV] and left ventricle [LV]).
- Presence of a pericardial or cardiac mass.
- Tamponade with early diastolic RA and RV collapse.



PATIENT

One Two Bernabe

Cytology is helpful in the diagnosis of lymphoma, septic pericarditis, and idiopathic effusion, but not in cases of neoplasia.

SPECIES

Canine

According to a study that found troponin I levels to be higher in dogs with neoplastic pericardial effusion, the cardiac troponin I assay can be helpful in the diagnosis hemangiosarcoma.

BREED

French Bulldog

Prognosis:

SEX

Neutered Male

- Cardiac hemangiosarcoma: < 8 months with surgical debulking and chemotherapy.
- Chemodectoma (aortic derived): MST 730 days post pericardectomy.
- Idiopathic: 50% complete resolution post cardiocentesis; curative with pericardectomy, which can be done via thoracotomy, or thoracoscopy, or using a balloon to tear the pericardium.
- Mesothelioma: Poor.
- Restrictive pericarditis: Poor, especially when the pericardium has not been surgical stripped.

AGE

12 Years

WEIGHT

30.9 Pounds

References:

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

Cagle LA, Epstein SE, Owens SD, et al. Diagnostic yield of cytology analysis of pericardial effusion in dogs. *J Vet Int Med* 2014;28:66-71.

IMAGING PERFORMED BY

Jessica Miller

Feigenbaum H. Pericardial disease. In: Feigenbaum H, ed. *Echocardiography*, 5th ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 1994:556-588.

HOSPITAL NAME

Summit Dog & Cat
Hospital

Jackson J, Richter KP, Launer DP. Thoroscopic partial pericardectomy in 13 dogs. *J Vet Int Med* 1999;13:529-33.

REFERRING VET

Dr. Levitian

Johnson MS, Martin M, Binns S. A retrospective study of clinical findings, treatment and outcome in 143 dogs with pericardial effusion. *J Small Anim Prac* 2004;45:546-52.

INVOICE

25852

Kienle RD, Thomas WP. Echocardiography. In: Nyland TG and Mattoon JS, eds. *Small Animal Diagnostic Ultrasound*, 2nd ed. Philadelphia, PA: WB Saunders; 2000:354-423.

DATE

9/29/21



PATIENT

One Two Bernabe

Miller MW, Sisson DD. Pericardial disorders. In: Ettinger SJ and Feldman EC, eds. *Textbook of Veterinary Internal Medicine, 5th ed.* Philadelphia, PA: WB Saunders; 2000:923-36.

SPECIES

Canine

Rajagopalan V, Jesty SA, Craig LE, et al. Comparison of presumptive echocardiographic and definitive diagnoses of cardiac tumors in dogs. *J Vet Int Med* 2013;27:1092-96.

BREED

French Bulldog

Shaw SP, Rozanski EA, Ruhs JE. Cardiac troponins I and T in dogs with pericardial effusion. *J Vet Int Med* 2004;18:322-24.

SEX

Neutered Male

Sidley JA, Atkins CE, Keene BW, et al. Percutaneous balloon pericardiectomy as a treatment for recurrent pericardial effusion in 6 dogs. *J Vet Intern Med* 2002;16:541.

AGE

12 Years

Sisson D, Thomas WP. Pericardial disease and cardiac tumors. In: Fox PR, Sisson D, Moise NS, eds. *Textbook of Canine and Feline Cardiology, 2nd ed.* Philadelphia, PA: WB Saunders; 1999:679-701.

WEIGHT

30.9 Pounds

Sisson D, Thomas WP, Reed J, et al. Intrapericardial cysts in the dog. *J Vet Int Med* 1993;7:364-69.

INTERPRETED BY

Eric Lindquist, DMV
DABVP, Cert. IVUSS

**IMAGING
PERFORMED BY**

Jessica Miller

HOSPITAL NAME

Summit Dog & Cat
Hospital

REFERRING VET

Dr. Levitian

INVOICE

25852

DATE

9/29/21