



## PATIENT

Doodle Bug Common  
Sense for Animals

## SPECIES

Canine

## BREED

Mix

## SEX

Spayed female

## AGE

2 years

## WEIGHT

47.5 lbs

## INTERPRETED BY

Eric Lindquist, DMV  
DABVP, Cert. IVUSS

## HOSPITAL NAME

SDEP Lab

## INVOICE

68694

## DATE

11/14/25

## PRESENTING CLINICAL SIGNS

T 100.4, HR 150, RR 18, BCS4/9, PS 0/4, FS 3-4/5

## ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

### Urinary System

The **urinary bladder**, trigone, and pelvic urethra presented normal thicknesses and normal tone. The ureters were not visible which is normal. No uroliths or sediment were visualized and anechoic urine was present. No evidence of inflammatory or neoplastic changes was noted. Ureteral papillae were normal.

The **kidneys** revealed normal size and structure, corticomedullary definition and ratio for this age. The cortices presented largely uniform texture with normal echogenic relationship to liver and spleen. Medullary structure differed distinctly from the cortex and no evidence of pelvic dilation was present. The capsules were acceptably uniform without significant irregularities.

### Adrenal Glands

Both **adrenal glands** were visualized and recognized as having normal shape, size, position and echogenicity for this breed. The phrenic vasculature, glandular echogenicity and detail were unremarkable. Capsule, cortex, and medullary definition were normal for this age patient.

### Spleen

The **spleen** presented a smooth homogeneous parenchyma hyperechoic to liver and renal cortical parenchyma. The capsule was smooth without noticeable expansion or deviation from within the spleen or adjacent pathology. The splenic vasculature demonstrated normal volume without signs of congestion or thrombosis. No sonographic evidence of acute or chronic inflammatory, neoplastic, or infarctual changes was noted.

### Liver

The **liver** images submitted revealed subjectively normal liver size, contour, and structure. Parenchymal echogenicity was naturally coarse and hypoechoic to the spleen. Vascular and biliary tracts were of normal volume with no evidence of congestion. The gallbladder presented acceptably thin walls with primarily anechoic content. The cystic and common bile ducts were normal. No pathological hepatic lymphadenopathy was evident. No overt structural evidence of inflammatory, infiltrative or regenerative pathology was evident.

### Gastrointestinal

Examination of the **gastrointestinal tract** revealed a stomach and intestine free of stasis, of normal wall thickness, acceptable curvilinear mural detail, and peristaltic activity. Small and large intestine



<b>PATIENT</b>	demonstrated normal luminal chyme and stool consistency respectively. No obstructive or overt infiltrative disease was noted. No associated abnormal lymphatic activity was noted.
Doodle Bug Common Sense for Animals	
<b>SPECIES</b>	<b><i>Pancreas</i></b>
Canine	The base and limbs of the <b>pancreas</b> were observed to be largely isoechoic to surrounding omental fat. Pancreatic duct and capsular contour were acceptably normal and parenchyma respected normal curvilinear patterns. No overt evidence of active inflammatory or neoplastic disease was noted.
<b>BREED</b>	
Mix	
<b>SEX</b>	<b>ULTRASONOGRAPHIC EXAMINATION OF THE HEART</b>
Spayed female	The echocardiogram in this patient demonstrated normal <b>left atrial</b> size based on 3 separate methods of LA evaluation. The cranial and caudal <b>mitral</b> valve leaflets presented normal linear structure, extension in systole, and union in diastole with normal kinesis. The <b>left ventricle</b> presented thicknesses with linear contour and was not dilated nor restricted. The <b>myocardium</b> presented normal echogenicity without subjective evidence of significant fibrotic or ischemic disease. <b>Contractility</b> of the ventricular walls was adequate and in normal range for this patient evidenced by the fractional shortening measurement and subjective evaluation of the different regions of the myocardium. The <b>left ventricular outflow</b> tract demonstrated normal laminar flow and subjective structural integrity. The <b>right atrium</b> and auricle revealed normal size, structure and content. No evidence of masses was noted. <b>Tricuspid</b> valvular assessment demonstrated adequate linear morphology and kinesis. The <b>right ventricle</b> was of normal size (1/3 diameter of LV), chordae structure, myocardial echogenicity and thickness. <b>Pulmonary outflow</b> tract assessment revealed normal valve structure, laminar flow, and diameter (approx. 1:1 pa/ao ratio). No visible <b>pericardial</b> or free pleura fluid was noted. The cranial <b>mediastinum and pericardial and extra-cardiac regions</b> were free of masses in the visible window.
<b>AGE</b>	
2 years	
<b>WEIGHT</b>	
47.5 lbs	
<b>INTERPRETED BY</b>	<b>ULTRASONOGRAPHIC FINDINGS</b>
Eric Lindquist, DMV DABVP, Cert. IVUSS	<ul style="list-style-type: none"> <li>• Normal abdomen.</li> <li>• Normal echocardiogram.</li> </ul>
<b>HOSPITAL NAME</b>	<b>INTERPRETATION OF THE FINDINGS &amp; FURTHER RECOMMENDATIONS</b>
SDEP Lab	No specific therapy is recommended at this time.
<b>INVOICE</b>	<b>The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.</b>
68694	
<b>DATE</b>	Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.
11/14/25	
	<b>Eric Lindquist, DMV, DABVP, Cert. IVUSS, CEO of SonoPath.com</b>
	<a href="mailto:info@SonoPath.com">info@SonoPath.com</a>