

PATIENT PRESENTING CLINICAL SIGNS

Bonzei Obrejanu Recently diagnosed diabetic, possible cushings meds: caninsulin 5 IU BID, aventi liver

SPECIES Abnormal PE/Chem/CBC/UA Results: diabetic, proteinuria, pancreatitis chronic; mild elevation in UPCR
Canine

BREED ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Bichon X *Urinary System*

SEX The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.
Neutered Male

AGE The visualized areas of prostate and surrounding tissue appear normal. Unfortunately, the prostate is not fully visualized likely due to its intrapelvic location. Correlate with rectal exam findings.

5 Years

WEIGHT The left kidney has a normal shape and size (5.61 cm) with occasional small cortical cysts. Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.
20 Pounds

INTERPRETED BY

Kathleen Sennello DVM, MS, Diplomate ACVIM (Small Animal Internal Medicine)

The right kidney has a normal shape and size (5.94 cm) with occasional small cortical cysts. Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

IMAGING PERFORMED BY

Kelly Reschny

The left adrenal gland is normal/borderline “plump”, measuring 0.85 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

HOSPITAL NAME

Aldershot AH

The right adrenal gland is normal/borderline “plump”, measuring 0.77 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

REFERRING VET

Dr. Wallace

Spleen

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

INVOICE

44348


Liver

The liver is large with smooth peripheral margins. The parenchyma is hyperechoic and homogenous in echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

DATE

1/19/23

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. There is a moderate amount of non-organized echogenic debris. The cystic and common bile ducts are normal/not visible.


PATIENT
Gastrointestinal

Bonzei Obrejanu

The stomach is moderately dilated with fluid and irregular shadowing material most consistent with normal ingesta and gas. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layering is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

SPECIES

Canine

BREED

Bichon X

SEX

Neutered Male

AGE

5 Years

WEIGHT

20 Pounds

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measures 0.34 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The left limb of the pancreas is prominent and mottled compared to the surrounding isoechoic mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

INTERPRETED BY

 Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

PRIMARY FINDINGS
**IMAGING
 PERFORMED BY**

Kelly Reschny

- Prominent, mottled left limb of the pancreas – The pancreatic changes are most consistent with mild pancreatitis or a recent episode of pancreatic inflammation.
- Large hyperechoic liver – The diffuse hepatic changes are non-specific and can be seen with vacuolar hepatopathy, reactive change, nodular hyperplasia or, less likely, inflammatory/immune-mediated disease, infiltrative neoplasia, or other hepatopathy. This could be consistent with a diabetic hepatopathy.
- Moderate gallbladder debris – The significance of the aggregated gallbladder debris is unclear. This could represent an early mucocele, cholestasis, or may be secondary to fasting but seems unlikely to be causing a current issue. Recommend continued monitoring.
- Borderline bilateral adrenomegaly – The bilateral adrenomegaly could be consistent with bilateral hyperplasia (e.g., secondary to pituitary-dependent hyperadrenocorticism), bilateral infiltrative neoplasia, inflammatory adrenal disease, other. Correlation with clinical findings is recommended.

HOSPITAL NAME

Aldershot AH

REFERRING VET

Dr. Wallace

INVOICE


44348

DATE

1/19/23

ULTRASONOGRAPHIC FINDINGS

- Moderate shadowing ingesta within the gastric lumen – This is most consistent with a recent meal.



PATIENT

Bonzei Obrejanu

SPECIES

Canine

BREED

Bichon X

SEX

Neutered Male

AGE

5 Years

WEIGHT

20 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

**IMAGING
 PERFORMED BY**

Kelly Reschny

HOSPITAL NAME

Aldershot AH

REFERRING VET

Dr. Wallace

INVOICE

44348


DATE


1/19/23

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Today's scan appears relatively normal for a diabetic patient. The liver is large and hyperechoic, which we often see associated with a diabetic hepatopathy. Additionally, the pancreas is somewhat prominent and mottled. This could be consistent with current mild inflammation or previous episodes of inflammation, and there is some moderate gallbladder debris, which should be monitored.

The adrenals measure at the high end of normal but appear relatively normal otherwise. It is difficult to sometimes differentiate Cushing's from diabetes because the symptoms overlap so significantly. It is smart to try and evaluate for concurrent issues such as urinary tract infections, pancreatitis, obesity, etc., which will complicate management. I would consider gradually increasing the insulin dose and monitoring with glucose curves until insulin resistance is strongly suspected (typically between 9-13 units twice daily). If the response to insulin is still very poor at these higher dosages, then consider an ACTH stimulation test (less affected by non-adrenal illness) to look further into the possibility of Cushing's disease.





PATIENT

Bonzei Obrejanu

SPECIES

Canine

BREED

Bichon X

SEX

Neutered Male

AGE

5 Years

WEIGHT

20 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Kelly Reschny

HOSPITAL NAME

Aldershot AH

REFERRING VET


Dr. Wallace

INVOICE

44348

DATE

1/19/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com