



PATIENT

Sparky Jacqueline

SPECIES

Canine

BREED

Goldendoodle

SEX

Neutered Male

AGE

9 Years 4 Months

WEIGHT

60 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Kerri Becker

HOSPITAL NAME

Bergen County VC

REFERRING VET

Dr. Gess

INVOICE

72535

DATE

1/28/26

PRESENTING CLINICAL SIGNS

Focal area mid abd. thickened on palpation. Hematuria vs. bilirubinuria.

Poss FB H/o eating underwear.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The visualized areas of prostate and surrounding tissue appear normal. Unfortunately, the prostate is not fully visualized likely due to its intrapelvic location. Correlate with rectal exam findings.

The left kidney has a normal shape and size (7.37 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (6.67 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is large, measuring 0.90 cm at the cranial pole and 0.93 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is large, measuring 0.60 cm at the cranial pole and 1.12 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is large and irregular in shape, measuring 2.56 cm in width at the level of the hilus. The blood flow through the hilus and splenic parenchyma appears normal. There is a very large, mixed echogenicity, partially cavitated, hypoechoic mass effect visualized in the cranial abdomen measuring 11.92 cm x 4.83 cm. This appears to be arising from the cranial aspect of the spleen.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.



PATIENT

Sparky Jacquette

SPECIES

Canine

BREED

Goldendoodle

SEX

Neutered Male

AGE

9 Years 4 Months

WEIGHT

60 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Kerri Becker

HOSPITAL NAME

Bergen County VC

REFERRING VET

Dr. Gess

INVOICE

72535

DATE

1/28/26

Gastrointestinal

The stomach contains mild shadowing ingesta. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measures 0.21 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The area of the pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

There is scant free fluid noted. No lymphadenopathy. The omentum is hyperechoic around the large cranial abdominal mass.

Other

The right auricle and pericardium were visualized and were unremarkable. No obvious pathology is visualized. If cardiac function evaluation is desired a full echocardiogram is warranted.

PRIMARY FINDINGS

- Large, mixed echogenicity, partially cavitated, hypoechoic cranial abdominal mass strongly suspected to arise from the spleen – A large, heterogenous mass with cavitations is present within the splenic parenchyma. The mass distorts the splenic capsule. Differentials for the mass include neoplasia (e.g., hemangiosarcoma, hemangioma), hematoma, abscess, other. A neoplastic process is favored.
- Heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.

SECONDARY FINDINGS

- Focal shadowing ingesta visualized within the gastric lumen – Findings could be consistent with ingesta, ingested foreign material, etc.
- Mild bilateral adrenomegaly – The bilateral adrenomegaly could be consistent with bilateral hyperplasia (e.g., secondary to pituitary-dependent hyperadrenocorticism), bilateral infiltrative neoplasia, inflammatory adrenal disease, other. Correlation with clinical findings is recommended.



PATIENT

Sparky Jacquette

SPECIES

Canine

BREED

Goldendoodle

SEX

Neutered Male

AGE

9 Years 4 Months

WEIGHT

60 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

**IMAGING
 PERFORMED BY**

Kerri Becker

HOSPITAL NAME

Bergen County VC

REFERRING VET

Dr. Gess

INVOICE

72535

DATE

1/28/26


INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is a very large, mixed echogenicity, partially cavitated cranial abdominal mass. This overlaps the spleen, and a vascular connection is strongly suspected but cannot be 100% confirmed. Findings are concerning for a neoplastic lesion, but a benign lesion is possible. Recommend splenectomy for both diagnostic and therapeutic purposes.

The liver is somewhat heterogeneous. This could be consistent with a vacuolar hepatopathy or similar. No evidence of a metastatic process is visualized, but it cannot be ruled out. Recommend a biopsy of the liver at the time of surgery.

Both adrenals measure as large on today's exam. The significance of this is uncertain. If signs consistent with Cushing's disease are present, adrenal function testing could be considered in the future once this issue has resolved.

Recommend three view thoracic radiographs to evaluate for possible concurrent thoracic disease/involvement (disregard if this has already been done).





PATIENT

Sparky Jacquette

SPECIES

Canine

BREED

Goldendoodle

SEX

Neutered Male

AGE

9 Years 4 Months

WEIGHT

60 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Kerri Becker

HOSPITAL NAME

Bergen County VC

REFERRING VET


Dr. Gess

INVOICE

72535

DATE

1/28/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com